

12-17-2015

## Knowledge, Attitudes, and Intentions Toward Fertility Awareness and Oocyte Cryopreservation Among Obstetrics and Gynecology Resident Physicians

L. Yu

*Yale University School of Medicine*

Brennan Peterson

*Chapman University, bpeterson@chapman.edu*

M. C. Inhorn

*Yale University*

Julia K. Boehm

*Chapman University, jboehm@chapman.edu*

P. Patrizio

*Yale University School of Medicine*

Follow this and additional works at: [https://digitalcommons.chapman.edu/mft\\_articles](https://digitalcommons.chapman.edu/mft_articles)



Part of the [Family Medicine Commons](#), [Marriage and Family Therapy and Counseling Commons](#), [Medical Education Commons](#), [Obstetrics and Gynecology Commons](#), and the [Women's Health Commons](#)

---

### Recommended Citation

Yu, L., Peterson, B., Inhorn, M. C., Boehm, J. K., & Patrizio, P. (2015). Knowledge, attitudes, and intentions toward fertility awareness and oocyte cryopreservation among obstetrics and gynecology resident physicians. *Human Reproduction*, dev308. <http://doi.org/10.1093/humrep/dev308>

This Article is brought to you for free and open access by the Marriage and Family Therapy at Chapman University Digital Commons. It has been accepted for inclusion in Marriage and Family Therapy Faculty Articles and Research by an authorized administrator of Chapman University Digital Commons. For more information, please contact [laughtin@chapman.edu](mailto:laughtin@chapman.edu).

---

# Knowledge, Attitudes, and Intentions Toward Fertility Awareness and Oocyte Cryopreservation Among Obstetrics and Gynecology Resident Physicians

## Comments

This is a pre-copy-editing, author-produced PDF of an article accepted for publication in *Human Reproduction* following peer review. The definitive publisher-authenticated version is available online at DOI: [10.1093/humrep/dev308](https://doi.org/10.1093/humrep/dev308).

## Copyright

Oxford University Press

1 **Title:**

2 Knowledge, attitudes, and intentions toward fertility awareness and oocyte cryopreservation among  
3 obstetrics and gynecology (OB/GYN) resident physicians

4  
5 **Running Title:**  
6 Gynecologists views about age and oocyte storage

7  
8  
9 **Authors:**

10 L. Yu<sup>1</sup>, B. Peterson<sup>2</sup>, M.C. Inhorn<sup>3</sup>, J.K. Boehm<sup>4</sup>, P. Patrizio<sup>1,5,\*</sup>

11  
12 <sup>1</sup>Yale University School of Medicine, Obstetrics Gynecology and Reproductive Sciences, New Haven CT,  
13 U.S.A. <sup>2</sup>Chapman University, Crean College of Health and Behavioral Sciences, Orange CA, U.S.A. <sup>3</sup>Yale  
14 University, Department of Anthropology, New Haven CT, U.S.A. <sup>4</sup>Chapman University, Department of  
15 Psychology, Crean College of Health and Behavioral Sciences, Orange CA, U.S.A. <sup>5</sup>Yale University, Fertility  
16 Center, New Haven CT, U.S.A. pasquale.patrizio@yale.edu

17  
18  
19  
20  
21  
22  
23 **\*Correspondence address:** Pasquale Patrizio, M.D., M.B.E., Yale University Fertility Center, 150 Sargent  
24 Drive, New Haven, Connecticut 06511, (E-mail: [pasquale.patrizio@yale.edu](mailto:pasquale.patrizio@yale.edu)) Telephone: 1 (203) 785-4708

25  
26

27

28

29

30

31

32

33

34 **Abstract**

35 **Study question:** What knowledge, attitudes and intentions do US obstetrics and gynecology (OB/GYN)  
36 residents have toward discussing age-related fertility decline and oocyte cryopreservation with their  
37 patients?

38  
39 **Summary answer:** Most OB/GYN residents believe that age-related fertility decline, but not oocyte  
40 cryopreservation, should be discussed during well-woman annual exams; furthermore, nearly half of  
41 residents overestimated the age at which female fertility markedly declines.

42  
43 **What is known already:** Oocyte cryopreservation can be utilized to preserve fertility potential. Currently,  
44 no studies of US OB/GYN residents exist that question their knowledge, attitudes, and intentions toward  
45 discussing age-related fertility decline and oocyte cryopreservation with patients.

46  
47 **Study design, size, duration:** A cross-sectional online survey was conducted during the fall of 2014 among  
48 residents in American Council for Graduate (ACOG) Medical Education-approved OB/GYN residency  
49 programs. Program directors were emailed via the ACOG Council on Resident Education in Obstetrics and  
50 Gynecology server listing and asked to solicit resident participation.

51  
52 **Participants/materials, setting, methods:** Participants included 238 residents evenly distributed between  
53 post-graduate years 1-4 with varied post-residency plans; 90% of residents were women and 75% were 26-  
54 30 years old. The survey was divided into three sections: demographics, fertility awareness, and attitudes  
55 toward discussing fertility preservation options with patients. Descriptive and inferential statistics were  
56 conducted.

57  
58 **Main results and the role of chance:** A strong majority of residents (83%) believed an OB/GYN should  
59 initiate discussions about age-related fertility decline with patients (mean patient age 31.8), and 73%  
60 percent believed these discussions should be part of an annual exam. One third of residents overestimated  
61 the age at which there is a slight decline in female fertility, while nearly half of residents overestimated the  
62 age at which female fertility markedly declines. Over three-quarters of residents (78.4%) also  
63 overestimated the likelihood of success using assisted reproductive treatments (ARTs). Residents were  
64 likely to support oocyte cryopreservation in cancer patients irrespective of the woman's age, but much  
65 less likely to support elective oocyte cryopreservation. For elective oocyte cryopreservation, 40% believed  
66 OB/GYNs should initiate discussions with patients (mean age 31.1), while only 20% believed this topic  
67 should be part of an annual exam.

68  
69 **Limitations, reasons for caution:** Because the study invitation was sent through US OB/GYN residency  
70 program directors rather than directly to residents, it is possible that some residents did not receive the  
71 invitation to participate. This limits the generalizability of the findings.

72  
73 **Wider implications of the findings:** Within the US, there appears to be a critical need for improved  
74 education on fertility decline in OB/GYN residency programs. To promote informed reproductive decision-  
75 making among patients, efforts should be made to help OB/GYNs provide comprehensive fertility  
76 education to all women, while also respecting patient choices.

77  
78 **Study funding/competing interest(s):** None

79  
80 **Key words:** fertility awareness, fertility preservation, age-related fertility decline, elective oocyte  
81 cryopreservation, medical oocyte cryopreservation, oocyte storage, ovarian aging, reproductive health  
82 education, obstetrics and gynecology residents, physicians

**83 Introduction**

84 Studies assessing the relationship between female fertility and aging have utilized a variety of  
85 approaches, including observational data on study populations, statistical modeling, and biochemical  
86 assays (Howe et al., 1985; Menken and Larken, 1986; Dunson et al., 2002; Broekmans et al., 2006;  
87 Eijkemans et al., 2014). Although the specified age of onset of fertility decline varies among studies, there  
88 is widespread agreement that female fertility begins to decline by a woman's early 30s, and that the rate  
89 of decline markedly increases at age 37 and thereafter (Howe et al., 1985; Dunson et al., 2002; Te Velde  
90 and Pearson, 2002; ACOG, 2014; ASRM, 2013). However, a large number of international studies have  
91 consistently found that people who are likely to delay childbearing underestimate the impact of age on  
92 fertility as a potential risk factor for involuntary childlessness (Lampic et al., 2006; Tyden et al., 2006;  
93 Bretherick et al., 2010; Hashiloni-Dolev et al., 2011; Virtala et al., 2011; Peterson et al., 2012; Wyndham et  
94 al., 2012; Chan et al., 2015). These studies also found that participants overestimate the effectiveness of  
95 assisted reproductive treatments (ARTs) to overcome age-related infertility (Center for Disease Control &  
96 Prevention, 2014; Ferraretti et al., 2013; Leridon, 2004). It is therefore particularly important that women  
97 of childbearing age have access to accurate information regarding the impact of age on fertility, as well as  
98 the success rates of both ARTs and fertility preservation, so that they are empowered to make informed  
99 reproductive decisions.

100 Most women who want children report that their health care provider is the preferred and most  
101 reliable source of information about reproductive health, rather than other sources such as the media,  
102 peers, and the Internet (Peterson et al., 2012; Wyndham et al., 2012; Hodes-Wertz et al., 2013; Lundsberg  
103 et al., 2014; Azhar et al., 2015). However, women typically wait to seek information from their health care  
104 providers on fertility and conception until they are older, when their fertility may already be declining or  
105 compromised (Lundsberg et al., 2014). Furthermore, not all health care providers are familiar with or

106 comfortable counseling their patients about age-related fertility decline. This combination of patient and  
107 physician factors may lead to a relatively low percentage of patients who actually receive reproductive  
108 health-related information directly from their health care providers (Lundsberg et al., 2014). Primary care  
109 physicians, and to a greater extent obstetric and gynecology (OB/GYN) specialists in hospitals and general  
110 practice, have an important role to play in educating patients about the relationship between age and  
111 fertility. They are also in a position to discuss the implications of oocyte cryopreservation such as cost, risk,  
112 and the estimated number of eggs needed to give women a reasonable chance of having a baby as a result  
113 (Dondorp et al., 2012). For example, doctors can present the possibility of freezing one's eggs for future  
114 use at a time of maximum reproductive potential. Although oocyte cryopreservation is clearly gaining  
115 acceptance for use in patients diagnosed with cancer (Mertes and Pennings, 2011; Noyes et al., 2011),  
116 considerable controversy exists regarding the use of oocyte cryopreservation for non-medical reasons  
117 (Stoop et al., 2011; Stoop et al., 2014). In October 2012, the American Society for Reproductive Medicine  
118 (ASRM) removed the experimental label from oocyte cryopreservation for medical reasons, given that it  
119 has similar obstetric and perinatal outcomes compared to procedures using fresh oocytes (ASRM, 2013;  
120 Oktay et al., 2006; Grifo and Noyes, 2010; Herrero et al., 2011; Rienzi et al., 2012; Levi Setti et al., 2013;  
121 Cobo et al., 2014). However, because of the newness of the procedure, total success rates after long-term  
122 freezing remain unclear making it is difficult to counsel women in either group on the minimum number of  
123 oocytes required to have a reasonable chance of birth after oocyte cryopreservation.

124 To provide a more neutral and accurate description of this technology, we will refer to oocyte  
125 cryopreservation for non-medical reasons as “elective oocyte cryopreservation” (EOC). Although EOC has  
126 the potential to alter the landscape of female fertility decision-making, it is of paramount importance to  
127 assess whether OB/GYNs—considered the first-line providers of comprehensive reproductive health

128 education—feel that they have a responsibility to educate patients about fertility decline and EOC, and  
129 whether they have the necessary education to perform this function. As fertility preservation technologies  
130 become more available in the US, OB/GYN awareness of these technologies may have a major impact on  
131 whether fertility decline is discussed, and if EOC options are presented to women during their routine  
132 gynecologic exams.

133 OB/GYN residents, currently in post-graduate training, may be the most likely physicians to  
134 integrate new evidence-based medicine and technologies into their practice. Thus, this study was designed  
135 to examine the knowledge, attitudes, and intentions of US OB/GYN residents in providing patients with  
136 information on age-related fertility decline and oocyte cryopreservation. Our study aims to assess OB/GYN  
137 residents' knowledge and beliefs regarding age-related fertility decline and the use and availability of  
138 oocyte cryopreservation. It is based on three key research questions: 1) Do OB/GYN residents believe that  
139 it is the role of OB/GYNs to initiate discussions about age-related fertility decline and oocyte  
140 cryopreservation with their patients, and if so, at what ages and how frequently? 2) Do OB/GYN residents  
141 possess accurate knowledge regarding the relationship between female fertility decline and age, as well as  
142 the success rates of ARTs? And 3) Do OB/GYN residents differ in their attitudes toward oocyte  
143 cryopreservation for patients diagnosed with cancer or other medical conditions versus EOC? To our  
144 knowledge, this is the first study to examine these issues in a sample of US OB/GYN resident physicians.

## 145 **Methods**

146 The study used a cross-sectional design to examine the knowledge, attitudes, and intentions of US  
147 OB/GYN residents. The study was reviewed and approved by the Yale University Human Investigation  
148 Committee (HIC#1409014546). All 232 residency program directors listed on the website of the American  
149 College of Obstetrician Gynecologists (ACOG) were sent a hyperlink to an online survey and asked to



150 forward this link to their residents. Resident participants had an opportunity to enter a raffle for one of six  
151 \$50 incentives at the completion of the survey. Between September and October 2014, an initial invitation  
152 email and two reminder emails were sent to each program director with a request to forward the  
153 invitation to all residents.

#### 154 *Instrument Design*

155 The survey was based on existing instruments measuring fertility awareness (Lampic et al., 2006),  
156 and the clinical experiences of the study authors in the fields of obstetrics and gynecology, reproductive  
157 endocrinology, psychology, and anthropology. The instrument was pre-tested on a small group of  
158 graduate students for clarity and wording, and was refined through discussion with this focus group as  
159 well as a literature review examining the published research on fertility awareness, preconception  
160 counseling, and oocyte cryopreservation. The survey included demographic background questions and  
161 questions about residents' attitudes towards discussing age-related fertility decline and oocyte  
162 cryopreservation. Questions took the form of "yes/no" responses (e.g. "Should an OB/GYN initiate  
163 discussions with patients regarding childbearing intentions?"), and open-ended numerical questions (e.g.  
164 "If you answered YES to [the previous question], at what age would you initiate this discussion with  
165 patients?"). Participants were also asked whether these conversations were appropriate for annual well-  
166 woman exams, and data were collected regarding the reasons for or against discussing childbearing  
167 intentions at well-woman exams. Options for open-ended qualitative responses using an 'other' category  
168 were also given to more thoroughly assess residents' attitudes. As in previous studies of fertility  
169 awareness, knowledge-based questions included age of 'slight' versus 'marked' decline in a woman's  
170 ability to become pregnant, and the average success rate for couples undergoing a single round of in vitro  
171 fertilization (IVF) (Lampic et al., 2006; Peterson et al., 2012; Chan et al., 2015). Participants were asked to

172 assess their own familiarity with oocyte cryopreservation, and whether oocyte cryopreservation was  
173 offered at their training institution. They were also asked whether they would initiate discussions of  
174 oocyte cryopreservation with their patients, at what ages they would initiate such discussions, and  
175 whether such discussions should be part of a well-woman annual exam. Finally, residents were asked how  
176 likely they would be to discuss oocyte cryopreservation and support insurance coverage for the technology  
177 in different clinical situations (e.g., “A 25-year-old with cancer”) or for non-medical reasons (e.g., “A 25-  
178 year-old who wants a career first”).

### 179 *Statistical analysis*

180 Data from the online survey were analyzed using SPSS (Version 21). Characteristics of study  
181 participants were first analyzed with descriptive statistics. Next, descriptive analyses regarding issues  
182 related to fertility awareness, pre-conception planning, and oocyte cryopreservation were conducted.  
183 OB/GYN residents’ knowledge of fertility issues was then examined with analyses of variance (ANOVAs),  
184 which tested whether knowledge differed based on participants’ year in residency. Gender differences  
185 were not examined due to the small sample size of male residents.

## 186 **Results**

### 187 *Sample Characteristics*

188 Two hundred thirty nine residents participated in the online survey, or approximately 5% of all  
189 OB/GYN residents in the United States (based on 5,021 total OB/GYN residents reported by the  
190 Accreditation Council for Graduate Medical Education) (ACGME 2014). As shown in Table I, approximately  
191 75% of residents were between the ages of 26 and 30. In addition, a significantly higher percentage of  
192 respondents were women (90.3% versus 81%,  $z = 3.72$ ,  $p < .0001$ ) and white (71.7% versus 54%,  $z = 5.46$ ,  
193  $p < .0001$ ) when compared with the overall population of residents (ACGME, 2014). Respondents were

194 nearly equally split across year of residency and slightly more hailed from the northeastern and southern  
195 regions of the US compared with the western and mid-western regions. Half of respondents (50.2%)  
196 intended to pursue general practice in the future.

### 197 *Attitudes Toward Discussing Pre-Conception Planning and Fertility*

198 Findings regarding residents' tendencies to discuss pre-conception planning and fertility with  
199 patients are shown in Table II. Nearly all respondents (91.7%) indicated that OB/GYNs should initiate  
200 discussion with patients about childbearing intentions at a patient's mean age of 20.8 ( $SD = 5.4$ ). A  
201 majority of respondents also thought that OB/GYNs should initiate discussions about age-related fertility  
202 decline (82.9%), although beginning at a patients' mean age of 31.8 ( $SD = 3.5$ ). Furthermore, 72.4%  
203 thought that discussing age-related fertility decline should be part of a well-woman annual exam with an  
204 OB/GYN, as this would help to educate women about making informed reproductive decisions. Of the  
205 27.6% of residents who did not think that discussing age-related fertility decline should be part of a well-  
206 woman annual exam, 53% explained that an annual exam was too frequent, 53% did not want to be  
207 perceived as pushing childbearing on their patients, and 40% reported that such discussions might lead to  
208 emotional distress in patients.

### 209 *Awareness of Fertility Issues*

210 Residents' knowledge about the ages when female fertility declines and their estimate of chance of  
211 success with IVF are shown in Table III. One third overestimated the age when fertility starts to decline  
212 and almost half of residents (46.5%) overestimated when fertility declines markedly. Estimates of slight  
213 and marked decline in fertility did not differ based on year in residency.

214 Residents also overestimated the overall chance of success in having a child after undergoing one  
215 IVF treatment cycle, as more than three-quarters of residents (78.4%) believed that the success rate was

216 30% or higher. The respondents' mean estimate of overall success was 42.3% ( $SD = 18.4\%$ ; minimum = 5%;  
217 maximum = 80%). Estimates for success after IVF did not differ by residency year.

### 218 *Familiarity with Oocyte Cryopreservation and Attitudes Toward Use*

219 Residents' attitudes towards the use of oocyte cryopreservation are presented in Table IV. Only  
220 one in four residents (25.1%) indicated that they were either "familiar" or "very familiar" with oocyte  
221 cryopreservation. However, six in ten residents (62.6%) worked at a training institution that offered oocyte  
222 cryopreservation to patients, suggesting a lack of education within US institutions' OB/GYN residency  
223 programs about these new technologies.

224 Sixty percent of respondents did not think that OB/GYNs should initiate discussion of oocyte  
225 cryopreservation with their female patients. Of the 40% of respondents who did think that OB/GYNs  
226 should initiate discussion, the mean patient age at which such discussions would occur was 31.1 ( $SD = 4.2$ ).  
227 Even fewer respondents (20.4%) thought that discussion of oocyte cryopreservation should be part of an  
228 annual well-woman exam. Reasons given for discussing oocyte cryopreservation during an annual exam  
229 included educating women to make informed reproductive choices and helping them understand the  
230 implications of oocyte cryopreservation. Reasons against discussing this issue during an annual exam  
231 included wanting to be respectful of patient choices and not wanting to be perceived as pushing  
232 childbearing.

### 233 *Likelihood of Discussing or Supporting Oocyte Cryopreservation for Different Patient Situations*

234 When presented with different patient scenarios, residents showed varying levels of support for  
235 discussing oocyte cryopreservation. As shown in Figure 1, residents were very likely to discuss oocyte  
236 cryopreservation with patients who had received a cancer diagnosis, regardless of whether that patient  
237 was 25 or 35 years of age. In contrast, residents were either somewhat or very unlikely to discuss EOC with

238 patients who wanted to pursue a career before starting a family, especially for younger patients. Similar  
239 patterns were evident for residents' support of insurance coverage of oocyte cryopreservation, with many  
240 more residents likely to support insurance coverage for patients who had received a cancer diagnosis  
241 versus those who wished to pursue a career prior to starting a family (data not shown).

## 242 **Discussion**

243 To our knowledge, this is the first study to examine the attitudes, knowledge, and intentions  
244 regarding fertility awareness and oocyte cryopreservation among US OB/GYN residents. Nearly all  
245 residents (92%) who completed the survey believed that an OB/GYN should initiate discussions regarding  
246 their patients' childbearing intentions. Furthermore, 83% of residents said that they believed an OB/GYN  
247 should initiate discussions about age-related fertility decline with female patients, and 72% said that these  
248 discussions should be a part of an annual well-woman exam. These findings are encouraging, as numerous  
249 international studies have shown that women who are likely to delay childbearing also lack awareness of  
250 age-related fertility decline. A recent study sampling fertility patients from 79 countries found that  
251 accurate fertility knowledge was reported by only 56.9% of patients, supporting the need for more  
252 education (Bunting et al., 2013). Although physicians and other health care providers should be the first-  
253 line reproductive health educators for women (Peterson et al., 2012; Wyndham et al., 2012; Hodes-Wertz  
254 et al., 2013), studies show that many women report never having discussed the effect of age on their  
255 ability to conceive, even though they identify their health care providers as their top source of information  
256 on fertility and reproductive health (Lundsberg et al., 2014).

257 Although it is encouraging that the majority of OB/GYN residents believe that physicians should  
258 initiate discussions about fertility decline with their patients, a surprisingly large percentage of the  
259 residents surveyed were misinformed about fertility decline themselves. For example, 33% of residents

260 believed female fertility slightly declines at age 35 or after, and nearly half (46.5%) of residents indicated  
261 that fertility declines markedly at age 40 or after—when, in fact, the marked decline occurs on average  
262 around the age of 37 (Dunson et al., 2002; ASRM, 2013). Given that OB/GYNs are the gatekeepers of the  
263 dissemination of correct reproductive knowledge, it is concerning that nearly half of the residents in this  
264 study were so uninformed about these basic reproductive facts. Furthermore, given that prior studies  
265 suggest that provision of fertility information impacts patient knowledge and intentions toward delaying  
266 childbearing (Williamson et al., 2014), these findings highlight a critical need for improved education and  
267 curricular offerings on age-related fertility decline in OB/GYN residency programs in the US.

268 In addition to misconceptions about age-related infertility, residents in this study were also  
269 misinformed about the success rates of ARTs. Over three-quarters of residents (78.4%) overestimated the  
270 likely success of IVF in treating infertility. OB/GYN residents seem to share the common misconceptions—  
271 perpetuated by inaccurate media reports, especially of “celebrity moms”—that women can delay having  
272 children until after 40, and that any difficulties can be overcome through IVF (Wyndham et al., 2012). It is  
273 important to educate practitioners that ARTs such as IVF can only make up for half of the births lost by  
274 postponing a first attempt to conceive from age 30 to 35, so that they may correct any misperceptions  
275 that patients may have (Leridon, 2004; Wyndham et al., 2012; ASRM, 2013). ART success rates are directly  
276 related to the age of the patient. For example, women under 35 in the United States have a 41.5% chance  
277 for a live birth using IVF. However, for older women - who may have intentionally postponed childbearing  
278 under the false impression that ARTs could correct any difficulties with fertility - only 11.7% of women  
279 aged 41-42, and only 4.5% of women ages 43-44 had a live birth (CDC, 2014). In other words, women who  
280 use ART in their 40s are much less likely than younger women to have a live birth as a result.

281 When OB/GYN residents in this study were asked whether they should initiate discussions

282 regarding oocyte cryopreservation with patients, less than half (40%) believed that OB/GYNs should, and  
283 only one-fifth (20%) reported that it should be a part of an annual well-woman exam. In a 2013 study of  
284 183 women who had undergone at least one oocyte cryopreservation cycle, the mean age of patients who  
285 cryopreserved their oocytes was 38, an age at which oocytes already have reduced quality and  
286 reproductive potential (Hodes-Wertz et al., 2013). Furthermore, 79% of the women wished they had  
287 undergone EOC at an earlier age, and only one-third had discussed EOC with their gynecologist prior to the  
288 procedure. In the current study, residents believed OB/GYNs should initiate discussions about EOC with  
289 patients starting at age 31, an age when a woman's reproductive potential is greater than the current  
290 norm among actual EOC users, who are on average freezing their oocytes in their late 30s. Recent data  
291 from decision-analysis models propose that the highest probability of achieving a live birth may be when  
292 women undertake EOC at <34 years of age (Mesen et al., 2015). Also, cost-effectiveness studies show that  
293 freezing oocytes by age 35 in women who plan to delay childbearing until age 40 effectively reduces the  
294 cost per live birth (Devine et al. 2015).

295 In considering the role of physicians in discussing these issues with patients, it is important to note  
296 that childbearing decisions are also influenced by relational circumstances or other factors that are  
297 beyond a patient's immediate control. For example, in a study of women who underwent EOC to preserve  
298 their fertility, 161 (88%) had delayed childbearing because they lacked a partner (Hodes-Wertz et al.,  
299 2013). OB/GYNs should be sensitive to these possibilities while delivering information about fertility  
300 decline and EOC. However, EOC may offer some women relief from the pressure of entering into an  
301 unwanted relationship "for the sake of children," or to have children before they are ready. Counseling  
302 women about their fertility and the possibility of EOC requires both maximal sensitivity and respect for  
303 patients' reproductive autonomy. Yet, ideally, OB/GYNs should be initiating such discussions with their

304 patients at an age when patients' reproductive potential can be maximized (the late 20s to early 30s) and  
305 when women may have the greatest flexibility in reproductive decision making.

306 We acknowledge an absence of studies that examine how women may respond to such  
307 discussions. However, we support the conclusions of other studies that call for research to investigate if  
308 patients want physicians to initiate these types of discussions, and under what circumstances they would  
309 like them to take place (Buske et al., 2015). The results of such research would be useful for OB/GYN  
310 training programs around the world, in order to teach residents how to deliver information about age-  
311 related fertility decline and oocyte cryopreservation in a way that respects patient circumstances, while  
312 providing education required for informed decision-making.

313 Examining the attitudes and knowledge of physicians regarding fertility preservation is critical, and  
314 this need has been highlighted in several international studies. For example, in Germany, a survey of 120  
315 oncologists found that while nearly all of the physicians felt fertility preservation was an important issue,  
316 only half reported having a thorough understanding of it, and only 40% reported discussing it with patients  
317 routinely (Buske et al., 2015). A recent study of breast cancer specialists in Japan found that physicians  
318 who had more positive attitudes toward fertility preservation were more likely to discuss this with  
319 patients, and calls were made to improve interdisciplinary communication between physicians and  
320 infertility specialists to improve patient care (Shimizu et al., 2011). Countries around the world are also  
321 beginning to incorporate EOC into standard fertility care, with some nations considering oocyte  
322 cryopreservation to be cost effective and thus potentially covered by insurance or national health plans  
323 (Shkedi-Rafid et al., 2011; Van Loendersloot et al., 2011). In the current study, residents suggested that  
324 they would be more likely to support insurance coverage for oocyte cryopreservation in cancer patients  
325 than for age-matched patients seeking EOC. Thus, future studies must examine how financial coverage of



326 these technologies might impact attitudes toward and uses of both medical oocyte cryopreservation and  
327 EOC in countries throughout the world.

328 Finally, it is important to note that women do not typically make their reproductive decisions  
329 alone, and often include male partners. Men have also been found to significantly overestimate the ages  
330 at which female fertility declines (Peterson et al., 2012). Furthermore, some data have shown that  
331 women's desire for childbearing may be related in part to whether a male partner desires children  
332 (Holton, et al., 2011). Given that reproductive health and the impact of fertility treatments have  
333 increasingly been conceptualized as a couple's issue (Peterson et al., 2009; Peterson et al., 2011), it is  
334 important that providers of men's health care also be encouraged to seek appropriate education regarding  
335 age-related fertility decline. As noted in a recent review, men are often the 'forgotten partner' when  
336 couples are diagnosed with infertility – even in cases of male-factor infertility (Petok, 2015). Once a couple  
337 is diagnosed with infertility, both partners participate in the help seeking process (Johnson & Johnson,  
338 2009). Thus, leaving men out of reproductive counseling overlooks the significant contribution that men  
339 make to reproduction (Inhorn et al., 2009). Discussions of fertility decline and oocyte cryopreservation can  
340 include men as well as women, so that both individuals and couples are better informed about their full  
341 range of reproductive options (Azhar et al., 2015).

### 342 **Limitations**

343 The findings of this study must also be interpreted in light of the study's limitations. First, due to  
344 our sampling methods, we were unable to calculate the exact number of residents who actually received a  
345 recruitment email. Because residents completed the survey in an online, anonymous questionnaire  
346 format, we must take into account the possibility for bias in the study findings. Inherent to email-based  
347 sampling is self-selection bias; we were unable to evaluate characteristics of non-respondents and thus

348 cannot be sure that our findings may therefore not be generalizable to all residents. In addition, female  
349 residents were slightly over-represented in our sample when compared to the proportion of female  
350 OB/GYN residents in the US (AAMC, 2014; ACGME, 2014). The study also did not address residents'  
351 attitudes and intentions regarding the education of men about fertility decline and oocyte  
352 cryopreservation. Finally, due to the small number of male respondents, analyses lacked sufficient power  
353 to explore gender differences among the study responses.

### 354 **Conclusion**

355 This study is the first of its kind to examine the knowledge, attitudes, and intentions of US OB/GYN  
356 residents in providing patients with information on age-related fertility decline and oocyte  
357 cryopreservation. The findings highlight a critical need for improved education among US OB/GYN  
358 residents about issues related to age-related fertility decline and the use of oocyte cryopreservation for  
359 both medical and elective reasons. Although our study focused on OB/GYN residents in the US, topics of  
360 reproductive health and fertility are universal; thus, further research is needed to explore the role of  
361 education across cultures and in countries where OB/GYN training programs may differ in duration and  
362 method. If OB/GYNs are taught to present fertility decline and oocyte cryopreservation to patients in a  
363 way that is both respectful of individual patient autonomy and informative about new reproductive  
364 technologies, then they will maximize their patients' ability to make the most informed reproductive  
365 decisions possible.

### 367 **Acknowledgments:**

368 Special thanks to Ashley Hodgson for her help in preparing the manuscript.

370 **Authors' roles:**

371 P. Patrizio proposed the study, design and reviewed and analyzed data; L. Yu and B. Peterson designed the  
372 study instrument, implemented the study, analyzed data, and prepared the initial manuscript; J. Boehm  
373 conducted data analysis and statistics and assisted with manuscript preparation; M. Inhorn edited the  
374 manuscript and further enhanced data analysis.

375

376 **Funding:**

377 None

378 **Conflict of interest:**

379 None

## References

- 380
- 381 *AAMC Physician Specialty Databook*. 2014. Association of American Medical Colleges (AAMC), Washington  
 382 D.C.
- 383 ACOG. Female age-related fertility decline. Committee Opinion No. 589. *Obstetrics and Gynecology* 2014;  
 384 123:719-21.
- 385 American Society for Reproductive Medicine (ASRM). Practice Committees of the American Society for  
 386 Reproductive Medicine and Society for Assisted Reproductive Technology: mature oocyte  
 387 cryopreservation: a guideline. *Fertil Steril* 2013;**99**:37-43.
- 388 Azhar E, Seifer DB, Malzer K, Ahmed A, Weedon J, Minkiff H. Knowledge of ovarian reserve and  
 389 reproductive choices. *J Assist Reprod Genet* 2015;**32**:409-15.
- 390 Bretherick KL, Fairbrother N, Avila L, Harbord SH, Robinson WP. Fertility and aging: do reproductive-aged  
 391 Canadian women know what they need to know? *Fertil Steril* 2010;**93**:2162-68.
- 392 Broekmans FJ, Kwee J, Hendriks DJ, Mol BW, Lambalk CB. A systematic review of tests predicting ovarian  
 393 reserve and IVF outcome. *Hum Reprod Update* 2006;**12**:685-718.
- 394 Bunting L, Tsibulsky I, Boivin J. Fertility knowledge and beliefs about fertility treatment: findings from the  
 395 international fertility decision-making study. *Hum Reprod* 2013;**28**:385-97.
- 396 Buske D, Sender A, Richter D. Patient-physician communication and knowledge regarding fertility issues  
 397 from German oncologists' perspective-a quantitative survey. *J Canc Educ* 2015;**1**:1-8.
- 398 Centers for Disease Control and Prevention, American Society for Reproductive Medicine, Society for  
 399 Assisted Reproductive Technology. *2012 Assisted Reproductive Technology Fertility Clinic Success Rates*  
 400 *Report*. 2014. US Department of Health and Human Services, Atlanta, GA.
- 401 Chan CH, Chan TH, Peterson BD, Lampic C, Tam MY. Intentions and attitudes towards parenthood and

- 402 fertility awareness among Chinese university students in Hong Kong: a comparison with Western  
 403 samples. *Hum Reprod* 2015;**30**:364-72.
- 404 Cobo A, Serra V, Garrido N, Olmo I, Pellicer A, Remohi J. Obstetric and perinatal outcome of babies born  
 405 from vitrified oocytes. *Fertil Steril* 2014;**102**:1006-15.
- 406 Devine, K, Mumford SL, Goldman KN, Hodes-Wertz B, Druckenmiller S, Propst AM, Noyes N. Baby  
 407 budgeting: oocyte cryopreservation in women delaying reproduction can reduce cost per live birth.  
 408 *Fertil Steril* 2015;**103**:1446-53.
- 409 Dondorp W, de Wert G, Pennings G, Shenfield F, Devroey P, Tarlatzis B, Barri P, Diedrich K. ESHRE Task  
 410 Force on Ethics and Law: oocyte cryopreservation for age-related fertility loss. *Hum Reprod*  
 411 2012;**27**:1231-7.
- 412 Dunson D, Colombo B, Baird D. Changes with age in the level and duration of fertility in the menstrual cycle.  
 413 *Hum Reprod* 2002;**17**:1399-403.
- 414 Eijkemans MJC, Van Poppel F, Habbema DF, Smith KR, Leridon H, Te Velde ER. Too old to have children?  
 415 Lessons from natural fertility populations. *Hum Reprod* 2014;**29**:1304-12.
- 416 Ferraretti AP, Goosens V, Kupka M, Bhattacharya S, de Mouzon J, Castilla JA, Erb K, Korsak V, Andersen AN,  
 417 EIM, ESHRE. Assisted reproductive technology in Europe, 2009: results generated from European  
 418 registers by ESHRE. *Hum Reprod* 2013;**28**:2318-31.
- 419 Gibreel AG, Eladawi N, El-Gilany A-H, Allakkany N, Shams M. How do Egyptian gynecologists manage  
 420 infertility? Cross-sectional study. *Journal of Gynecological Research* 2015;**41**,1067-1073.
- 421 *Graduate Medical Education Data Resource Book*. 2014. Accreditation Council for Graduate Medical  
 422 Education (ACGME), Chicago, IL.
- 423 Grifo JA, Noyes N. Delivery rate using cryopreserved oocytes is comparable to conventional in vitro

- 424 fertilization using fresh oocytes: potential fertility preservation for female cancer patients. *Fertil Steril*  
425 2010;**93**:391-6.
- 426 Hashiloni-Dolev Y, Kaplan A, Shkedi-Rafid S. The fertility myth: Israeli students' knowledge regarding age-  
427 related fertility decline and late pregnancies in an era of assisted reproduction technology. *Hum Reprod*  
428 2011;**26**:3045-53.
- 429 Herrero L, Martinez M, Garcia-Velasco JA. Current status of human oocyte and embryo cryopreservation.  
430 *Obstet Gynecol* 2011;**23**:245-50.
- 431 Hodes-Wertz B, Druckenmiller S, Smith M, Noyes N. What do reproductive-age women who undergo  
432 oocyte cryopreservation think about the process as a means to preserve fertility? *Fertil Steril*  
433 2013;**100**:1343-9.
- 434 Holton, S, Rowe H, Fisher J. Women's health and their childbearing expectations and outcomes: a  
435 population-based survey from Victoria, Australia. *Women Health Iss* 2011;**21**:366-73.
- 436 Howe GWC, Vessey M, Yeates D. Effects of age, cigarette smoking, and other factors on fertility: findings in  
437 a large prospective study. *BMJ* 1985;**290**:1697-1700.
- 438 Inhorn MC, Tjornhoj-Thomsen T, Goldberg H, Mosegaard ML. Reconceiving the second sex, men,  
439 masculinity and reproduction. New York: Berghahn; 2009:2.
- 440 Johnson KM, Johnson DR. Partnered decisions? U.S. couples and medical help-seeking for infertility. *Fam*  
441 *Relat* 2009;**58**:431-44.
- 442 Lampic C, Svanberg AS, Karlstrom P, Tyden T. Fertility awareness, intentions concerning childbearing, and  
443 attitudes towards parenthood among female and male academics. *Hum Reprod* 2006;**21**:558-64.
- 444 Leridon, H. Can assisted reproduction technology compensate for the natural decline in fertility with age? A  
445 model assessment. *Hum Reprod* 2004;**19**:1548-1553.

- 446 Levi Setti PE, Albani E, Morengi E, Morreale G, Piane LD, Scaravelli G, Patrizio P. Comparative analysis of  
447 fetal and neonatal outcome of pregnancies from fresh and cryopreserved/thawed oocytes in the same  
448 group of patients. *Fertil Steril* 2013;**100**:396-401.
- 449 Lundsberg LS, Pal L, Gariepy AM, Xu X, Chu MC, Illuzzi JL. Knowledge, attitudes, and practices regarding  
450 conception and fertility: a population-based survey among reproductive-age United States women.  
451 *Fertil Steril* 2014;**101**:767-74.
- 452 Menken JTJ, Larken, U. Age and infertility. *Sci* 1986;**233**:1389-94.
- 453 Mertes H, Pennings G. Social egg freezing: for better, not for worse. *Reprod Biomed Online* 2011;**23**:824-9.
- 454 Mesen TB, Mersereau JE, Kane JB, Steiner AZ. Optimal timing for elective egg freezing. *Fertil Steril*  
455 2015;**103**:1551-6.
- 456 Noyes N, Knopman JM, Melzer K, Fino ME, Friedman B, Westphal LM. Oocyte cryopreservation as a fertility  
457 preservation measure for cancer patients. *Reprod Biomed Online* 2011;**23**:323-33.
- 458 Oktay K, Cil AP, Bang H. Efficiency of oocyte cryopreservation: a meta-analysis. *Fertil Steril* 2006;**86**:70-80.
- 459 Peterson BD, Pirritano M, Block J, Schmidt L. Marital benefit and coping strategies in men and women  
460 undergoing unsuccessful fertility treatments over a 5-year period. *Fertil Steril* 2011;**95**:1759-63.
- 461 Peterson BD, Pirritano M, Christensen, U, Boivin J, Block J, Schmidt L. The longitudinal impact of partner  
462 coping in couples following five years of unsuccessful fertility treatments. *Hum Reprod* 2009;**24**:1656-  
463 64.
- 464 Peterson BD, Pirritano M, Tucker L, Lampic C. Fertility awareness and parenting attitudes among American  
465 male and female undergraduate university students. *Hum Reprod* 2012;**27**:1375-82.
- 466 Petok WD. Infertility counseling (or the lack thereof) of the forgotten male partner. *Fertil Steril*  
467 2015;**104**:260-6.

- 468 Rienzi L, Cobo A, Paffoni A, Scarduelli C, Capalbo A, Vajta G, Remohi J, Ragni G, Ubaldi FM. Consistent and  
469 predictable delivery rates after oocyte vitrification: an observational longitudinal cohort multicentric  
470 study. *Hum Reprod* 2012;**27**:1606-12.
- 471 Shimizu C, Bando H, Kato T, Mizota Y, Yamamoto S, Fujiwara Y. Physicians' knowledge, attitude, and  
472 behavior regarding fertility issues for young breast cancer patients: a national survey for breast care  
473 specialists. *Breast Cancer* 2013;**20**:230-40.
- 474 Shkedi-Rafid S, Hashiloni-Dolev Y. Egg freezing for age-related fertility decline: preventive medicine or a  
475 further medicalization of reproduction? Analyzing the new Israeli policy. *Fertility and Sterility* 2011;  
476 96:291-4.
- 477 Stoop D, Nekkebroeck J, Devroey P. A survey on the intentions and attitudes towards oocyte  
478 cryopreservation for non-medical reasons among women of reproductive age. *Hum Reprod*  
479 2011;**26**:655-61.
- 480 Stoop D, Van der Veen F, Deneyer M, Nekkebroeck J, Tournaye H. Oocyte banking for anticipated gamete  
481 exhaustion (AGE) is a preventive intervention, neither social nor nonmedical. *Reprod Biomed Online*  
482 2014;**28**:548-51.
- 483 Te Velde E, Pearson PL. The variability of female reproductive ageing. *Hum Reprod Update* 2002;**8**:141-54.
- 484 Tyden T, Svanberg AS, Karlstrom PO, Lihoff L, Lampic C. Female university students' attitudes to future  
485 motherhood and their understanding about fertility. *Eur J Contracept Reprod Health Care* 2006;**11**:181-  
486 9.
- 487 Van Loendersloot LL, Moolenaar LM, Mol BW, Repping S, van der Veen F, Goddijn M. Expanding  
488 reproductive lifespan: a cost-effectiveness study on oocyte freezing. *Human Reproduction* 2011;  
489 26:3054-60.



- 490 Virtala A, Vilska S, Huttunen T, Kunttu K. Childbearing, the desire to have children, and awareness about the  
491 impact of age on female fertility among Finnish university students. *Eur J Contracept Reprod Health*  
492 *Care*, 2011;**16**:108-15.
- 493 Williamson LEA, Lawson KL, Downe PJ, Pierson RA. Informed reproductive decision-making: the impact of  
494 providing fertility information on fertility knowledge and intentions to delay childbearing. *J Obstet*  
495 *Gynecol Canada* 2014;**36**:400-5.
- 496 Wyndham N, Marin Figueira PG, Patrizio P. A persistent misperception: assisted reproductive technology  
497 can reverse the "aged biological clock". *Fertil Steril* 2012;**97**:1044-7.
- 498
- 499




Figure 1: Likelihood of discussing medically indicated and elective oocyte cryopreservation by age

500

Table 1. Characteristics of the Sample

Characteristic	<i>N</i>	%
<b>Age</b>		
18-25	7	2.9
26-30	179	74.9
31-35	51	21.3
36-40	2	0.8
41-45	0	0
<b>Gender</b>		
Female	214	90.3
Male	23	9.7
Other	0	0
<b>Racial/Ethnic Background</b>		
White/Caucasian	170	71.7
Black/African American	16	6.8
Asian and Pacific Islander	27	11.4
Hispanic/Latino	12	5.1
Middle Eastern	3	1.3
Multiracial	6	2.5
Other	3	1.3
<b>Current Relationship Status</b>		
Single/Never Married	83	34.7
Committed Relationship	52	21.8
Married	101	42.3

Separated	3	1.3
Other	0	0
<hr/>		
Currently Have Children		
Yes	33	13.9
No	205	86.1
<hr/>		
Religious Affiliation		
Christianity	110	46.6
Judaism	16	6.8
Islam	5	2.1
Hinduism	9	3.8
Buddhism	1	0.4
Other	7	3
None	88	37.3
<hr/>		
Post-Graduate Year		
1	62	25.9
2	66	27.6
3	53	22.2
4	57	23.8
Other	1	0.4
<hr/>		
Geographic Location		
West	35	14.8
Midwest	45	19
Northeast	90	38
South	59	24.9
Other	8	3.4
<hr/>		

---

Professional Plans

Maternal Fetal Medicine	18	7.5
Reproductive Endocrinology and Infertility	18	7.5
Urogynecology	17	7.1
Gynecologic Oncology	26	10.9
Family Planning	12	5
Other Fellowship	10	4.2
General Practice	120	50.2
Other	18	7.5

---

Table 2. OB/GYN Residents' Attitudes Toward Discussing Pre-Conception Planning and Fertility

Item	N	%	
Should an OB/GYN initiate discussions with patients about their potential childbearing intentions?			
Yes	198	91.7	
No	18	8.3	
Should an OB/GYN initiate discussions about age-related fertility decline with patients?			
Yes	180	82.9	
No	37	17.1	
Should discussing the natural decline in fertility with age be part of a well-woman annual exam with a gynecologist?			
Yes	157	72.4	
No	60	27.6	
Reasons for Yes	Educating women about this helps women make informed reproductive decisions	141	89.8
	I want to provide comprehensive health education to my patients	127	80.9
	Women should be aware of the correct relationship between fertility and age	112	71.3
	I can help dispel many of the myths in society/media regarding fertility and age	85	54.1
	Other	0	0
Reasons for No	Bringing this issue up annually is too frequent, but I am not opposed to discussing this issue with patients every three to four years	32	53.3
	I don't want to be perceived as pushing childbearing on patients	32	53.3
	Bringing up this issue annually may lead to emotional distress in my patients	24	40
	I want to be able to fully respect patient choices	23	38.3
	I don't have enough time	7	11.7
	Other	7	11.7
	It is not my primary responsibility	2	3.3

Table III Obstetrics and gynecology (OB/GYN) Residents' Awareness of Fertility Issues

Fertility Issue	All Residents (N = 217)	1st Year Residents (N = 55)	2nd Year Residents (N = 58)	3rd Year Residents (N = 51)	4th Year Residents (N = 53)	p
At what age is there a <i>slight</i> decrease in women's ability to become pregnant?	31.67 (3.10)	31.55 (3.47)	31.43 (2.74)	31.37 (2.95)	32.36 (3.21)	0.32
15-24	0.9%	1.8%	0%	0%	1.9%	
25-29*	9.7%	9.1%	10.3%	15.7%	3.8%	
30-34*	53.0%	50.9%	60.3%	54.9%	45.3%	
35-59	33.1%	38.2%	29.3%	29.4%	49.1%	
At what age is there a <i>marked</i> decrease in women's ability to become pregnant?	37.58 (2.53)	37.95 (2.54)	37.17 (2.62)	37.20 (2.51)	38.00 (2.39)	0.15
25-34	0%	0%	0%	0%	0%	
35-39*	52.5%	47.3%	60.3%	58.8%	43.4%	
40-44	46.5%	50.9%	37.9%	41.2%	56.6%	
45-59	0.9%	1.8%	1.7%	0%	0%	
What is the overall chance, on average, that a couple who undergoes treatment with in vitro fertilization will have a child after one treatment?	42.30 (18.37)	41.20 (19.52)	45.02 (18.75)	42.02 (18.61)	40.75 (16.62)	0.61
0-19%	5.1%	7.3%	1.7%	5.9%	5.7%	
20-29%*	16.6%	16.4%	17.2%	13.7%	18.9%	
30-39%	24.0%	29.1%	19.0%	29.4%	18.9%	
40-100%	54.4%	47.3%	62.1%	51.0%	56.6%	

\*Asterisk indicates the correct category based on published literature. For the slight decline in fertility, literature suggests this decline can begin in the late 20s to early 30s. *Note.* Means and standard deviations are presented in the first row for each fertility issue; percentages are presented in subsequent rows. Significance values come from one-way analyses of variance testing for differences based on year in residency.

Table 4. OB/GYN Residents' Attitudes Toward Use of Oocyte Cryopreservation

Item	N	%	
Should an OB/GYN initiate discussions regarding oocyte cryopreservation with female patients?			
Yes	83	39.9	
No	125	60.1	
Should discussing oocyte cryopreservation be part of a well-woman annual exam with a gynecologist?			
Yes	42	20.4	
No	164	79.6	
Reasons for Yes	Educating women about this issue helps women make more informed reproductive decisions	33	78.6
	Understanding the implications of oocyte cryopreservation increases women's childbearing choices	31	73.8
	I want to provide comprehensive health education to all my patients	28	66.7
	Other	1	2.4
Reasons for No	Bringing this issue up annually is too frequent, but I am not opposed to discussing this issue with patients every three to four years	79	48.2
	I don't want to be perceived as pushing childbearing on patients	49	29.9
	Other	45	27.4
	Bringing up this issue annually may lead to emotional distress in my patients	44	26.8
	I want to be able to fully respect patient choices	40	24.4
	It is not my primary responsibility	28	17.1
	I don't have enough time	25	15.2