

2013

Differential Allocation to Photosynthetic and Non-Photosynthetic Nitrogen Fractions among Native and Invasive Species

Jennifer L. Funk

Chapman University, jlfunk@chapman.edu

Lori A. Glenwinkle

Columbia University

Lawren Sack

University of California - Los Angeles

Follow this and additional works at: https://digitalcommons.chapman.edu/sees_articles



Part of the [Botany Commons](#), [Other Ecology and Evolutionary Biology Commons](#), [Other Plant Sciences Commons](#), and the [Plant Biology Commons](#)

Recommended Citation

Funk JL, Glenwinkle LA, Sack L (2013) Differential Allocation to Photosynthetic and Non-Photosynthetic Nitrogen Fractions among Native and Invasive Species. *PLoS ONE* 8(5): e64502. doi:10.1371/journal.pone.0064502

This Article is brought to you for free and open access by the Science and Technology Faculty Articles and Research at Chapman University Digital Commons. It has been accepted for inclusion in Biology, Chemistry, and Environmental Sciences Faculty Articles and Research by an authorized administrator of Chapman University Digital Commons. For more information, please contact laughtin@chapman.edu.

Differential Allocation to Photosynthetic and Non-Photosynthetic Nitrogen Fractions among Native and Invasive Species

Comments

This article was originally published in *PLoS ONE*, volume 8, issue 5, in 2013. DOI: [10.1371/journal.pone.0064502](https://doi.org/10.1371/journal.pone.0064502)

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

Copyright

The authors

Differential Allocation to Photosynthetic and Non-Photosynthetic Nitrogen Fractions among Native and Invasive Species

Jennifer L. Funk^{1*}, Lori A. Glenwinkel², Lauren Sack³

1 School of Earth and Environmental Sciences, Chapman University, Orange, California, United States of America, **2** Department of Biological Sciences, Columbia University, New York, New York, United States of America, **3** Department of Ecology and Evolutionary Biology, University of California Los Angeles, Los Angeles, California, United States of America

Abstract

Invasive species are expected to cluster on the “high-return” end of the leaf economic spectrum, displaying leaf traits consistent with higher carbon assimilation relative to native species. Intra-leaf nitrogen (N) allocation should support these physiological differences; however, N biochemistry has not been examined in more than a few invasive species. We measured 34 leaf traits including seven leaf N pools for five native and five invasive species from Hawaii under low irradiance to mimic the forest understory environment. We found several trait differences between native and invasive species. In particular, invasive species showed preferential N allocation to metabolism (amino acids) rather than photosynthetic light reactions (membrane-bound protein) by comparison with native species. The soluble protein concentration did not vary between groups. Under these low irradiance conditions, native species had higher light-saturated photosynthetic rates, possibly as a consequence of a greater investment in membrane-bound protein. Invasive species may succeed by employing a wide range of N allocation mechanisms, including higher amino acid production for fast growth under high irradiance or storage of N in leaves as soluble protein or amino acids.

Citation: Funk JL, Glenwinkel LA, Sack L (2013) Differential Allocation to Photosynthetic and Non-Photosynthetic Nitrogen Fractions among Native and Invasive Species. *PLoS ONE* 8(5): e64502. doi:10.1371/journal.pone.0064502

Editor: Mark van Kleunen, University of Konstanz, Germany

Received: November 11, 2012; **Accepted:** April 16, 2013; **Published:** May 20, 2013

Copyright: © 2013 Funk et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This research was supported by a grant from the USDA Biology of Weedy and Invasive Species In Agroecosystems Program (2008-35320-18721). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: jlfunk@chapman.edu

Introduction

The invasion of non-native species poses a rapidly growing threat to biodiversity and ecosystem function in many parts of the world, a threat equal to or stronger than that of climate change [1]. Thus, the need to understand the mechanisms of invasion has never been greater. Recent work has suggested that native and invasive plant species have similar carbon capture strategies but are aligned on opposite sides of a leaf economics spectrum [2]. For example, while native and invasive species show similar relationships among variables such as leaf nitrogen (N) concentration, photosynthetic rate (A), and leaf mass per unit area (LMA), invasive species cluster on the “high-return” end of the leaf economics spectrum with higher values of leaf N and A and lower values of LMA relative to natives [3,4]. Many of these traits, such as LMA, leaf N and A, correlate with relative growth rate [5,6], but see [7,8] and higher relative growth rates facilitate invasion in some systems [9], but see [10]. Intra-plant N allocation should reflect these trait differences [11]; however, the biochemical basis underlying these patterns has not been examined in native and invasive species.

Within leaves, N occurs in soluble and membrane-bound proteins which are involved in carbon assimilation and light harvesting, respectively, as well as non-photosynthetic compounds such as cell wall proteins, amino acids, nucleic acids, defense compounds (e.g., alkaloids), and inorganic N (NO_3^- , NH_4^+). Plant

species vary greatly in how N is allocated among these pools. For example, plants may allocate 5–32% of leaf N to Rubisco and 2–30% to cell walls, with higher amounts of cell wall protein in longer-lived leaves [12–17]. Notably, studies examining intra-leaf N pools have tended to focus primarily on cell-wall protein as the primary non-photosynthetic fraction without quantifying N allocation to amino acids, nucleic acids, inorganic N and secondary N-containing compounds [14–16,18,19], but see [20,21]. This level of resolution may be critical for determining important traits of invasive species. If invasive species occupy the “high-return” of the leaf economics spectrum with high A and low LMA, they should allocate more N to photosynthetic enzymes, and amino acids and nucleic acids as precursors of protein synthesis and growth. It is also possible that allocation of N to amino acids and nucleic acids (required for protein synthesis and growth) may have an equal or stronger effect on plant performance than increasing the amount of soluble protein involved in photosynthesis [21–22]. This hypothesis is supported by many studies where fertilization of woody species (primarily fast-growing forestry species) resulted in small changes in leaf N, but large changes in relative growth rate [23–29]. Conversely, if native species occupy the “slow-return” end of the leaf economics spectrum, requiring tougher, longer-lived leaves for a positive carbon balance, these species should allocate more N to structure

(cell-wall protein) at the expense of metabolic processes such as photosynthesis and growth.

The goal of this study was to examine intra-leaf N pools within a suite of native and invasive species and determine how these N pools correlate with leaf-level morphological and physiological traits, such as A and LMA. We characterized 34 leaf traits, and in particular leaf N partitioning with a high level of resolution, in five native and five invasive woody species that co-occur in Hawaii. All species establish on young, N-poor volcanic soils ranging in age from 100 to 10,000 years with ample water availability [30]; thus, plant growth is limited primarily by N availability in open areas, and co-limited by N and irradiance under closed canopies. While relative growth rates were not directly measured for these species, data collected in our study demonstrated that the invasive species selected generally had lower LMA than natives, aligning them closer to the “high-return” end of the leaf economics spectrum.

We examined seven intra-leaf N pools including soluble protein, membrane-bound protein, cell-wall protein, total nucleic acid, amino acid and inorganic N (NO_3^- , NH_4^+). The soluble protein fraction contains Calvin cycle and photorespiratory enzymes of which up to 50% is Rubisco [12]. All ten species were grown under shaded conditions ($300 \mu\text{mol photon m}^{-2} \text{s}^{-1}$ at mid-day) to simulate natural conditions for seedlings establishing in the understory of native mesic forests in Hawaii (dominated by *Metrosideros polymorpha*). Previous theoretical estimates based on gas exchange data have indicated that shade leaves should have a 1:1 ratio of soluble to membrane-bound protein [12,31]. We expected that invasive species would have higher A and lower LMA and, consequently, would allocate more N to soluble protein, amino acids and nucleic acids and less N to cell wall protein.

Materials and Methods

Plant material

Seeds were collected on the Island of Hawaii in November 2007 for the native species *Acacia koa* (Fabaceae), *Dodonaea viscosa* (Sapindaceae), *Osteomeles anthyllidifolia* (Rosaceae), *Pipturus albidus* (Urticaceae) and *Sophora chrysophylla* (Fabaceae), and for the invasive species *Falcataria moluccana* (Fabaceae), *Leucaena leucocephala* (Fabaceae), *Psidium cattleianum* (Myrtaceae), *Pyracantha angustifolia* (Rosaceae), and *Schinus terebinthifolius* (Anacardiaceae); nomenclature follows Wagner et al. [32]. All necessary permits were obtained for the described field studies (Tim Tunison, Hawaii Volcanoes National Park). All ten species are woody shrubs or trees with C_3 photosynthesis. All five invasive species are considered high risk weeds (A Global Compendium of Weeds, <http://www.hear.org/gcw/index.html>). Only one of the five native species (*Dodonaea*) is indigenous or naturalized outside of Hawaii. Because many of the worst invaders in Hawaii are legumes, we included two invasive (*Falcataria*, *Leucaena*) and two common native (*Acacia*, *Sophora*) legume species.

All non-legumes co-occur within Hawaii Volcanoes National Park and were chosen based on abundance, growth habit, and viability of seed. Seeds were germinated in potting soil in March 2008 and grown at the UCLA Plant Growth Facility greenhouse. After 3–6 months, plants were transferred to 3.6 L pots to avoid pot-binding. All plants received fertilization (0.2 mg) during irrigation every two days [Scotts Peters Professional water soluble fertilizer, N-P-K ratio 20-20-20; N as NH_4^+ (4.1%), NO_3^- (5.5%), urea (10.4%); P as P_2O_5 ; K as K_2O]. Pots were watered to field capacity. Irradiance levels within the greenhouse ranged from 60–550 $\mu\text{mol photon m}^{-2} \text{s}^{-1}$, with an average value of 300 $\mu\text{mol photon m}^{-2} \text{s}^{-1}$ at mid-day. Our efforts focused on intensive

physiological and biochemical measures and relative growth rate data were not collected in this study.

Gas exchange measurements

In January 2009, photosynthetic rates and chlorophyll fluorescence were measured with a LI-6400 portable photosynthesis system with a fluorescence chamber (LI-COR, Lincoln, NE, USA). All physiological and biochemical measures were conducted on fully-expanded, recently mature leaves. We conducted light response curves on two individuals per species and determined that 1000 $\mu\text{mol photon m}^{-2} \text{s}^{-1}$ was a saturating irradiance for all species. We conducted all measures at this irradiance. Relative humidity was maintained between 40–60% and ambient temperature was 25°C.

The effective quantum yield of PSII (ϕ_{PSII}), measured at 1000 $\mu\text{mol photon m}^{-2} \text{s}^{-1}$, was calculated as $(F_m' - F_s)/F_m'$, where F_s is the fluorescence yield of a light-adapted leaf and F_m' is the maximal fluorescence during a saturating light flash. CO_2 -response curves were determined by varying chamber CO_2 concentration between 50 and 2000 $\mu\text{L L}^{-1}$ (50, 100, 200, 400, 600, 1000, 1500, 2000). Maximum carboxylation rate (V_{max}) and maximum photosynthetic electron transport rate (J_{max}) were estimated from the CO_2 -response curves based on [33,34] using temperature responses outlined in ref. [35].

Following gas exchange measures, we measured leaf thickness with digital calipers at three points along the leaf to account for variation in thickness. Leaves were then harvested, scanned for leaf area, dried, and weighed to determine leaf mass per unit area (LMA). Leaf density (g cm^{-3}) was calculated by dividing LMA by leaf thickness. Total leaf N concentration was determined with an elemental analyzer (CE Instruments Flash EA 1112, CE Elantech). All physiology measurements were conducted on five individuals per species.

Biochemical measurements

Leaves adjacent to gas exchange leaves were harvested, immersed in liquid nitrogen and stored at -80°C prior to processing. We pulverized 30 to 100 mg of leaf tissue in liquid nitrogen prior to each extraction procedure. All reagents and solutions were made with ultrapure water (Barnstead NANOpure Infinity™ Ultrapure Water System. Prod. >17.00 M Ω –CM). Pigments were isolated in acetone and determined colorimetrically with a UV/VIS spectrophotometer (Beckman DU-640) following the methods of [36]. Chlorophyll a and b concentration were determined using a multi-wavelength analysis at 645, 662, and 710 nm. We did not have enough tissue to perform pigment analysis on *Acacia koa*.

Water-soluble, SDS-soluble, and SDS-insoluble protein fractions were extracted using a modified version of the procedure described in ref. [16] in which the SDS-soluble fraction consists of membrane-bound proteins and the SDS-insoluble fraction consists of cell-wall proteins. Previous work has shown a tight correlation between soluble protein and Rubisco concentration with Rubisco accounting for approximately 50% of the soluble protein fraction [31]. We used 4% polyvinylpyrrolidone (PVPP, or cross-linked insoluble PVP) in place of PVP with a 100 mM Tris Extraction buffer, pH 7.4 for the effective absorption of polyphenols [37]. The pH of 7.4 allows for maximum extraction efficiency combined with maximum polyphenol absorption by PVPP [38] and significantly increases water soluble protein yields [39]. While this protein fractionation method has been used in a diverse array of studies, e.g., [16,18], the method underestimates soluble and membrane-bound protein fractions when phenolic compounds are present. We corrected for the interference by phenolics for species

with total phenolic concentration greater than 12% dry leaf mass (*Pipturus*, *Psidium*, and *Schinus*) (see Methods S1). Our correction resulted in a $\pm 20\%$ error for protein fractions in these three species. This error is similar to or smaller than that associated with calculations of intra-leaf N partitioning which rely on estimates of Rubisco activity and bioenergetics derived from gas exchange measures [14,40–45].

The three protein fractions were hydrolyzed with barium hydroxide [46] and then quantified using an updated version of the ninhydrin method [47]. Because the ninhydrin method is poor for detecting hydroxyproline or proline [16], which are abundant in cell-wall proteins [17], this assay likely underestimates the amount of structural protein. However, this method is appropriate for assessing the relative differences in protein fractions among species. We used bovine serum albumin for standards.

Nucleic acids were extracted using a modified version of the method in ref. [20]. Samples were first extracted in 2 ml of chloroform:methanol (2:1). Samples were then sequentially extracted twice with chloroform:methanol:water (1:1:0.8), once with cold 80% methanol, and once with cold 5% trichloroacetic acid (TCA). All sequential extractions were performed for 15 min with 1 ml of extraction medium. After each extraction the samples were centrifuged at maximum speed for 2 minutes and the supernatant discarded. The final pellet was extracted twice with 2.5% TCA for one hour at 95°C. For total nucleic acid-nitrogen (TNA) quantification, the supernatant containing TNA was digested using the persulfate digest method [48] with 4.9 ml of water, 100 μ l of TNA extract, and 666 μ l of persulfate reagent. Each reaction vial was autoclaved at 120°C for 1.5 h [49], cooled at 4°C and buffered with 1 ml of 0.7 M ammonium chloride, pH 8.5 [48]. We added 0.5 to 0.6 g of spongy cadmium to each reaction tube and placed vials on a shaker table in the dark for 1.5 hours. We then added 250 μ l of color reagent [48] and allowed the samples to develop for 10 minutes in the dark. We read the absorbance at 540 nm using a microplate reader. Potassium nitrate and calf thymus DNA (Sigma Aldrich D4522) were used as standards.

Amino acids were extracted with 80% methanol [50] and quantified using the Ninhydrin method [47], modified for use with a microplate reader. We added 120 μ l of ninhydrin solution to 120 μ l of amino acid extract. The samples were then incubated at 100°C for 15 minutes and cooled on ice. We added 440 μ l of cold 50% isopropanol to each sample and read the absorbance at 570 nm. Ammonium was extracted as in ref. [51] using 2 ml of 80% methanol and quantified by fluorimetry using HPLC. Nitrate was extracted as in ref. [52] and absorbance was read at 405 nm using a microplate reader.

Statistical analysis

To determine differences in biochemical and gas exchange traits among five native and five invasive species, we used a nested ANOVA with ‘origin’ (native and invasive species) as a fixed effect and ‘species’ nested within origin. To determine differences in biochemical and gas exchange traits among legumes and non-legumes, we used a nested ANOVA with ‘legume’ as a fixed effect and ‘species’ nested within legume. Data that violated the ANOVA assumptions of normality and homogeneity of variance were Box–Cox transformed.

For traits that correlated with leaf mass per area (LMA), we tested whether the differences between native and invasive species could be explained by differences in LMA. We tested the trait-LMA relationships between native and invasive species for differences in slope and intercepts using SMATR (<http://www.bio.mq.edu.au/ecology/SMATR/index.html>; [53]). For this test,

we used raw and log-transformed data (i.e., fitting a power-law relationship).

A multivariate analysis was conducted on biochemical traits using principal components analysis (PCA). All traits were standardized prior to analysis using the formula [(trait – trait mean)/trait SD]. Pearson product–moment correlation coefficients were generated to evaluate the linear association among biochemical, physiological and morphological traits. All ANOVA, PCA and correlation analyses were performed in JMP 8 (SAS Institute, Inc., Cary, NC, U.S.A.).

Results

Nitrogen fraction recovery

Across species, we recovered between 38 to 108% of total leaf N in our seven fractions (soluble protein, membrane-bound protein, cell-wall protein, total nucleic acid, amino acid, NH_4^+ , and NO_3^- , see Results S1). Lower recovery in some species could be due to the presence of nitrogenous compounds not measured (e.g., alkaloids) or interference during N fraction extraction. These deviations did not occur in any systematic way; that is, the sum of N fractions was not substantially lower at low or high leaf N, for native versus invasive species, or for legumes versus non-legumes (see Results S1). For consistency across the results, we present the percentage of leaf N allocated to various N fractions using the value obtained by summing our seven fractions rather than that obtained by elemental analysis. Throughout, we use the term ‘concentration’ to refer to a parameter expressed per unit dry mass and ‘content per area’ for the area basis.

General patterns across species

The allocation of leaf N to protein (81.7 to 91.8%) varied relatively narrowly across species (Figure 1). However, species varied substantially in their distribution of protein among soluble, membrane-bound and cell-wall fractions. Total nucleic acid and amino acid concentrations and contents per area varied significantly among species (Figure 1). The fraction of leaf N allocated to nucleic acid ranged twofold from 5.4 to 11.2% across species while the allocation to total amino acid concentration ranged fivefold from 1.1 to 5.4% (Figure 1). Variation among species in inorganic N was driven by a few species as discussed below.

Variation in nitrogen partitioning between native and invasive species

Invasive species differed from natives in several leaf physiological and biochemical traits (Table 1). Invasive species had lower LMA and, consequently, lower leaf N content per area compared to native species. Invasive species also had lower photosynthetic function per area (A_{area} , V_{max} , J_{max} , Table 1) compared to native species. With respect to biochemical traits, native species had high absolute values of membrane-bound, cell-wall, and total protein concentrations and content per area compared to invasive species as well as a higher fraction of total leaf N allocated to membrane-bound protein (Table 1). Soluble protein concentration, content per area, and the ratio of soluble to membrane-bound protein (invasive, 1.33; native, 1.00) did not differ significantly between native and invasive species (Table 1). Invaders allocated a higher percentage of leaf N to total amino acid and NH_4^+ compared to native species, driven by higher NH_4^+ in *Psidium* and *Schinus* (Table 1, Table S1). Chlorophyll concentration and chlorophyll content per area were similar between native and invasive species.

These results were not confounded with differences in LMA across native and invasive species (data not shown). At a given LMA, most traits showed differences between native and invasive




Figure 1. The fraction of total leaf N present in six biochemical fractions for five native and five invasive species grown at 300 $\mu\text{mol photon m}^{-2} \text{s}^{-1}$. The percent of total leaf N represented by all three protein fractions is provided on the right. Inorganic N includes nitrate and ammonium. Species abbreviations are: *Acacia koa* (acko), *Dodonaea viscosa* (dovi), *Falcataria moluccana* (famo), *Leucaena leucocephala* (lele), *Osteomeles anthyllidifolia* (osan), *Pipturus albidus* (pial), *Psidium cattleianum* (psca), *Pyracantha angustifolia* (pyan), *Schinus terebinthifolius* (scte), and *Sophora chrysophylla* (soch). Legumes are marked with an asterisk (*). doi:10.1371/journal.pone.0064502.g001

species: for N_{area} and cell-wall protein concentration because the slopes of the trait against LMA differed between native and invasive species (P -values = 0.02–0.03) and for membrane-bound protein concentration and NH_4^+ content per area the slopes were statistically similar but the intercept was higher for native or invasive species (P -values = 0.01–0.03). For one trait, the percent of total leaf N allocated to amino acids, native and invasive showed the same relationship with LMA (slope $P=0.58$; intercept $P=0.97$), so the differences in that case were explained by LMA.

A multivariate analysis showed that two invasive species were biochemically distinct from all other species (Figure 2). *Psidium* tended to have high concentrations of soluble and NH_4^+ , and low concentrations of membrane-bound and cell-wall proteins. *Schinus* differed from other species in its high NH_4^+ concentration and low nucleic acid concentration.

Variation in nitrogen partitioning between legumes and non-legumes

Legumes generally had higher total protein and soluble protein concentrations and content per area (Table 1). Non-legumes allocated a larger fraction of leaf N to membrane-bound and cell-wall protein fractions than did legumes (Table 1). Legumes had substantially higher ratios of soluble to membrane-bound protein than non-legumes (1.71 vs. 0.81, Table 1). Total nucleic acids and amino acids were higher in legumes than non-legumes on both a mass and area basis (Table 1, Table S1). Leaf NO_3^- concentration was significantly higher in two legumes, *Leucaena* and *Sophora*, than in all other species (Figure 1, Table S1).

The multivariate analysis showed that legumes clustered together with high soluble protein and nucleic acid concentration when expressed as a percentage of total leaf N (Figure 2). The exception was *Sophora*, one of the native legumes, which clustered with the majority of non-legumes when traits were expressed on a mass basis.

Functional significance of N partitioning

Biochemical traits were more strongly inter-correlated on a mass basis than an area basis (Table 2). On a mass basis, soluble protein was strongly positively correlated with most biochemical fractions, with the exception of NH_4^+ , with which it was negatively correlated. Few biochemical fractions correlated with physiological function or leaf morphological traits. Membrane-bound protein correlated positively with light use efficiency (Φ_{PSII}), although this did not translate into higher rates of photosynthesis (Table 3). Species with high soluble and membrane-bound protein concentration tended to have higher rates of photosynthesis (A_{mass}), but this was not statistically significant. Leaf NH_4^+ was negatively correlated with all four measures of physiological function (A , V_{max} , J_{max} , Φ_{PSII}). Amino acid and chlorophyll concentrations (area basis) were negatively correlated with PNUE. Lastly, species with high LMA and density had higher area-based cell wall protein concentration, but lower mass-based soluble protein, nucleic acid and chlorophyll concentration (Table 3).

Table 1. Group means (and standard error in parentheses) for invasive and native species, and for legumes and non-legumes grown at 300 $\mu\text{mol photon m}^{-2} \text{s}^{-1}$.

	Native species (n = 5)	Invasive species (n = 5)	Effect of origin	Legumes (n = 4)	Non-legumes (n = 6)	Effect of legume
Physiology						
A _{area}	12.5 (0.7)	10.5 (0.6)	5.13*	11.8 (0.6)	11.3 (0.6)	0.49
A _{mass}	277 (31)	295 (28)	0.17	335 (30)	254 (29)	3.81
V _{max}	100.9 (5.4)	78.7 (5.0)	11.09**	106.8 (5.1)	78.4 (5.1)	17.65**
J _{max}	173 (9)	131 (9)	11.60**	171 (9)	139 (9)	8.70**
Φ _{PSII}	0.23 (0.01)	0.21 (0.01)	1.55	0.24 (0.01)	0.20 (0.01)	4.42*
N _{area}	1.72 (0.11)	1.34 (0.10)	6.53*	1.71 (0.11)	1.41 (0.10)	7.36**
N _{mass}	34.2 (1.2)	33.0 (1.1)	0.20	43.8 (1.1)	26.8 (1.1)	113.49**
PNUE	120 (13)	124 (12)	0.00	107 (12)	131 (12)	1.68
LMA	54.9 (3.1)	44.3 (2.8)	4.83*	40.7 (3.0)	55.5 (2.9)	10.37**
Leaf thickness	0.019 (0.001)	0.016 (0.001)	18.25**	0.016 (0.001)	0.019 (0.001)	20.77**
Leaf density	2.86 (0.18)	2.70 (0.16)	0.13	2.49 (0.17)	2.98 (0.17)	2.15
Biochemistry						
P _{sol} (mass)	11.3 (1.0)	10.6 (1.0)	0.08	17.20 (0.99)	6.80 (0.97)	46.58**
P _{sol} (area)	0.55 (0.09)	0.41 (0.08)	1.35	0.67 (0.09)	0.36 (0.09)	17.25**
P _{sol} (% N)	34.8 (2.5)	39.0 (2.4)	1.44	45.3 (2.5)	31.3 (2.4)	16.69**
P _{mem} (mass)	11.63 (0.98)	9.05 (0.93)	7.34**	11.96 (0.95)	9.26 (0.93)	5.81*
P _{mem} (area)	0.64 (0.07)	0.36 (0.07)	7.78**	0.49 (0.07)	0.51 (0.07)	0.09
P _{mem} (% N)	43.4 (2.2)	36.6 (2.1)	5.10*	33.8 (2.1)	44.1 (2.1)	12.26**
P _{cw} (mass)	2.22 (0.19)	1.71 (0.18)	5.58*	2.11 (0.19)	1.87 (0.18)	1.16
P _{cw} (area)	0.12 (0.01)	0.07 (0.01)	7.76**	0.08 (0.01)	0.10 (0.01)	1.59
P _{cw} (% N)	8.88 (0.65)	7.73 (0.61)	1.15	6.01 (0.63)	9.84 (0.61)	19.05**
A/P _{sol}	41.7 (7.1)	44.9 (6.6)	0.02	25.3 (6.9)	55.3 (6.6)	11.53**
P _{sol} /P _{mem}	1.00 (0.18)	1.33 (0.17)	3.24	1.71 (0.18)	0.81 (0.17)	15.90**
P _{total} (mass)	25.1 (1.5)	21.4 (1.4)	6.81*	31.3 (1.5)	17.9 (1.4)	45.68**
P _{total} (area)	1.31 (0.15)	0.85 (0.14)	14.43**	1.24 (0.15)	0.97 (0.14)	10.76**
AA (mass)	0.78 (0.09)	0.93 (0.08)	2.03	1.26 (0.08)	0.59 (0.08)	65.13**
AA (% N)	2.60 (0.30)	3.88 (0.28)	8.49**	3.43 (0.29)	3.12 (0.28)	7.08*
TNA (mass)	2.40 (0.16)	2.42 (0.15)	1.38	3.35 (0.16)	1.78 (0.15)	57.94**
TNA (% N)	8.49 (0.68)	9.17 (0.64)	0.49	9.44 (0.66)	8.42 (0.65)	1.51
NH ₄ ⁺ (mass)	0.11 (0.02)	0.21 (0.02)	3.64	0.05 (0.02)	0.23 (0.02)	71.50**
NH ₄ ⁺ (% N)	0.44 (0.20)	1.67 (0.19)	9.01**	0.19 (0.19)	1.63 (0.19)	126.53**
NO ₃ ⁻ (mass)	0.36 (0.11)	0.42 (0.11)	0.00	0.59 (0.11)	0.25 (0.11)	1.51
NO ₃ ⁻ (% N)	1.44 (0.45)	2.00 (0.44)	1.56	1.87 (0.45)	1.62 (0.43)	2.30
Chl (mass)	780 (76)	856 (62)	0.70	1043 (74)	712 (62)	11.58**
Chl (area)	40.9 (5.2)	35.9 (4.2)	0.66	41.1 (5.0)	36.6 (4.2)	1.02

Statistical effects of 'origin' (native/invasive) and 'legume' (legume/non-legume) are F-ratios with statistically significant values denoted by * $P < 0.05$ and ** $P < 0.01$. For both analyses, 'species' (df = 8) was nested within 'origin' (df = 1) or 'legume' (df = 1). Trait abbreviations: A_{area}, photosynthetic rate ($\mu\text{mol CO}_2 \text{ m}^{-2} \text{ leaf s}^{-1}$); A_{mass}, photosynthetic rate ($\text{nmol g}^{-1} \text{ s}^{-1}$); V_{max}, maximum rate of carboxylation ($\mu\text{mol CO}_2 \text{ m}^{-2} \text{ leaf s}^{-1}$); J_{max}, maximum electron transport rate ($\mu\text{mol electrons m}^{-2} \text{ s}^{-1}$); Φ_{PSII}, effective quantum yield of PSII, ($\Delta F/F_m$); N_{area}, leaf N (g N m^{-2}); N_{mass}, leaf N (mg N g^{-1}); PNUE, photosynthetic nitrogen use efficiency ($\mu\text{mol CO}_2 \text{ mol}^{-1} \text{ N s}^{-1}$); LMA, leaf mass per area (g m^{-2}); Leaf thickness (mm); Leaf density (g cm^{-3}); P_{sol}, soluble protein (mg N g^{-1} and g N m^{-2}); P_{mem}, membrane-bound protein (mg N g^{-1} and g N m^{-2}); P_{cw}, cell-wall protein (mg N g^{-1} and g N m^{-2}); AA, amino acid (mg N g^{-1}); TNA, total nucleic acid (mg N g^{-1}); NH₄⁺, ammonium (mg N g^{-1}); NO₃⁻, nitrate (mg N g^{-1}); Chl, Chlorophyll A + B ($\mu\text{g g}^{-1}$ and mg m^{-2}); % N, percent of total leaf N.

doi:10.1371/journal.pone.0064502.t001

Discussion

Intra-leaf N partitioning should reflect trade-offs on the leaf economics spectrum with faster growing species allocating more N to metabolic processes at the expense of structure. Thus, we hypothesized that invasive species, which are generally located on the "high-return" end of the leaf economics spectrum, would have

lower LMA and higher A relative to native species, with greater allocation to leaf N pools associated with photosynthesis and growth. In support of this idea, across all species, LMA was negatively correlated with soluble protein, total nucleic acid and chlorophyll content and positively correlated with cell wall protein. Invasive species as a group did have lower LMA and cell wall




Figure 2. Results of a principal components analysis for six N fractions from five native and five invasive (*) species grown at 300 $\mu\text{mol photon m}^{-2} \text{s}^{-1}$ on a mass basis (A) and as a percentage of total leaf N (B). Legumes tended to cluster together (dashed square) in both panels. Two invasive species differed from the other species: *Psidium* had lower membrane-bound and cell-wall protein and higher amino acid concentration whereas *Schinus* had high NH_4^+ concentration. Species abbreviations are provided in Figure 1. doi:10.1371/journal.pone.0064502.g002

protein with higher amino acid content, consistent with allocation to growth at the expense of structure. However, our hypothesis that invasive species would allocate more resources to carbon assimilation and growth at the expense of structure was only partially supported. Photosynthetic rates were either similar (mass-based) or higher (area-based) in native species relative to invasive species, soluble protein did not differ between groups, and native species had higher amounts of total N and membrane-bound protein.

Our finding that native and invasive species did not differ in the allocation of N to soluble protein contrasts with previously published results that invasive populations of *Ageratina adenophora* allocated more N to soluble protein at the expense of cell-wall protein compared with native populations [18]. Despite no difference in soluble protein between native and invasive species, native species had higher A_{area} and V_{max} compared with invasive species and this suggests that Rubisco content or activity may have

been higher in native species. Our soluble protein fraction includes Rubisco and many other proteins, but Rubisco was not directly measured in this study.

The low irradiance used in our experiment (which reflects actual growing conditions of seedlings in Hawaiian forest understory) resulted in relatively even allocation of N to light harvesting (e.g., membrane-bound protein) and carbon assimilation (e.g. soluble protein) functions. Our data matched theoretical estimates modeled from gas exchange data, indicating that shade leaves should have a 1:1 ratio of soluble and membrane-bound protein whereas sun leaves should have 2–3 times higher soluble than membrane-bound protein [12,31]; in our light-limited growth environment, we found roughly a 1:1 ratio of soluble to membrane-bound protein with no significant difference between native and invasive species. However, native species allocated more N to membrane-bound protein (43.4% of total N) than invasive species (36.6% of total N). Because the membrane-bound

Table 2. Correlation coefficients (r) for relationships among leaf N fractions for five native and five invasive species grown at 300 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ and high nutrient availability (n = 10 species).

	P_{sol}	P_{mem}	P_{cw}	AA	TNA	NH_4^+	NO_3^-	Chl	N
P_{sol}	–	0.78**	0.50	0.71*	0.88**	–0.45	–0.04	0.76*	0.75*
P_{mem}	0.79**	–	0.84**	0.40	0.72*	–0.20	0.06	0.62+	0.68*
P_{cw}	0.63*	0.85**	–	0.30	0.59+	0.06	–0.10	0.44	0.56+
AA	0.75*	0.40	0.30	–	0.70*	–0.09	0.19	0.68*	0.72*
TNA	0.93**	0.76*	0.72*	0.75*	–	–0.43	–0.07	0.66+	0.53
NH_4^+	–0.65*	–0.57+	–0.54	–0.38	–0.59+	–	0.11	0.13	0.01
NO_3^-	0.29	0.29	–0.05	0.30	0.25	–0.14	–	0.50	0.16
Chl	0.86**	0.64+	0.35	0.63+	0.78*	–0.25	0.70*	–	0.74*
N	0.93**	0.68*	0.41	0.74*	0.77*	–0.63+	0.40	0.86**	–

Correlations between area- and mass-based traits are on the upper half and lower half of the table, respectively. Correlations were performed on log-transformed data. Significant correlations at $P < 0.10$ (+), $P < 0.05$ (*) and $P < 0.01$ (**) are shown. Trait abbreviations are given in Table 1. doi:10.1371/journal.pone.0064502.t002

Table 3. Correlation coefficients (r) for relationships among leaf biochemical, physiological and morphological traits for five native and five invasive species grown at 300 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ and high nutrient availability (n = 10 species).

	A_{area}	A_{mass}	V_{max}	J_{max}	ϕ_{PSII}	PNUE	LMA	Density	Thick
P _{sol} (mass)	0.18	0.53	0.33	0.06	0.49	-0.05	-0.59+	-0.54	-0.18
P _{sol} (area)	0.14	0.10	0.44	0.16	0.56+	-0.41	-0.02	0.05	0.05
P _{sol} (% N)	-0.07	0.48	0.04	-0.01	0.13	-0.05	-0.66*	-0.50	-0.43
P _{mem} (mass)	0.43	0.53	0.50	0.14	0.71*	0.15	-0.42	-0.49	0.10
P _{mem} (area)	0.33	-0.04	0.54	0.23	0.64*	-0.25	0.27	0.12	0.37
P _{mem} (% N)	0.29	-0.12	0.16	0.15	0.11	0.29	0.37	0.25	0.31
P _{cw} (mass)	0.39	0.34	0.44	0.04	0.67*	0.14	-0.24	-0.38	0.27
P _{cw} (area)	0.18	-0.67	0.36	0.11	0.43	-0.31	0.58+	0.35	0.55+
P _{cw} (% N)	-0.02	-0.50	-0.13	-0.04	-0.23	0.09	0.63*	0.50	0.36
AA (mass)	0.01	0.19	0.15	0.06	0.23	-0.33	-0.30	-0.28	-0.10
AA (area)	-0.09	-0.23	0.16	0.11	0.17	-0.64*	0.19	0.15	0.10
AA (% N)	-0.30	-0.24	-0.19	0.01	-0.30	-0.47	0.07	0.12	-0.12
TNA (mass)	0.10	0.47	0.21	-0.02	0.39	0.00	-0.59+	-0.68*	0.09
TNA (area)	0.03	0.03	0.29	0.07	0.40	-0.33	-0.05	-0.26	0.40
TNA (% N)	-0.29	0.08	-0.25	-0.20	-0.22	-0.01	-0.37	-0.57+	0.30
NH ₄ ⁺ (mass)	-0.57+	-0.58+	-0.81**	-0.67*	-0.71*	-0.24	0.31	0.23	0.21
NH ₄ ⁺ (area)	-0.53	-0.78**	-0.66*	-0.50	-0.64*	-0.43	0.63*	0.52	0.33
NH ₄ ⁺ (% N)	-0.52	-0.64*	0.72*	-0.48	-0.74*	-0.19	0.45	0.40	0.18
NO ₃ ⁻ (mass)	-0.06	0.28	0.18	0.14	-0.06	0.00	-0.43	-0.42	-0.11
NO ₃ ⁻ (area)	-0.18	-0.21	0.18	0.24	-0.22	-0.37	0.14	0.09	0.10
NO ₃ ⁻ (% N)	-0.28	-0.20	-0.19	0.01	-0.54	-0.05	0.07	0.13	-0.13
Chl (mass)	-0.28	0.31	-0.05	-0.27	0.01	-0.16	-0.62+	-0.69*	0.06
Chl (area)	-0.47	-0.49	0.00	-0.16	-0.08	-0.81**	0.30	0.10	0.45

Correlations were performed on log-transformed data. Significant correlations at $P < 0.10$ (+), $P < 0.05$ (*) and $P < 0.01$ (**) are shown. Trait abbreviations are given in Table 1.

doi:10.1371/journal.pone.0064502.t003

protein fraction includes pigment-protein complexes which promote effective light harvesting and electron transport, higher amounts of membrane-bound protein in native species may have contributed to the higher photosynthetic rates (A_{area} , J_{max}) under our light-limited growth conditions. While membrane-bound protein was higher in native species relative to invasive species, chlorophyll content was similar and this suggests that the membrane-bound protein fraction may represent protein that may or may not be involved in carbon assimilation.

Two invasive species (*Falcataria*, *Psidium*) and one native species (*Acacia*) displayed high soluble protein concentration and, consequently, low photosynthetic rates per unit soluble protein (A/P_{sol}), which conflicts with predictions of optimal N partitioning models. These models, run across irradiance environments, suggest that shade leaves should allocate less N to soluble protein as photosynthesis becomes limited by light harvesting rather than carbon assimilation at low irradiance [54]. Although a high production of soluble protein seems like a wasteful use of N at low irradiance, it is possible that higher protein and amino acid concentration in *Acacia*, *Falcataria* and *Psidium* represented N storage [55,56] and this could allow rapid growth in a heterogeneous light environment (e.g., tree gaps, forest edges), although the rate of amino acid production rather than pool size is more commonly related to relative growth rate. A recent study supports the idea that *Acacia* has high phenotypic plasticity in response to light availability [57] which suggests

that stored N may be useful for rapid growth under changing light conditions.

Our hypotheses for N allocation outside of protein were partially supported. We predicted that invasive species would allocate more N to amino acids and nucleic acids which are required in greater amounts in rapidly growing tissues [21,22]. While invasive species allocated a larger percentage of leaf N to amino acids compared with native species, total nucleic acid concentration did not vary between groups. Strong species-level variation was evident, and thus differences among native and invasive species for a given N fraction were often driven by a single species. Considering all the biochemical fractions, two invaders were distinct from all other species. *Psidium* tended to have high amino acid concentration, high NH_4^+ concentration, and low membrane-bound and cell-wall protein concentration while *Schinus* differed from other species primarily in its high NH_4^+ concentration and low nucleic acid concentration. These findings at the biochemical level are analogous to other studies that showed great trait variation within groups of native and/or invasive species: such a pattern has been reported for leaf nutrient concentrations, LMA, canopy height, seed mass, relative growth rate, and water use [2,58–60]. Thus, the results of comparisons of native and invasive species will depend on the particular species considered. The overall average differences between natives and invasives are instructive, but species-level comparisons provide higher resolution that might indicate outcomes of these trait differences at the

plant level. Thus, the pertinent question is: what are the functional consequences of this inter-specific variation in N partitioning?

The partitioning patterns for *Psidium* were consistent with general expectations concerning invasive species: more N was allocated to growth (amino acids) at the expense of structure (cell-wall protein). *Psidium* may succeed in dense understories by increasing amino acid production for transport out of the leaf to fuel the production of new structures. Increasing whole-plant leaf area will maximize light interception in Hawaii's light-limited mesic forests. Our results concur with those of Niinemets *et al.* [28] who found that an invasive species (*Rhododendron ponticum*) diluted leaf N at a high fertility, forested site, making more leaves rather than increasing leaf N concentration. The authors concluded that, at a plant level, greater light interception may be more advantageous than a higher Chl/N ratio in a forest understory.

Conversely, the patterns in *Schinus*, particularly its high levels of NH_4^+ , suggested stress rather than optimal N partitioning. NH_4^+ accumulation in leaves is toxic and under normal conditions leaf NH_4^+ levels are relatively constant in plants. Differences in leaf NH_4^+ among species may indicate a decrease in amino acid production most likely resulting from stress associated with low irradiance. Low total protein concentration in *Schinus* drove the overall higher total protein concentration of natives relative to invasives. While *Schinus* seedlings often recruit within dense patches of established adult plants, our results suggest that this species spreads by a mechanism other than superior leaf biochemistry, possibly vegetative reproduction into high light environments (J Funk, personal observation).

We found that nitrogen-fixation strongly influenced intra-leaf N allocation. Legumes had higher total protein content, higher soluble protein content, and a higher ratio of soluble to membrane-bound protein relative to non-legumes. As noted above, this pattern of N allocation may be disadvantageous in a light-limited habitat where membrane-bound protein will increase light harvesting. However, we should note that A was not correlated with membrane-bound protein across species in this study. A study of nitrogen-fixing and non-fixing plant species in the United States found that nitrogen-fixing species tended to be less shade tolerant than non-fixers [61]; thus, high soluble protein in these species may be beneficial in open canopies or for rapid growth when gaps appear in the canopy. As discussed above, legumes may store N as soluble protein for use when light becomes available in the understory.

We acknowledge limitations in our study design. First, we were unable to find seeds for co-occurring, phylogenetically related pairs of native and invasive species or pairs of invasive and noninvasive exotic species. Because more closely related taxa share similar trait values, phylogenetic comparative designs minimize trait differences associated with comparing unrelated species and disparate life forms. Thus, future biochemical studies should include more taxonomically based species comparisons. A second limitation was that we did not directly measure relative growth rates of our species. While many studies infer inter-specific differences in hard-to-measure processes like growth from easy-to-measure traits like LMA based on established leaf economics spectrum relationships, e.g., [5], we assumed that our invasive species were located closer to the "high return" end of the leaf economics spectrum relative to native species based on differences in LMA. Consequently, we were unable to directly link species differences in amino acid and nucleic acid content to relative growth rates. Third, our study was conducted at high N

availability and we may expect to see greater differences in N allocation, and stronger differences between native and invasive species in N partitioning and leaf-level traits, under low N conditions. Our experiment initially included a low N treatment but high mortality resulted in too few replicates for our final analyses.

Lastly, our estimates of protein fractions for the three species with leaf phenolic content greater than 12% (*Pipturus*, *Psidium*, *Schinus*) and low percent of N recovery for some species added error to the data and, consequently, our interpretation. In particular, we only recovered 38% of total leaf N from *Schinus* which, along with *Psidium*, was an outlier with respect to several N pools. *Schinus* is the only species we surveyed that contains leaf resin which may have interfered with extraction and quantification of soluble protein. Underestimates of soluble protein in *Schinus* and *Psidium* would result in lower amounts of NH_4^+ and amino acids than we report here, making the two species more similar in biochemistry to the other invasive species measured. Additionally, *Sophora* had lower soluble protein content relative to other legumes, and low N recovery (64%) for this species may reflect an underestimate of soluble protein.

Despite these limitations, our study is the first to examine intra-leaf N partitioning in a suite of native and invasive species. Our data suggest that invasive species employ a wide range of mechanisms in N allocation. For example, some invasive species (particularly *Falcataria* and *Leucaena*) may succeed as invaders in low-light environments by storing N as protein and amino acids that can be used when high-light conditions become available. Conversely, our results suggest that one invader (*Psidium*) may succeed by allocating N to growth at the expense of higher leaf-level carbon assimilation. More studies are needed to confirm these patterns across a larger number of native and invasive species and to evaluate the potential importance of biochemistry, in combination with other factors (e.g., clonality, enemy release, seed dispersal), in contributing to the success of invasive species.

Supporting Information

Table S1 Trait values for five native and five invasive species.

(PDF)

Methods S1 Correction of protein yields for interference by polyphenols.

(PDF)

Results S1 Nitrogen recovery in biochemical fractions.

(PDF)

Acknowledgments

Many thanks to Crissy Anderson, Warren De Bruyn, Brittany Hickok, Savannah Lane, Cynthia Neben, Jessica Pasquet-Kok, Erick Reisinger, Christyn Takara, Jaclyn Vanmansart, Martin Vignovich and Ginny Zachary for lab and greenhouse assistance.

Author Contributions

Conceived and designed the experiments: JF. Performed the experiments: JF LG. Analyzed the data: JF LG LS. Contributed reagents/materials/analysis tools: JF LS. Wrote the paper: JF LS.

References

- Millennium Ecosystem Assessment (2005) Ecosystems and Human Well-Being: Biodiversity Synthesis. Washington, DC:World Resources Institute.
- Leishman MR, Thomson VP, Cooke J (2010) Native and exotic invasive plants have fundamentally similar carbon capture strategies. *Journal of Ecology* 98: 28–42.
- Leishman MR, Haslehurst T, Ares A, Baruch Z (2007) Leaf trait relationships of native and invasive plants: community- and global-scale comparisons. *New Phytologist* 176:635–643
- Penuelas J, Sardans J, Llusia J, Owen SM, Carnicer J, et al. (2010) Faster return on 'leaf economics' and different biogeochemical niche in invasive compared with native plant species. *Global Change Biology* 16: 2171–2185.
- Reich PB, Wright IJ, Lusk CH (2007) Predicting leaf physiology from simple plant and climate attributes: A global glonpnet analysis. *Ecological Applications* 17: 1982–1988.
- Wright IJ, Westoby M (2000) Cross-species relationships between seedling relative growth rate, nitrogen productivity and root vs leaf function in 28 Australian woody species. *Functional Ecology* 14: 97–107.
- Poorter L, Wright SJ, Paz H, Ackerly DD, Condit R, et al. (2008) Are functional traits good predictors of demographic rates? Evidence from five neotropical forests. *Ecology* 89: 1908–1920.
- Shipley B (2002) Trade-offs between net assimilation and specific leaf area in determining relative growth rate: Relationship with daily irradiance. *Functional Ecology* 16: 682–689.
- Grotkopp E, Erskine-Ogden J, Rejmanek M (2010) Assessing potential invasiveness of woody horticultural plant species using seedling growth rate traits. *Journal of Applied Ecology* 47: 1320–1328.
- Daehler CC (2003) Performance comparisons of co-occurring native and alien invasive plants: implications for conservation and restoration. *Annual Review of Ecology, Evolution and Systematics* 34: 183–211.
- Field CB, Mooney HA (1986) The photosynthesis-nitrogen relationship in wild plants. In: Givnish TJ, editor. *On the Economy of Plant Form and Function*. Cambridge, England:Cambridge University Press. pp. 25–55.
- Evans JR (1989) Photosynthesis and nitrogen relationships in leaves of C3 plants. *Oecologia* 78: 9–19.
- Warren CR, Adams MA (2001) Distribution of N, Rubisco and photosynthesis in *Pinus pinaster* and acclimation to light. *Plant, Cell and Environment* 24: 597–609.
- Harrison MT, Edwards EJ, Farquhar GD, Nicotra AB, Evans JR (2009) Nitrogen in cell walls of sclerophyllous leaves accounts for little of the variation in photosynthetic nitrogen-use efficiency. *Plant, Cell and Environment* 32: 259–270.
- Onoda Y, Hikosaka K, Hirose T (2004) Allocation of nitrogen to cell walls decreases photosynthetic nitrogen-use efficiency. *Functional Ecology* 18: 419–425.
- Takashima T, Hikosaka K, Hirose T (2004) Photosynthesis or persistence: nitrogen allocation in leaves of evergreen and deciduous *Quercus* species. *Plant, Cell and Environment* 27: 1047–1054.
- Reiter WD (1998) The molecular analysis of cell wall components. *Trends in Plant Science* 3: 27–32.
- Feng Y, Lei Y-B, Wang Y-P, Callaway RM, Valiente-Banuet A, et al. (2009) Evolutionary tradeoffs for nitrogen allocation to photosynthesis versus cell walls in an invasive plant. *Proceedings of the National Academy of Sciences* 106: 1853–1856.
- Hikosaka K, Shigeno A (2009) The role of Rubisco and cell walls in the interspecific variation in photosynthetic capacity. *Oecologia* 160: 443–451.
- Chapin III FS, Kedrowski RA (1983) Seasonal changes in nitrogen and phosphorus fractions and autumn retranslocation in evergreen and deciduous taiga trees. *Ecology* 64: 376–391.
- Chapin III FS, Shaver GR, Kedrowski RA (1986) Environmental controls over carbon, nitrogen and phosphorus fractions in *Eriophorum vaginatum* in Alaskan tussock tundra. *Journal of Ecology* 74: 167–195.
- Lambers H, Poorter H (1992) Inherent variation in growth rate between higher plants: a search for physiological causes and ecological consequences. *Advances in Ecological Research* 23: 187–261.
- Brix H (1981) Effects of fertilizer source and application rates on foliar nitrogen concentration, photosynthesis, and growth of Douglas-fir. *Canadian Journal of Forest Research* 11: 775–780.
- Walters MB, Reich PB (1989) Response of *Ulmus americana* seedlings to varying nitrogen and water status. 1. Photosynthesis and growth. *Tree Physiology* 5: 159–172.
- Wendler R, Millard P (1996) Impacts of water and nitrogen supplies on the physiology, leaf demography and nitrogen dynamics of *Betula pendula*. *Tree Physiology* 16: 153–159.
- Coleman MD, Dickson RE, Isebrands JG (1998) Growth and physiology of aspen supplied with different fertilizer addition rates. *Physiologia Plantarum* 103: 513–526.
- Warren CR, Chen ZL, Adams MA (2000) Effect of N source on concentration of Rubisco in *Eucalyptus diversicolor*, as measured by capillary electrophoresis. *Physiologia Plantarum* 110: 52–58.
- Niinemets U, Valladares F, Ceulemans R (2003) Leaf-level phenotypic variability and plasticity of invasive *Rhododendron ponticum* and non-invasive *Ilex aquifolium* co-occurring at two contrasting European sites. *Plant, Cell and Environment* 26: 941–956.
- Funk JL, Jones CG, Lerdau MT (2007) Leaf- and shoot-level plasticity in response to varying nutrient and water availabilities. *Tree Physiology* 27: 1731–1739.
- Vitousek PM, Walker LR, Whiteaker LD, Matson PA (1993) Nutrient limitations to plant growth during primary succession in Hawaii Volcanoes National Park. *Biogeochemistry* 23: 197–215.
- Terashima I, Evans JR (1988) Effects of light and nitrogen nutrition on the organization of the photosynthetic apparatus in spinach. *Plant and Cell Physiology* 29: 143–155.
- Wagner WL, Herbst DR, Sohmer SH (1999) *Manual of the Flowering Plants of Hawaii*. Honolulu:University of Hawaii Press.
- Farquhar GD, von Caemmerer S, Berry JA (1980) A biochemical model of photosynthetic CO₂ assimilation in leaves of C3 species. *Planta* 149: 78–90.
- Wullschlegel SD (1993) Biochemical limitations to carbon assimilation in C3 plants - a retrospective analysis of the A/Ci curves from 109 species. *Journal of Experimental Botany* 44: 907–920.
- Long SP, Bernacchi CJ (2003) Gas exchange measurements, what can they tell us about the underlying limitations to photosynthesis? Procedures and sources of error. *Journal of Experimental Botany* 54: 2393–2401.
- Lichtenthaler HK, Buschmann C (2001) Chlorophylls and carotenoids: measurement and characterization by UV-VIS spectroscopy. *Current Protocols in Food Analytical Chemistry*. New York: John Wiley and Sons. pp. F4.3.1–F4.3.8.
- Isaacson T, Damasceno CMB, Saravanan RS, He Y, Catala C, et al. (2006) Sample extraction techniques for enhanced proteomic analysis of plant tissues. *Nature Protocols* 1: 769–774.
- Loomis WD, Battaile J (1966) Plant phenolic compounds and the isolation of plant enzymes. *Phytochemistry* 5: 423–428.
- Warren CR (2000) Is photosynthesis related to concentrations of nitrogen and Rubisco in leaves of Australian native plants? *Australian Journal of Plant Physiology* 27: 407–416.
- Mao QZ, Watanabe M, Imori M, Kim YS, Kita K, et al. (2012) Photosynthesis and nitrogen allocation in needles in the sun and shade crowns of hybrid larch saplings: Effect of nitrogen application. *Photosynthetica* 50: 422–428.
- Frak E, Le Roux X, Millard P, Dreyer E, Jaouen G, et al. (2001) Changes in total leaf nitrogen and partitioning of leaf nitrogen drive photosynthetic acclimation to light in fully developed walnut leaves. *Plant, Cell and Environment* 24: 1279–1288.
- Warren CR, Adams MA (2004) Evergreen trees do not maximize instantaneous photosynthesis. *Trends in Plant Science* 9: 270–274.
- Evans JR (1996) Developmental constraints on photosynthesis: effects of light and nutrition. In: Baker NR, editor. *Photosynthesis and the Environment*. The Netherlands: Kluwer Academic Publishers. pp. 281–304.
- Evans JR, Poorter H (2001) Photosynthetic acclimation of plants to growth irradiance: the relative importance of specific leaf area and nitrogen partitioning in maximizing carbon gain. *Plant, Cell and Environment* 24: 755–767.
- Pons TL, Westbeek MHM (2004) Analysis of differences in photosynthetic nitrogen-use efficiency between four contrasting species. *Physiologia Plantarum* 122: 68–78.
- McGrath R (1972) Protein measurement by ninhydrin determination of amino acids released by alkaline hydrolysis. *Analytical Biochemistry* 49: 95–102.
- Sun S-W, Lin Y-C, Weng Y-M, Chen M-J (2006) Efficiency improvements on ninhydrin method for amino acid quantification. *Journal of food composition and analysis* 19: 112–117.
- Bronk DA, Lomas MW, Gilbert PM, Schukert KJ, Sanderson MP (2000) Total dissolved nitrogen analysis: comparisons between the persulfate, UV and high temperature oxidation methods. *Marine Chemistry* 69: 163–178.
- Purcell LC, King CA (1996) Total nitrogen determination in plant material by persulfate digestion. *Agronomy Journal* 88: 111–113.
- Noctor G, Bergot GL, Mauve C, Thominet D, Lelarge-Trouverie C, et al. (2007) A comparative study of amino acid measurement in leaf extracts by gas chromatography-time of flight-mass spectrometry and high performance liquid chromatography with fluorescence detection. *Metabolomics* 3: 161–174.
- Husted S, Hebborn CA, Mattsson M, Schjoerring JK (2000) A critical experimental evaluation of methods for determination of NH₄⁺ in plant tissue, xylem sap and apoplastic fluid. *Physiologia Plantarum* 109: 167–179.
- Leleu O, Vuylstecker C (2004) Unusual regulatory nitrate reductase activity in cotyledons of *Brassica napus* seedlings: enhancement of nitrate reductase activity by ammonium supply. *Journal of Experimental Botany* 55: 815–823.
- Warton DI, Wright IJ, Falster DS, Westoby M (2006) Bivariate line-fitting methods for allometry. *Biological Reviews* 81: 259–291.
- Hikosaka K, Terashima I (1995) A model of the acclimation of photosynthesis in the leaves of C3 plants to sun and shade with respect to nitrogen use. *Plant, Cell and Environment* 18: 605–618.
- Nasholm T, Edfast A-B, Ericsson A, Norden L-G (1994) Accumulation of amino acids in some boreal forest plants in response to increased nitrogen availability. *New Phytologist* 126: 137–143.

56. Warren CR, Adams MA (2000) Capillary electrophoresis for the determination of major amino acids and sugars in foliage: application to the nitrogen nutrition of sclerophyllous species. *Journal of Experimental Botany* 51: 1147–1157.
57. Funk JL (2008) Differences in plasticity between invasive and native plants from a low resource environment. *Journal of Ecology* 96: 1162–1174.
58. Cavaleri MA, Sack L (2010) Comparative water use of native and invasive plants at multiple scales: a global meta-analysis. *Ecology* 91: 2705–2715.
59. Ordonez A, Wright IJ, Olff H (2010) Functional differences between native and alien species: a global-scale comparison. *Functional Ecology* 24: 1353–1361.
60. van Kleunen M, Weber E, Fischer M (2010) A meta-analysis of trait differences between invasive and non-invasive plant species. *Ecology Letters* 13: 235–245.
61. Menge DNL, DeNoyer JL, Lichstein JW (2010) Phylogenetic constraints do not explain the rarity of nitrogen-fixing trees in late-successional temperate forests. *PLOS One* 5: e12056. doi:10.1371/journal.pone.0012056.