


2006

Interdependency of Pharmacokinetic Parameters: A Chicken-and-Egg Problem? Not!

Reza Mehvar

Chapman University, mehvar@chapman.edu

Follow this and additional works at: https://digitalcommons.chapman.edu/pharmacy_articles

 Part of the [Numerical Analysis and Scientific Computing Commons](#), [Other Mathematics Commons](#), [Pharmacy and Pharmaceutical Sciences Commons](#), and the [Software Engineering Commons](#)

Recommended Citation

Mehvar R, "Interdependency of Pharmacokinetic Parameters: A Chicken-and-Egg Problem? Not!" *J. Pharm. Pharm. Sci.*, 9:113-118 (2006).

This Article is brought to you for free and open access by the School of Pharmacy at Chapman University Digital Commons. It has been accepted for inclusion in Pharmacy Faculty Articles and Research by an authorized administrator of Chapman University Digital Commons. For more information, please contact laughtin@chapman.edu.

Interdependency of Pharmacokinetic Parameters: A Chicken-and-Egg Problem? Not!

Comments

This article was originally published in *Journal of Pharmacy and Pharmaceutical Sciences*, volume 9, in 2006.

Copyright

Canadian Society for Pharmaceutical Sciences

Interdependency of pharmacokinetic parameters: A chicken-and-egg problem? Not!

Reza Mehvar*

School of Pharmacy, Texas Tech University Health Sciences Center, Amarillo, Texas, USA

Received 24 February 2006, Accepted 27 February 2006,
Published 08 March 2006

Abstract Pharmacokinetic (PK) software packages are widely used by scientists in different disciplines to estimate PK parameters. However, their use without a clear understanding of physiological parameters affecting the PK parameters and how different PK parameters are related to each other may result in erroneous interpretation of data. Often, mathematical relationships used for the estimation of PK parameters obscure the true physiological relationships among these parameters, prompting a discussion of which parameter came first and giving the appearance of the-chicken-and-the-egg dilemma. In this article, the author attempts to show how different PK parameters are related to physiological parameters and each other by using various scenarios and examples. In particular, the relationship between clearance and the rate of elimination and that among the other major PK parameters are explored. It is concluded that there is no dilemma in interdependency of the PK parameters, and the relationships among the PK parameters and between PK and physiological parameters are clear.

INTRODUCTION

The origin of pharmacokinetics is attributed to an article written by Torsten Torell in 1937 (1). After an initial fascination with mathematical relationships defining this discipline, came the introduction of a large number of easy-to-use pharmacokinetic (PK) software packages. Because of widespread application of pharmacokinetics in other disciplines, such as biology, pharmacology, and physiology, and the availability of software, the use of pharmacokinetics in biological sciences has grown substantially in the last two decades. However, the use of the PK software without a clear understanding of physiological parameters affecting the PK parameters and how different PK parameters

are related to each other may result in erroneous interpretation of data.

One of the most common errors made by biological scientists, whose main focus and education are not in the area of pharmacokinetics, is the distinction between mathematical and physiological relationships among PK parameters (2). In other words, when two PK parameters are changed, which parameter is the cause and which one is the effect. For example, a recent heated discussion was erupted among some of the subscribers to the PharmPK LISTSERV[†] about whether the clearance of a drug is dependent on the rate of elimination or vice versa, with apparently no final resolution. The advocates of the dependency of *CL* on the rate of elimination (dA_e/dt) cited the following equation as evidence:

$$CL = \frac{dA_e/dt}{C} \quad (1)$$

The argument is that because in practice, *CL* is sometimes determined from the rate of elimination, it is obvious that the latter influences the former. The proponents of the dependency of the rate of elimination to *CL*, on the other hand, state that the correct physiological presentation of the above equation is:

$$dA_e/dt = CL \times C \quad (2)$$

with a definition of *CL* as the proportionality constant relating the rate of elimination to the blood concentration. This group argues that although in practice *CL* may be estimated using Equation 1, it is the rate of elimination that is dependent on *CL*, not vice versa.

This may seem a matter of semantics or give the impression of the-chicken-and-the egg dilemma. However, neither is true because there is indeed a clear physiologic cause and effect relationships among PK parameters, including clearance and rate of elimination, as demonstrated in the following sections.

It should be noted that the concepts presented here are not new and have been known by pharmacokineticists for many years. However, the aim of this presentation is to highlight the interdependence of PK parameters in an integrated and focused manner with some examples.

Corresponding Author: Reza Mehvar, Ph.D., School of Pharmacy, Texas Tech University Health Sciences Center, 1300 Coulter, Amarillo, TX 79106. reza.mehvar@ttuhsc.edu.

[†] A Listserv maintained by Dr. David Bourne at the University of Oklahoma (<http://www.boomer.org/pkin/>). The message thread was "Clearance and Elimination."

THE RELATIONSHIP BETWEEN CLEARANCE AND RATE OF ELIMINATION

For illustration purposes, we shall select a hypothetical drug with elimination through hepatic metabolism only, although the same principles are valid for drugs with exclusive renal elimination or a combination of renal and hepatic elimination. Figure 1 demonstrates the cellular events in the liver that lead to the metabolism of drugs.




Figure 1: Factors affecting the metabolism of a hypothetical drug by the liver enzymes.

The intrinsic capability of the liver enzymes in the hepatocytes to metabolize the drug (CL_{int}) is dependent on the Michaelis-Menten constants, V_{MAX} (maximum rate of metabolism) and K_M (drug concentration producing half of V_{MAX}), and drug concentration at the site of metabolism (C_u) as defined by the following equation:

$$CL_{int} = \frac{V_{MAX}}{K_M + C_u} \tag{3}$$

For majority of drugs, therapeutic concentrations result in C_u values much lower than K_M , hence for these drugs CL_{int} becomes a constant independent of the blood concentrations within the therapeutic range:

$$CL_{int} \approx \frac{V_{MAX}}{K_M} \tag{4}$$

This is not true for drugs such as phenytoin where therapeutic concentrations are close to the K_M

values. For these drugs, CL_{int} decreases with an increase in the blood concentrations. For simplicity, we shall assume that our drug is metabolized by a single enzyme with a V_{MAX} of 1.5 mg/min and a K_M of 0.1 mg/L. Further, its therapeutic concentration ranges from 1 to 10 $\mu\text{g/L}$ and its free fraction in blood (f_{ub}) is 0.1. This means that at therapeutic concentrations, the free drug concentrations (C_u) (0.1-1 $\mu\text{g/L}$) are much lower than K_M (0.1 mg/L). Therefore, linear metabolism is expected within this range as CL_{int} stays close to 15 L/min:

$$CL_{int} \approx \frac{1.5 \text{ mg/min}}{0.1 \text{ mg/L}} \approx 15 \text{ L/min} \tag{5}$$

It should be noted that this parameter (CL_{int}) is the intrinsic capability of the liver to remove the drug in the absence of any supply (liver blood flow) limitation. However, as demonstrated in Fig. 1, the access of the liver enzymes to the drug is limited by other parameters such as the liver blood flow (Q), free fraction of the drug in the blood (f_{ub}), and permeability of the hepatocytes to the drug. However, in most cases, permeability of the hepatocytes to the drug is not the rate-limiting step in the metabolism of most drugs.

In the following sections, we will look at the metabolism of this hypothetical drug at a macro level by presenting three different scenarios.

Scenario 1: the basics

The metabolism of our hypothetical drug during one pass through the liver is depicted in Fig. 2.




Figure 2: The metabolism of a hypothetical drug in one single pass through the liver based on Scenario 1 (see text for details).

For demonstration purposes, we assume that the concentrations of the drug entering and leaving the liver are 6 and 3 $\mu\text{g/L}$, respectively. This indicates that 50% of the drug entering the liver is converted to a metabolite or, in other words, the fraction of the drug extracted by one single-pass through the liver

(extraction ratio, E) is 0.5. As demonstrated in Fig. 1, assuming high permeability, the extraction ratio (E) is dependent on Q , f_{ub} , and CL_{int} . Several models have been proposed (3-5) to define the relationship between E and its three determinants. One of the widely-used models is the well-stirred or venous equilibrium model, which defines this relationship using the following equation (3):

$$E \approx \frac{f_{ub} \cdot CL_{int}}{Q + f_{ub} \cdot CL_{int}} = \frac{0.1 \times 15}{1.5 + (0.1 \times 15)} = 0.5 \quad (6)$$

Considering that the blood leaving the liver contains a concentration half of that entering the liver (C_{in}) (Fig. 2), one may state that half of the blood is totally cleared of the drug and the other half has the same concentration as C_{in} . In other words, half of the blood is cleared of the drug per unit of time. This is one of the definitions of clearance, which in this case is equivalent to 0.75 L/min (0.5×1.5 L/min), forming the basis of the following equation:

$$CL = Q \times E \quad (7)$$

Equations 6 and 7 clearly indicate both E and CL are dependent on f_{ub} , CL_{int} , and Q . Substituting Equation 4 and 6 into 7 would clearly show the determinants of CL for a drug with linear metabolism:

$$CL = Q \cdot \frac{f_{ub} \cdot \frac{V_{MAX}}{K_M}}{Q + f_{ub} \cdot \frac{V_{MAX}}{K_M}} \quad (8)$$

Any change in Q , f_{ub} , V_{MAX} , and/or K_M may potentially affect the CL of the drug. The degree of dependency of CL on any of these parameters, however, is influenced by the initial values of these parameters relative to each other, and its discussion is outside the scope of this communication.

Now that we have dealt with CL , let us consider the rate of elimination in our example (Fig. 2). Considering a Q of 1.5 L/min and the inlet and outlet concentrations of 6 and 3 $\mu\text{g/L}$, respectively, this means that the rate of elimination (metabolism) of the drug is 4.5 $\mu\text{g/min}$ (1.5 L/min \times 3 $\mu\text{g/L}$). The same value may also be obtained using Equation 2:

$$dAe/dt = CL \times C = 0.75 \text{ L/min} \times 6 \mu\text{g/L} = 4.5 \mu\text{g/min}$$

Scenario 2: a change in the blood concentration

Let us assume that the drug concentration entering the liver is now changed from 6 $\mu\text{g/L}$ to 4 $\mu\text{g/L}$ (Fig. 3). Because of the linear metabolism of the drug, the change in C_{in} is not expected to affect CL_{int} , Q , or f_{ub} of the drug. Therefore, as Equation 6 suggests, the E of the drug (0.5) does not change. Consequently, the outlet concentration will be 2 $\mu\text{g/L}$ in this case (Fig. 3). The CL of the drug, defined as the volume of blood cleared of drug per unit of time, also remains the same as that in Scenario 1 (0.75 L/min, Fig. 3). This is also consistent with the calculation of CL based on Equation 7 or 8, as none of the determinants of CL were changed. However, the rate of elimination of the drug in this case (1.5 L/min \times 2 $\mu\text{g/L}$ or 3 $\mu\text{g/min}$) will be lower than that in Scenario 1 (1.5 L/min \times 3 $\mu\text{g/L}$ or 4.5 $\mu\text{g/min}$). Again, the rate of elimination may also be estimated using Equation 2:




Figure 3: The metabolism of a hypothetical drug in one single pass through the liver based on Scenario 2 (a decrease in the inlet drug concentration; see text for details).

$$dAe/dt = CL \times C = 0.75 \text{ L/min} \times 4 \mu\text{g/L} = 3.0 \mu\text{g/min}$$

This observation indicates that the rate of elimination is dependent on the blood concentration of the drug, whereas CL is independent of blood concentration for a drug with linear pharmacokinetics.

Scenario 3: a change in the clearance

For this scenario, we shall change the CL of the drug and keep the inlet concentration the same as that in Scenario 1. Based on Equation 7, CL is dependent on Q and E . Additionally, E is dependent on Q , f_{ub} , and CL_{int} (Equation 6), with the latter being dependent on V_{MAX} and K_M (Equation 4). A change in any of the determinants of CL , which are Q , f_{ub} , V_{MAX} and K_M (Equation 8), can potentially alter CL . Let us assume that the V_{MAX} of the drug is increased by a factor of 2 from 1.5 mg/min to 3

mg/min as a result of an interacting drug. This in turn results in a two-fold increase in CL_{int} from 15 L/min to 30 L/min:

$$CL_{int} \approx \frac{V_{MAX}}{K_M} \approx \frac{3.0}{0.1} = 30 \text{ L/min}$$

The increased CL_{int} will cause an increase in E and CL as demonstrated in Figure 4 and by Equations 6 and 7:

$$E \approx \frac{f_{ub} \cdot CL_{int}}{Q + f_{ub} \cdot CL_{int}} = \frac{0.1 \times 30}{1.5 + (0.1 \times 30)} = 0.67$$

$$CL = 1.5 \times 0.67 = 1.0 \text{ L/min}$$




Figure 4: The metabolism of a hypothetical drug in one single pass through the liver based on Scenario 3 (an increase in the clearance; see text for details).

Figure 4 also clearly shows that the rate of elimination of the drug in this case is 6 µg/min ($1.5 \text{ L/min} \times 4 \text{ µg/mL}$). As in the previous scenarios, the rate of elimination can also be calculated using Equation 2:

$$dAe/dt = CL \times C = 1.0 \text{ L/min} \times 6 \text{ µg/L} = 6.0 \text{ µg/min}$$

This scenario shows that when CL is changed, the rate of elimination of the drug changes proportionally.

Summary of scenarios

The three scenarios discussed above for a drug with linear pharmacokinetics suggest the following conclusions:

1. When the blood concentration of the drug changes, CL remains unchanged, whereas the rate of elimination of the drug changes proportionally. In other words, the change in the rate of elimination does not affect CL (Scenario 2). This shows the dependency of the rate of elimination to the blood concentration. Further, it shows the independence of CL from

the rate of elimination.

2. When the blood concentration is kept constant and one of the determinants of CL is changed, the change in CL is proportionally reflected in a change in the rate of elimination, indicating the dependency of the rate of elimination on the CL and its determinants (Scenario 3).
3. Consequently, the rate of elimination is dependent on both the CL and blood concentration of the drug. Although CL is not affected by the rate of elimination, it is dependent on its determinants CL_{int} , f_{ub} , and Q .

Coming back to the-chicken-and-the-egg dilemma, it is clear that there is no dilemma regarding the rate of elimination and CL ; it is the rate of elimination that is dependent on CL and not vice versa. Therefore, although both Equations 1 and 2 are mathematically correct, only Equation 2 is physiologically valid. Researchers often use Equation 1 to estimate CL . However, it should not be forgotten that it is CL that determines the rate of elimination.

THE RELATIONSHIPS AMONG OTHER PHARMACOKINETIC PARAMETERS

As the above discussion suggests, CL is a major PK parameter that is related to the efficiency of the eliminating organs, such as the liver and kidneys, to remove the drug from the body. Another major PK parameter, which is at the same level of importance as CL , is the volume of distribution (V), which is an indication of the extent of the distribution of the drug within the body. Similar to CL , V is determined by the physiological parameters of the patient and the physicochemical characteristics of the drug. The volume of distribution of drugs at steady state (V_{SS}) is dependent on the volumes of blood (V_B) and tissue (V_T) and free fractions of the drug in blood (f_{ub}) and tissues (f_{ut}) according to the following equation (6):

$$V_{SS} = V_B + \frac{f_{ub}}{f_{ut}} V_T \tag{9}$$

Both V and CL are independent parameters, meaning that a change in one does not necessarily result in a change in the other parameter, although there may be situations when a change in an underlying physiologic factor would affect both parameters (such as a change in f_{ub}).

A third major PK parameter is the elimination half life ($t_{1/2}$) or rate constant (k). However, it should be noted that, in contrast to CL and V , $t_{1/2}$ or

k does not represent any single physiological process. Instead, it is a composite parameter, reflecting both CL and V processes (2). Therefore, although these three major parameters are mathematically related to each other, one should be aware that $t_{1/2}$ or k is dependent on both CL and V and not vice versa. Therefore, although one may use the mathematical relationship presented in Equation 10 to estimate CL , one should be aware that the proper equations describing the interdependency of these three parameters are presented by Equations 11 and 12:

$$CL = k \cdot V \tag{10}$$

$$k = \frac{CL}{V} \tag{11}$$

$$t_{1/2} = \frac{0.693V}{CL} \tag{12}$$

This issue has been discussed in detail in a recent article (2), hence will not be covered in more detail here.

CONCLUSIONS

The arguments presented here are summarized in Figure 5 in terms of hierarchy of the major pharmacokinetic parameters (constants) and their interdependence. Furthermore, the effects of these kinetic parameters on the blood concentrations and elimination rate after a single intravenous dose are demonstrated.

Although the hypothetical drug used here only underwent hepatic metabolism, the principles discussed here are equally applicable for drugs undergoing renal and/or hepatic elimination. When different organ clearances are involved in the drug elimination, the total CL will be a summation of the individual clearances. This, however, does not change the dependency of the rate of elimination on total CL .




Figure 5: The relationship among physiological and PK parameters for a hypothetical drug eliminated only by hepatic metabolism after administration of a single intravenous bolus dose.

ACKNOWLEDGMENT

The author would like to acknowledge the PharmPK members who participated in the discussion of "Clearance and Elimination," and Dr. David Bourne, who maintains the PharmPK Listserve, for inspiring the author to write this article.

ABBREVIATION LIST

C	Blood Concentration
C_u	Unbound Concentration
C_{in}	Inlet Concentration
C_{out}	Outlet Concentration
CL	Clearance
CL_{int}	Intrinsic Clearance
dA_e/dt	Rate of Elimination
E	Hepatic Extraction Ratio
f_{ub}	Drug Unbound Fraction in Blood
f_{ut}	Drug Unbound Fraction in Tissue
k	Elimination Rate Constant
K_M	Michaelis-Menten Constant
Q	Hepatic Blood Flow
$t_{1/2}$	Elimination Half Life
V	Volume of Distribution
V_B	Blood Volume
V_{MAX}	Maximum Rate of Metabolism
V_{SS}	Volume of Distribution at Steady State
V_T	Tissue Volume

REFERENCES

- [1] Gibaldi M., Perrier D. Pharmacokinetics. New York, NY: Marcel Dekker, 1982.
- [2] Mehvar R., The relationship among pharmacokinetic parameters: effects of altered kinetics on the drug plasma concentration-time profiles. *Am J Pharm Educ*, 68 (2): article 36, 2004.
- [3] Pang K., Rowland M., Hepatic clearance of drugs. I. Theoretical considerations of a "well-stirred" model and a "parallel tube" model. Influence of hepatic blood flow, plasma and blood cell binding, and the hepatocellular enzymatic activity on hepatic drug clearance. *J Pharmacokinet Biopharm*, 5:625-53, 1977.
- [4] Roberts M., Rowland M., Hepatic elimination--dispersion model. *J Pharm Sci*, 74:585-7, 1985.
- [5] Gray M., Tam Y., The series-compartment model for hepatic elimination. *Drug Metab Dispos*, 15:27-31, 1986.
- [6] Mehvar R., Role of protein binding in pharmacokinetics. *Am J Pharm Educ*, 69 (5): article 103, 2005.