

Pharmacy Faculty Articles and Research

School of Pharmacy

2011

Comparative Pharmacodynamics of Ceftobiprole, Daptomycin, Linezolid, Telavancin, Tigecycline, and Vancomycin in the Treatment of Methicillin Resistant *Staphylococcus aureus*: A Monte Carlo Simulation Analysis

Ahmed H. Salem University of Minnesota

Ehab A. Noreddin Hampton University

George G. Zhanel University of Manitoba

Ayman M. Noreddin Chapman University, noreddin@chapman.edu

Follow this and additional works at: https://digitalcommons.chapman.edu/pharmacy_articles

Part of the Pharmacy and Pharmaceutical Sciences Commons

Recommended Citation

Salem A, Noreddin E, Zhanel G, Noreddin A. Comparative pharmacodynamics of ceftobiprole, daptomycin, linezolid, telavancin, tigecycline, and vancomycin in the treatment of methicillin resistant staphylococcus aureus: A monte carlo simulation analysis. *2nd World Congress on Bioavailability & Bioequivalence: Pharmaceutical R & D Summit-2011 and International Conference on Pharmaceutics & Novel Drug Delivery Systems.* 2011. doi: 10.4172/2153-2435.10000S3

This Article is brought to you for free and open access by the School of Pharmacy at Chapman University Digital Commons. It has been accepted for inclusion in Pharmacy Faculty Articles and Research by an authorized administrator of Chapman University Digital Commons. For more information, please contact laughtin@chapman.edu.

Comparative Pharmacodynamics of Ceftobiprole, Daptomycin, Linezolid, Telavancin, Tigecycline, and Vancomycin in the Treatment of Methicillin Resistant *Staphylococcus aureus*: A Monte Carlo Simulation Analysis

Comments

This article was originally published in the proceedings of the 2nd World Congress on Bioavailability & Bioequivalence: Pharmaceutical R & D Summit-2011 and International Conference on Pharmaceutics & Novel Drug Delivery Systems in 2011. DOI: 10.4172/2153-2435.10000S3

Creative Commons License



This work is licensed under a Creative Commons Attribution 3.0 License.

Copyright The authors

Joint Meeting



2nd World Congress on Bioavailability & Bioequivalence: Pharmaceutical R & D Summit-2011

International Conference on Pharmaceutics & Novel Drug Delivery Systems

Comparative pharmacodynamics of Ceftobiprole, Daptomycin, Linezolid, Telavancin, Tigecycline, and Vancomycin in the treatment of Methicillin Resistant Staphylococcus aureus: A Monte Carlo simulation Analysis

Ahmed H. Salem¹, Ehab A. Noreddin³, George G. Zhanel², Ayman M. Noreddin³

¹College of Pharmacy, University of Minnesota, USA

²Faculty of Medicine, University of Manitoba, Canada

³School of Pharmacy, Hampton University, USA

Background/Objectives: Appropriate initial treatment choices for methicillin resistant Staphylococcus aureus (MRSA) infections are very critical. The aim of this study was to compare the ability of Ceftobiprole, Daptomycin, Linezolid, Telavancin, Tigecycline, and Vancomycin to achieve their requisite pharmacokinetic/ pharmacodynamic (PK/PD) target against clinical MRSA isolates.

Methods: Monte Carlo Simulations were performed to simulate the PK/PD indices of the investigated antimicrobials. Population Pharmacokinetic data and Pharmacodynamic indices were integrated into Monte Carlo Simulation routine with 10,000 iterations. Probability of target attainment (PTA) was estimated at MIC values ranging from 0.03-32 μ g/ml to define the PK/PD susceptibility breakpoints. Cumulative fraction of response (CFR) was computed using MIC data from the Canadian National Ward (CAN-Ward) study collected in 2007, 2008 and 2009.

Results: Analysis of the simulation results suggested the breakpoints of 8μ g/ml for Ceftobiprole, 0.12 µg/ml for Daptomycin and Tigecycline, 0.5 µg/ml for Telavancin and 1 µg/ml for Linezolid and Vancomycin. The estimated CFR were 100, 66.5, 84, 89.1, 98.2, 60, 97.5 % for Ceftobiprole, Daptomycin (4mg/kg/day), Daptomycin (6mg/kg/day), Linezolid, Telavancin, Tigecycline, Vancomycin (2gm/day) and Vancomycin (3gm/day), respectively.

Conclusions: Ceftobiprole and Telavancin have the highest probability of achieving favorable outcome against MRSA infections. The susceptibility results suggested a further reduction of the vancomycin breakpoint to $1 \mu g/ml$.