

Summer 8-5-2020

The Poincaré Duality Theorem and its Applications

Natanael Alpay

Chapman University, alpay100@mail.chapman.edu

Melissa Sugimoto

Chapman University, sugim105@mail.chapman.edu

Mihaela Vajiac

Chapman University, mbvajiac@chapman.edu

Follow this and additional works at: <https://digitalcommons.chapman.edu/surf>

 Part of the [Geometry and Topology Commons](#), and the [Other Mathematics Commons](#)

Recommended Citation

Alpay, Natanael; Sugimoto, Melissa; and Vajiac, Mihaela, "The Poincaré Duality Theorem and its Applications" (2020). *SURF Posters and Papers*. 8.

<https://digitalcommons.chapman.edu/surf/8>

This Poster is brought to you for free and open access by the Center for Undergraduate Excellence at Chapman University Digital Commons. It has been accepted for inclusion in SURF Posters and Papers by an authorized administrator of Chapman University Digital Commons. For more information, please contact laughtin@chapman.edu.

The Poincaré Duality Theorem and its Applications

Comments

Presented at the 2020 SURF Virtual Summer Research Conference.



What is a Vector Space

A **vector space** is a set V , with addition and multiplication such that the following holds for all $u, v, w \in V$ and $a, b \in F$: Commutative, Associative, Additive identity, Additive inverse, Multiplicative identity, and the Distribution laws

A **linear map** from V to W is a function $T : V \rightarrow W$ with the following properties for all $u, v \in V$ and $\lambda \in F$:

$$T(u + v) = Tu + Tv, \quad T(\lambda v) = \lambda T(v)$$

An **isomorphism** is an invertible linear map

$\mathcal{L}(V, W)$ is the set of all linear maps from V to W .

A **linear functional** on V is a linear map from V to F , that is element of $\mathcal{L}(V, F)$.

A **dual space** of V , denoted by V^* , is the vector space of all linear functionals on V .

The Five Lemma

A sequence of maps d_0, d_1, \dots, d_n , is an **exact sequence** if

$$Im(d_{k-1}) = Ker(d_k)$$

A **short exact sequence** is of the form:

$$0 \rightarrow A \xrightarrow{f_1} B \xrightarrow{f_2} C \xrightarrow{0}$$

A **long exact sequence** is of f_0, f_1, \dots, f_n , has the form

$$\dots \xrightarrow{f_0} A_0 \xrightarrow{f_1} A_1 \xrightarrow{f_2} A_2 \rightarrow \dots$$

The five lemma Given a commutative diagram of Abelian groups and group homomorphisms as in Figure 1 below, in which the rows are exact sequence, if the maps α, β, δ , and ε are isomorphism, then γ is also an isomorphism.




Figure 1: A commutative diagram to show the Five Lemma

Smooth Manifold

A **diffeomorphism** is a map $f : X \rightarrow Y$ such that f is a homeomorphism, and both f and f^{-1} are smooth (differentiable).

A **smooth manifold of dimension m** is a subset $M \subset \mathbb{R}^n$ such that for each $x \in M$, x has a neighborhood $W \cap M$ that is diffeomorphic to an open subset U of the euclidean space \mathbb{R}^m .

A basis (b_1, \dots, b_n) determines some **orientation** as basis (b'_1, \dots, b'_n) if: $b'_i = \sum_j a_{i,j} b_j$, $det(a_{i,j}) > 0$.

A **oriented smooth manifold** consists of a manifold M and a choice of orientation for each tangent TM_x .

A **good cover** is an open cover $U = \{U_\alpha\}$ of a manifold M of dimension m . M where all nonempty finite intersections $U_{\alpha_0} \cap \dots \cap U_{\alpha_p}$ are diffeomorphic to \mathbb{R}^m .

A **finite good cover** is a good cover U of M which is finite. Equivalently M is **of finite type**.

Outline

In this talk I will explain the duality between the deRham cohomology of a manifold M and the compactly supported cohomology on the same space. This phenomenon is entitled "Poincaré duality" and it describes a general occurrence in differential topology, a duality between spaces of closed, exact differentiable forms on a manifold and their compactly supported counterparts. In order to define and prove this duality I will start with the simple definition of the dual space of a vector space, with the definition of a positive definite inner product on a vector space, then define the concept of a manifold. I will continue with the definition of differential forms on a differentiable manifold and their corresponding spaces necessary to this analysis. I will then introduce the concepts of a good cover of a manifold, manifolds of finite type, and orientation, all necessary concepts towards the goal of defining and proving Poincaré duality. I will finish with the proof of the Poincaré duality in the case of M orientable and admits a finite good cover, with examples.




Figure 2: A smooth manifold

Poincaré duality for deRham cohomology

Lemma 5.6 The two Mayer-Vietoris sequences ... and ..., may be paired together to form a sign-commutative diagram **Theorem.** For an oriented manifold M there is a pairing

$$\int : H^q(M) \otimes H_c^{n-q}(M) \rightarrow \mathbb{R},$$

given by the integral of the wedge product of two forms. Then the Poincaré duality asserts that this pairing is nondegenerate whenever M is orientable and has a finite good cover; equivalently

$$H^q(M) \simeq (H_c^{n-q}(M))^*$$

Proof idea. The proof is a proof by induction as follows:

- Let M be a manifold $M = \bigcup_{k=1}^l U_k$.
- Induction basis: By lemma 5.6 we have $U_1 \cup U_2$.
- Induction Hypothesis: Assume $(U_1 \cup \dots \cup U_k)$.
- Induction Step: $(U_1 \cup \dots \cup U_k) \cup U_{k+1}$
- We have $H^*(U_1 \cup \dots \cup U_k) \cup H^*(U_{k+1}) \implies H^{*+1}((U_1 \dots U_k) \cap U_{k+1}) \implies H^{*+1}((U_1 \dots U_k) \cap U_{k+1})$
- Then by the induction step and the Five Lemma: we get

$$H^{*+1}(U_1 \dots U_{k+1})$$

Other forms of the Poincaré duality

The theorem can be extended to any orientable manifold by the Mayer-Vietoris theorem, as follows:

Theorem. If M is an orientable manifold of dimension n , whose cohomology is not necessarily finite dimension, then

$$H^q(M) \simeq (H_c^{n-q}(M))^*$$

for any integer q .

Proof idea. The finiteness assumption on the good cover is not necessary, then by closer of analysis of topology of a manifold can be extended by the Mayer-Vietoris theorem.

Remark. One should note that the the reverse implication that te following is not always true:

$$H_c^q(M) \simeq (H^{n-q}(M))^*$$

The Euclidian space \mathbb{R}^n

Example. By the Five Lemma if Poincaré duality holds for U, V , and $U \cap V$, then it holds for $U \cup V$. By induction on the cardinality of a good cover. Considering M diffeomorphic to \mathbb{R}^n , and from the Poincaré lemmas

$$H^*(\mathbb{R}^n) = \begin{cases} \mathbb{R} & \text{in dimension } 0 \\ 0 & \text{elsewhere} \end{cases}, H_c^*(\mathbb{R}^n) = \begin{cases} \mathbb{R} & \text{in dimension } n \\ 0 & \text{elsewhere} \end{cases}$$

The Poincaré duality follows.

The Sphere space \mathbb{S}^n

Let S^n are the point $(x_1, \dots, x_{n+1}) \in \mathbb{R}^{n+1}$, such that

$$x_1^2 + \dots + x_{n+1}^2 = 1$$

Example. Let $S^n = U \cup V$ where $U \cap V$ is diffeomorphic to $S^{n-1} \times \mathbb{R}$. Then, through the Mayer-Vietoris sequence,

$$H^*(S^n) = \begin{cases} \mathbb{R} & \text{in dimensions } 0, n \\ 0 & \text{otherwise.} \end{cases}$$

Which can be written as:

$$\begin{aligned} H^0(\mathbb{S}^n) &= \mathbb{R} \\ H^n(\mathbb{S}^n) &= \mathbb{R} \\ H^k(\mathbb{S}^n) &= 0, \quad k \neq 0, n \end{aligned}$$

Hence we have by the Poincaré dual we know $H^q(S^n) \simeq (H^{n-q}(S^n))^*$. For $q = 0$ we have $H^n(S^n) = \mathbb{R}$. For $q = n$, $H^0(S^n) \simeq \mathbb{R}$. And since $\mathbb{R} = \mathbb{R}^*$, we obtain, $H_c^n(S^n) = \mathbb{R}$.

Poincaré duals of a point in \mathbb{R}^n

Since $H^n(\mathbb{R}^n) = 0$, the closed Poincaré dual is μ_p is trivial, and can be represented by any closed n -form on \mathbb{R}^n , but the compact Poincaré dual is the nontrivial class in $H_c^n(\mathbb{R}^n)$ represented by a bump form with total integral 1.

Möbius strip

Counter example. One may suspect that for cohomology with we compact support would have: $H_c^*(E) \simeq H_c^{*-n}(M)$. However this is not generally true; the open Möbius strip which is a vector bundle over S^1 , is a counter example. The compact cohomology of the Möbius strip is identically 0; but S^1 does not match that, hence the Poincaré duality will not hold.




Figure 3: Möbius strip

Example. But if E and M are finite orientable manifolds, and thus the equation would hold using the Poincaré duality (P.D):

$$\begin{aligned} H_c^*(E) &\simeq (H^{m+n-*}(E))^* && \text{By applying the P.D thoerm on } E \\ &\simeq (H^{m+n-*}(M))^* && \text{By deRham cohomology homopoy} \\ &\simeq H_c^{*-n}(M) && \text{By P.D on } M \end{aligned}$$

Conclusion

Poincaré duality describes a general occurrence in differential topology, a duality between spaces of closed, exact differentiable forms on a manifold and their compactly supported counterparts.

$$H^q(M) \simeq (H_c^{n-q}(M))^*$$

The duality between the deRham cohomology of a manifold M and the compactly supported cohomology on the same space.

References

- [1] S. Acler, *Linear Algebra Done Right, third edition.*
- [2] R. Bott, L. W. Tu, *Differential Forms in algebraic Topology*
- [3] J. W. Milnor, *Topology From the Differentiable Viewpoint*