

2007

Synthesis And Antifungal Properties Of Alpha-Methoxy And Alpha-Hydroxyl Substituted 4-Thiatetradecanoic Acids

Nestor Carballeira
University of Puerto Rico

Rosann O'Neill
University of Puerto Rico

Keykavous Parang
Chapman University, parang@chapman.edu

Follow this and additional works at: http://digitalcommons.chapman.edu/pharmacy_articles

 Part of the [Cell Biology Commons](#), [Fungi Commons](#), [Medical Biochemistry Commons](#), and the [Other Chemicals and Drugs Commons](#)

Recommended Citation

Carballeira, Néstor M, Rosann O'Neill, and Keykavous Parang. "Synthesis And Antifungal Properties Of Alpha-Methoxy And Alpha-Hydroxyl Substituted 4-Thiatetradecanoic Acids." *Chemistry And Physics Of Lipids* 150.1 (2007): 82-88
DOI:10.1016/j.chemphyslip.2007.06.218

This Article is brought to you for free and open access by the School of Pharmacy at Chapman University Digital Commons. It has been accepted for inclusion in Pharmacy Faculty Articles and Research by an authorized administrator of Chapman University Digital Commons. For more information, please contact laughtin@chapman.edu.

Synthesis And Antifungal Properties Of Alpha-Methoxy And Alpha-Hydroxyl Substituted 4-Thiatetradecanoic Acids

Comments

NOTICE: this is the author's version of a work that was accepted for publication in *Chemistry And Physics Of Lipids*. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in *Chemistry And Physics Of Lipids*, volume 150, issue 1, 2007. DOI: [10.1016/j.chemphyslip.2007.06.218](https://doi.org/10.1016/j.chemphyslip.2007.06.218)

The Creative Commons license below applies only to this version of the article.

Creative Commons License



This work is licensed under a [Creative Commons Attribution-Noncommercial-No Derivative Works 4.0 License](https://creativecommons.org/licenses/by-nc-nd/4.0/).

Copyright

Elsevier



Published in final edited form as:

Chem Phys Lipids. 2007 November ; 150(1): 82–88.

Synthesis and antifungal properties of α -methoxy and α -hydroxyl substituted 4-thiatetradecanoic acids

Néstor M. Carballeira^{*,1}, Rosann O'Neill¹, and Keykavous Parang²

¹Department of Chemistry, University of Puerto Rico, P.O. Box 23346, San Juan, Puerto Rico 00931-3346

²Department of Biomedical and Pharmaceutical Sciences, College of Pharmacy, University of Rhode Island, Kingston, USA.

Abstract

4-Thiatetradecanoic acid exhibited weak antifungal activities against *Candida albicans* (ATCC 60193), *Cryptococcus neoformans* (ATCC 66031), and *Aspergillus niger* (ATCC 16404) (MIC = 4.8-12.7 mM). It has been demonstrated that α -methoxylation efficiently blocks β -oxidation and significantly improve the antifungal activities of fatty acids. We examined whether antifungal activity of 4-thiatetradecanoic acid can be improved by α -substitution. The unprecedented (\pm)-2-hydroxy-4-thiatetradecanoic acid was synthesized in four steps (20% overall yield), while the (\pm)-2-methoxy-4-thiatetradecanoic acid was synthesized in five steps (14% overall yield) starting from 1-decanethiol. The key step in the synthesis was the hydrolysis of a trimethylsilyloxynitrile. In general, the novel (\pm)-2-methoxy-4-thiatetradecanoic acid displayed significantly higher antifungal activities against *Candida albicans* (ATCC 60193), *Cryptococcus neoformans* (ATCC 66031), and *Aspergillus niger* (ATCC 16404) (MIC = 0.8-1.2 mM), when compared with 4-thiatetradecanoic acid. In the case of *C. neoformans* the (\pm)-2-hydroxy-4-thiatetradecanoic acid was more fungitoxic (MIC = 0.17 mM) than the α -methoxylated analog, but not as effective against *A. niger* (MIC = 5.5 mM). The enhanced fungitoxicity of the (\pm)-2-methoxy-4-thiatetradecanoic acid, as compared to decylthiopropionic acid, might be the result of a longer half-life in the cells due to a blocked β -oxidation pathway which results in more time to exert its toxic effects. Thus, these novel fatty acids may have applications as probes to study fatty acid metabolic routes in human cells.

Keywords

antifungal; decylthiopropionic acid; methoxylated fatty acids; synthesis; thia fatty acids

1. Introduction

Thia fatty acids with the sulfur atom at the 3 and 4 positions have been of considerable interest for their effects on the metabolism of fatty acids (Dyroy et al., 2006; Skrede et al., 1997). The 3-thia fatty acids have been reported to decrease the serum concentration of triacylglycerols and cholesterol and to induce in hepatocytes peroxisomal β -oxidation more efficiently than normal fatty acids (Hovik et al., 1990; Skrede et al., 1995). In particular, tetradecylthioacetic acid is also known to have beneficial antioxidant properties, to lower hyperglycemia, and

*Corresponding author. Tel.: 787-764-0000 x 4791; fax: 787-756-8242 E-mail address: nmcarballeira@uprrp.edu (N.M. Carballeira)

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

displays a positive effect on insulin levels (Dyroy et al., 2006). On the other hand, 4-thia fatty acids inhibit β -oxidation (or it occurs rather slowly) in peroxisomes and mitochondria thereby inducing hepatic lipidosis when administered to rats, i.e., the hepatic levels of triacylglycerols are increased (Hvattum et al., 1992). The difference between the 3- and 4-thia fatty acids can be ascribed as to the position of the sulfur atom in the fatty acyl chain; the 3-thia fatty acids can not be β -oxidized and are rather metabolized by ω -oxidation (Gudbrandsen et al., 2005). On the other hand, the 4-thia fatty acids can be β -oxidized once, but the resulting metabolites (alkylthioacrylyl-CoA) accumulate in the mitochondria thereby inhibiting fatty acid oxidation (Hvattum et al., 1992).

Despite much of the above studies on the metabolism of thia fatty acids in humans nothing is known as to their antifungal properties. However, earlier work by Gordon and collaborators (Kishore et al. 1993) indicated that the myristoyl-CoA:protein *N*-myristoyltransferase (NMT) of the yeast *Saccharomyces cerevisiae*, an enzyme that catalyzes the transfer of myristic acid (C_{14:0}) from myristoyl-CoA to the α -amino nitrogen of glycine, has a greater affinity for 4-thiatetradecanoic acid than for the 3-thiatetradecanoic acid (Kishore et al., 1993). Moreover, when the authors compared human NMT versus *S. cerevisiae* NMT they found that the 4-thiatetradecanoic acid has a greater affinity for the *S. cerevisiae* NMT than for the human NMT. On the other hand, the 3-thiatetradecanoic acid had almost the same affinity for both types of NMTs (Kishore et al., 1993).

Our studies showed that 4-thiatetradecanoic acid has weak antifungal activities against *Candida albicans* (ATCC 60193), *Cryptococcus neoformans* (ATCC 66031), and *Aspergillus niger* (ATCC 16404) (MIC = 4.8-12.7 mM). We have previously shown that α -methoxylation of tetradecanoic acid can significantly improve the antifungal activities of tetradecanoic acid (Carballeira et al., 2005). We investigated whether α -substitution of 4-thiatetradecanoic acid can improve the antifungal activities of 4-thiatetradecanoic acid by inhibition of β -oxidation or fungi NMT.

It was our expectation that the α -methoxy substitution would essentially block the only β -oxidation pathway left for the 4-thia fatty acid thus allowing for only by ω -oxidation to be the most likely catabolic pathway for this novel fatty acid, which could render the α -methoxy 4-thia fatty acid metabolically similar to a 3-thia fatty acid and thus become a good tool for further lipid metabolic studies. Furthermore, based on our previous findings (Carballeira et al., 2005), it was expected that the α -methoxy-4-thia fatty acid to be a better antifungal compound than the parent 4-thia fatty acid. Therefore, in this work we present the first total synthesis for a 2-methoxy-4-thia fatty acid, namely the (\pm)-2-methoxy-4-thiatetradecanoic acid (**1**), and report on its antifungal properties and compare it to other analogs of similar chain lengths.

2. Materials and methods

2.1 Instrumentation

¹H NMR (300 MHz) and ¹³C NMR (75 MHz) were recorded on a Bruker DPX-300 spectrometer. ¹H NMR chemical shifts are reported with respect to internal (CH₃)₄Si (77.0 ppm), ¹³C NMR chemical shifts are reported in parts per million relative to CDCl₃ (77.0 ppm). GC/MS analyses were recorded at 70 eV using a Hewlett Packard 5972A MS equipped with a 30 m \times 0.25 mm special performance capillary column (HP-5MS) of polymethyl siloxane crosslinked with 5% phenyl methylpolysiloxane. High resolution mass spectral data was performed at the Emory University Mass Spectrometry Center on a thermo LTQ-FTMS using flow injection analysis.

2.2. 2-(2-Thiadodecanoyl)-1,3-dioxolane (2)

Into a two necked 100 mL round-bottomed flask equipped with a magnetic stirrer and a condenser was added NaH (0.54g, 23.0 mmol) and diluted with THF (10mL). The solution was kept under argon. The temperature was lowered to 0°C and the 1-decanethiol (4.6 mL, 23.0 mmol) was slowly added to the solution. The reaction mixture was warmed to room temperature in a period of 35 min, then cooled to 0°C and the 2-(2-bromomethyl)-1,3-dioxolane (1.5 mL, 15.0 mmol) was added. The reaction was refluxed for 5 h and extracted with ether (2 × 15 mL), dried over Na₂SO₄, and filtered. The solvent was removed in vacuo affording 3.59 g of 2-(2-thiadodecanoyl)-1,3-dioxolane (2) (92% yield). ¹H NMR (CDCl₃) δ 5.04 (1H, t, *J* = 4.6 Hz, H-2), 4.00 (2H, AA' BB', -OCH₂-), 3.88 (2H, AA' BB', -OCH₂-), 2.70 (2H, d, *J* = 4.6 Hz, H-1'), 2.60 (2H, t, *J* = 7.4 Hz, H-3'), 1.57 (2H, quintet, *J* = 7.4 Hz, H-4'), 1.35 (2H, m, H-5), 1.25 (12H, brs, -CH₂-), 0.86 (3H, t, *J* = 7.0 Hz, -CH₃-); ¹³C NMR (CDCl₃) δ 104.3 (d, C-2), 65.2 (t), 35.4 (t), 32.9 (t), 31.9 (t), 29.6 (t), 29.51 (t), 29.47 (t), 29.3 (t), 29.2 (t), 28.8 (t), 22.6 (t), 14.1 (q, CH₃); GC-MS (70 eV) *m/z* (relative intensity) 260 (*M*⁺, 2), 187 (1), 87 (1), 86 (1), 83 (1), 75 (1), 74 (3), 73 (100), 69 (2), 67 (1), 61 (3), 60 (1), 59 (1), 57 (2), 56 (1), 55 (5); HRMS (ESI) Calcd for C₁₄H₁₇O₂S [M-H]⁺ 259.1726, found 259.1724.

2.3. 3-Thiatetradecanal (3)

Into a 100 mL round-bottomed flask equipped with a magnetic stirrer was added the dioxolane 2 (2.0g, 8.0 mmol) diluted with 20 mL of a solution of water/acetone (1:1). Amberlyst 15® (excess) was added to the solution which was stirred at 60°C for 24h. The reaction mixture was filtered, extracted with ethylacetate (2 × 20 mL) and dried over Na₂SO₄. The solvent was removed in vacuo affording 1.05 g (60 % yield) of 3-thiatetradecanal (3). ¹H NMR (CDCl₃) δ 9.45 (1H, t, *J* = 3.5 Hz, CHO), 3.17 (2H, d, *J* = 3.5 Hz, H-2), 2.42 (2H, t, *J* = 7.4 Hz, H-4), 1.55 (2H, quintet, *J* = 7.4 Hz, H-5), 1.34 (2H, m, H-6), 1.25 (12H, brs, -CH₂-), 0.87 (3H, t, *J* = 6.9 Hz, -CH₃); ¹³C NMR (CDCl₃) δ 193.9 (d, C-1), 41.4 (t), 31.8 (t), 31.6 (t), 30.9 (t), 29.5 (t), 29.4 (t), 29.2 (t), 29.1 (t), 28.6 (t), 22.6 (t), 14.1 (q, CH₃); GC-MS (70 eV) *m/z* (relative intensity) 216 (*M*⁺, 16), 188 (6), 187 (44), 173 (25), 171 (3), 143 (4), 129 (18), 115 (7), 112 (3), 101 (16), 99 (3), 97 (23), 96 (4), 95 (3), 89 (5), 88 (5), 87 (28), 85 (5), 84 (8), 83 (57), 82 (7), 81 (6), 77 (3), 76 (6), 75 (7), 71 (10), 70 (14), 69 (54), 68 (8), 67 (12), 65 (2), 63 (4), 62 (4), 61 (90), 60 (26), 57 (32), 56 (18), 55 (100).

2.4. (±)-2-Trimethylsilyloxy-4-thiatetradecanonitrile (4)

Into a 100 mL round-bottomed flask equipped with a magnetic stirrer was added the 3-thiatetradecanal (3) (1.48 g, 7.0 mmol) diluted in dichloromethane (10 mL). The temperature was lowered to 0°C and trimethylsilyl cyanide (1.19 mL, 9.0 mmol) was added. After 1 min three drops of Et₃N were added and the solution was stirred at 0°C for 3 h. The solution was extracted with ether (2 × 15 mL), dried over Na₂SO₄ and the solvent removed in vacuo. The reaction afforded 1.64 g (74% yield) of nitrile 4 as a viscous oil, which was used for the next step without further purification. ¹H NMR (CDCl₃) δ 4.46 (1H, t, *J* = 6.7 Hz, H-2), 2.84 (2H, d, *J* = 6.7 Hz, H-3), 2.64 (2H, t, *J* = 7.4 Hz, H-5), 1.58 (2H, m, H-6), 1.37 (2H, m), 1.26 (12H, brs, -CH₂-), 0.87 (3H, t, *J* = 6.4 Hz, -CH₃), 0.2 (9H, s, -Si(CH₃)₃); ¹³C NMR (CDCl₃) δ 119.3 (d, C-2), 62.8 (t), 37.5 (t), 33.3 (t), 31.9 (t), 29.6 (t), 29.5 (t), 29.4 (t), 29.3 (t), 29.2 (t), 28.7 (t), 22.7 (t), 14.1 (q), -0.38 (q); GC-MS (70 eV) *m/z* (relative intensity) 315 (*M*⁺, 4), 300 (3), 189 (5), 188 (13), 187 (100), 144 (2), 133 (6), 117 (4), 97 (13), 87 (3), 85 (4), 84 (5), 83(29), 81 (2), 75 (9), 73 (22), 71 (6), 70 (2), 69 (17), 61 (30), 59 (4), 57 (13), 55 (22).

2.5. (±)-2-Hydroxy-4-thiatetradecanoic acid (5)

Into a 50 mL round-bottomed flask equipped with a magnetic stirrer was added the nitrile 4 (0.60 g, 2.0 mmol) diluted in 5.0 mL of concentrated HCl. The solution was stirred overnight at room temperature followed by 2h at 65°C. The reaction mixture was washed with water,

extracted with ether (2 × 10 mL), and dried over Na₂SO₄. The solvent was removed in vacuo affording 0.26 g (50% yield) of (±)-2-hydroxy-4-thiatetradecanoic acid (**5**) as a white solid. mp 84–86°C; ¹H NMR (CDCl₃) δ 4.40 (1H, dd, *J* = 4.2 Hz, *J* = 6.3 Hz, H-2), 3.05 (1H, dd, *J* = 4.2 Hz, *J* = 14.2 Hz, H-3), 2.90 (1H, dd, *J* = 6.3 Hz, *J* = 14.2 Hz, H-3), 2.55 (2H, t, *J* = 7.2 Hz, H-5), 1.58 (2H, quintet, *J* = 7.2 Hz, H-6), 1.35 (2H, m), 1.25 (12H, brs, -CH₂-), 0.87 (3H, t, *J* = 5.9 Hz, -CH₃); ¹³C NMR (CDCl₃) δ 176.6 (s, C-1), 69.8 (d), 36.2 (t), 32.9 (t), 31.9 (t), 29.5 (t), 29.3 (t), 29.2 (t), 28.8 (t), 22.7 (t), 14.1 (q); HRMS (ESI) Calcd for C₁₃H₂₅O₃S [M-H]⁺ 261.1535, found 261.1527.

2.6. Methyl (±)-2-hydroxy-4-thiatetradecanoate (**5a**)

Was prepared from **5** to confirm GC purity by means of the reaction of the acid with methanolic HCl followed by purification using column chromatography on Si gel eluting with hexane/ether (9:1). GC-MS (70 eV) *m/z* (relative intensity) 276 (M⁺, 11), 258 (4), 226 (2), 217 (8), 198 (2), 189 (5), 188 (13), 187 (100), 184 (5), 173 (10), 171 (7), 146 (3), 135 (2), 119 (5), 118 (8), 103 (7), 97 (28), 95 (3), 89 (12), 87 (18), 83 (67), 71 (16), 69 (55), 67 (11), 61 (90), 59 (18), 57 (40), 56 (12), 55 (97).

2.7. (±)-2-Methoxy-4-thiatetradecanoic acid (**1**)

Into a 50 mL round-bottomed flask was placed NaH (0.13 g, 5.0 mmol) diluted in THF (5 mL). Then 2-hydroxy-4-thiatetradecanoic acid (**5**) (0.20 g, 0.76 mmol) in 5 mL of THF, was added to the reaction mixture. The reaction was stirred for 20 min, then cooled to 0°C and iodomethane (0.5 mL, 6.0 mmol) was slowly added. The reaction was stirred for 4 h at room temperature, and then HCl(conc) was added to make the solution acidic. The reaction mixture was extracted with ether (2 × 10 mL), dried over Na₂SO₄, and the solvent was removed in vacuo. The crude product was purified using silica gel column chromatography and eluting first with hexane/EtOAc (9:1) to remove the impurities, followed by hexane/EtOAc (7:3) to obtain 0.15 g (70% yield) of the (±)-2-methoxy-4-thiatetradecanoic acid (**1**) as a viscous oil. ¹H NMR (CDCl₃) δ 3.98 (1H, dd, *J* = 4.4 Hz, *J* = 6.9 Hz, H-2), 3.48 (3H, s, -OCH₃), 2.95 (1H, dd, *J* = 4.4 Hz, *J* = 14.0 Hz, H-3), 2.85 (1H, dd, *J* = 6.9 Hz, *J* = 14.0 Hz, H-3), 2.59 (2H, t, *J* = 7.4 Hz, H-5), 1.55 (2H, quintet, *J* = 7.4 Hz, H-6), 1.34 (2H, m), 1.26 (12H, brs, -CH₂-), 0.87 (3H, t, *J* = 6.4 Hz, -CH₃); ¹³C NMR (CDCl₃) δ 175.9 (s, C-1), 80.7 (d, C-2), 58.7 (q, OCH₃), 34.1 (t), 33.2 (t), 31.9 (t), 29.53 (t), 29.49 (t), 29.3 (t), 29.2 (t), 29.0 (t), 28.8 (t), 22.6 (t), 14.1 (q, CH₃); GC-MS (70 eV) *m/z* (relative intensity) 276 (M⁺, 1), 248 (1), 247 (3), 246 (21), 231 (1), 189 (1), 188 (2), 187 (13), 185 (1), 176 (1), 175 (6), 174 (14), 173 (100), 171 (2), 161 (3), 143 (1), 140 (1), 129 (3), 120 (1), 119 (4), 112 (1), 111 (2), 107 (3), 106 (13), 105 (3), 101 (2), 97 (7), 96 (1), 91 (2), 90 (1), 89 (14), 88 (8), 87 (6), 85 (2), 84 (3), 83 (11), 82 (2), 81 (1), 79 (1), 75 (1), 73 (3), 71 (2), 70 (4), 69 (8), 68 (1), 67 (3), 61 (6), 60 (3), 59 (2), 57 (5), 56 (4), 55 (12). HRMS (ESI) Calcd for C₁₄H₂₇O₃S [M-H]⁺ 275.1675, found 275.1680.

2.8 4-Thiatetradecanoic acid

Was obtained in a 75% yield from the reaction of 3-bromopropionic acid (1g, 6 mmol) and decanethiol (2.4 mL, 12 mmol) using NaH in THF and following the procedure described by Skrede (Skrede et al., 1997).

2.9 Microorganisms

C. albicans ATCC 60193, *A. niger* ATCC 16404, and *C. neoformans* ATCC 66031 were obtained from American Type Culture Collection, Manassas, Virginia. Stock cultures were kept on Sabouraud dextrose agar (SDA; Becton-Dickinson and Co., Sparks, Maryland). Subcultures were prepared on SDA at 35–37°C. Suspension cultures were prepared by inoculation of single colonies in 7 ml of normal saline solution. Prior to preparation of susceptibility assays, yeast cells were resuspended in normal saline to make a transmittance of

73-75% at 530 nm. This transmittance provides a concentration of 10^6 cells/ml for *C. albicans* or *C. neoformans* and 10^6 spores/ml for *A. niger* in saline medium when compared to the control tube. The media was Sabourad dextrose broth (SDB; Becton Dickinson and Co., Sparks, Maryland).

2.10 Chemicals and antifungal agents

Amphotericin B (AMB) was purchased from Acros, New Jersey, USA and was kept as a 0.005 mM stock in DMSO at 0°C and used during one week of preparation. Fluconazole (FLC) was purchased from Medisa Inc., New York, USA, or was provided from Vera Laboratories Ltd, Hyderabad, India, and was kept as a 0.02 mM stock solution at 0°C. Working dilutions were made in SDB medium. The final maximum concentration of DMSO in the assays was 5% (v/v). DMSO was not inhibitory to the organisms in the concentrations tested.

2.11 Susceptibility testing

Microdilutions for control experiments with *C. albicans*, *A. niger*, and *C. neoformans* were the modified method of National Committee for Clinical Laboratory Standards (NCCLS) method as described by Galgani (Galgani, 1993) and by the more recent NCCLS M27-A microdilution methods as described previously (Carballeira et al., 2004; Nam et al., 2004). Dilutions were prepared in 0.1 ml of SDB; the inocula were either 10^4 *C. albicans* or *C. neoformans* cells or *A. niger* spores. The tubes were incubated for 24-28 h at 36 ± 1 °C, and turbidity was read visually. MICs were calculated in comparison to growth control as the lowest concentration that shows inhibition for AMB, FLC, and the test compounds.

3. Results and discussion

A five-step synthesis for the (\pm)-2-methoxy-4-thiatetradecanoic acid (**1**) was accomplished as outlined in Scheme 1. This synthesis started with commercially available 1-decanethiol that was coupled with 2-(2-bromomethyl)-1,3-dioxolane under basic conditions using NaH in dry THF affording the 2-(2-thiadodecanoyl)-1,3-dioxolane (**2**) in a 92% yield. The acetal in dioxolane **2** was removed by treatment with acidic Amberlyst 15® (wet) ion-exchange resin in acetone-water (1:1) for 24h, which afforded the 3-thiatetradecanal (**3**) in a 60% yield. The aldehyde **3** was then reacted with trimethylsilyl cyanide and triethylamine in dichloromethane at 0°C affording the (\pm)-2-trimethylsilyloxy-4-thiatetradecanonitrile (**4**) in a 74% yield. Hydrolysis of the nitrile **4** was carried out in the presence of concentrated HCl by stirring for 24 h at room temperature followed by 2h at 65°C, which resulted in a 50% yield of the desired (\pm)-2-hydroxy-4-thiatetradecanoic acid (**5**). The hydrolysis of the nitrile **4** was the key step in this synthesis. The desired α -methoxylated 4-thia fatty acid **1** was then obtained in a 70% yield in the reaction of the α -hydroxy acid **5** with sodium hydride and iodomethane in tetrahydrofuran (THF) which only afforded methylation at the α -hydroxy group and not at the carboxylic acid. The overall yield for this five-step synthesis was 14%. Our developed synthetic methodology can be applied to synthesize other methoxylated 4-thia fatty acid analogs with different chain lengths by just changing the length of the starting 1-alkanethiol.

The fungitoxicity of the novel α -substituted 4-thia fatty acids **5** and **1**, as compared to 4-thiatetradecanoic acid and other analogs, is presented in Table 1. It is interesting to note that the 4-thiatetradecanoic acid is quite benign (MIC = 5-12 mM) since it is less toxic to the studied fungi than even myristic acid. As compared to the parent 4-thiatetradecanoic acid both α -hydroxylation and α -methoxylation increased the fungitoxicity of the 4-thia fatty acid. In general, α -methoxylation was more effective in increasing the antifungal activity (MIC = 0.8-1.2 mM) of the 4-thiatetradecanoic acid towards all the studied fungal strains (Table 1). However, there were some interesting differences between the strains. For example, in the case of *C. albicans* (ATCC 60193) α -hydroxylation and α -methoxylation were equally effective in

increasing the antifungal activity (MIC = 1.0-1.2 mM) of the 4-thiatetradecanoic acid. However, in the case of *C. neoformans* (ATCC 66031) α -hydroxylation (MIC = 0.17 mM) was about seven fold more effective than α -methoxylation (MIC = 1.24 mM) in increasing the fungitoxicity of the 4-thia fatty acid. In the case of *A. niger* (ATCC 16404) α -hydroxylation did not improve significantly the antifungal activity when compared to the parent analog. However, α -methoxylation enhanced the antifungal activity towards *A. niger* (MIC = 0.8 mM) approximately eight fold higher than that of 4-thiatetradecanoic acid (MIC = 6.3 mM).

In Table 1 the previously reported antifungal activities of (\pm)-2-methoxy-4-oxatetradecanoic acid (Carballeira et al., 2005) have been given to compare the effect of 4-oxa substitution vs. 4-thia substitution on antifungal activities. With respect to *C. neoformans* (ATCC 66031) and *A. niger* (ATCC 16404) the (\pm)-2-methoxy-4-oxatetradecanoic acid was a better antifungal fatty acid (MIC = 0.08-0.22 mM) than **1** (MIC = 0.8-1.2 mM). However, against *C. albicans* (ATCC 60193) the (\pm)-2-methoxy-4-thiatetradecanoic acid (**1**) was about three fold more fungitoxic (MIC = 1.24 mM). Overall, the (\pm)-2-methoxytetradecanoic acid continues to be the most effective antifungal fatty acid of the series (MIC = 0.11 mM) against the three fungal strains tested.

From our results it is clear that α -methoxylation increases the fungitoxicity of 4-thiatetradecanoic acid towards the studied fungal strains. The rationale behind the increased fungitoxicity of **1** versus the 4-thiatetradecanoic acid could be the result of several factors. There are clear differences in the way acid **1** is metabolized by the fungi as compared to 4-thiatetradecanoic acid. The presence of the additional α -methoxy group should introduce sufficient steric effect at the α -position to delay the β -oxidation of acid **1**, thus leaving ω -oxidation and sulfur oxidation as the most likely metabolic pathways for **1**. This steric blockade should increase the half-life of acid **1** in the cell as compared to the half-life of the 4-thiatetradecanoic acid, which can go through one cycle of β -oxidation. Since **1** is expected to have longer half-life in the fungal cells, it possibly gets incorporated more efficiently into the membrane phospholipids of the fungi and causes membrane disruption and eventual antifungal activities (Hvattum et al., 1992). Presumably, sulfur oxidation occurs more rapidly than methoxy hydroxylation (removal of the α -methoxy group) or ω -oxidation (Alterman et al., 1995), and metabolites such as the 2-methoxydecylsulfoxypropionic acid and 2-methoxycarboxymethylsulfoxypropionic acid could also be accumulating in the fungal cells (Hvattum et al., 1992). Alternatively fatty acid **1** or its oxidized metabolites may be inhibiting some key fatty acid metabolic enzymes in the fungal cells or the NMT enzyme of some of these fungi.

As seen in Table 1 there are also clear differences in antifungal activities between the different α -methoxylated myristic acid analogs, in particular between the 2-methoxy-4-oxatetradecanoic acid and **1**. This difference in antifungal activity could also be ascribed to differences in the pKa of the different substrates. It is evident that α -methoxylation increases the pKa of these acids, which could explain the difference between the α -methoxylated fatty acids and the non-methoxylated fatty acids. However, since 4-oxa substitution increases the acidity of myristic acid from 4.91 to 4.37 (Parang et al., 1996) and oxygen is more electronegative than sulfur, this difference in electronegativity could account for the better fungitoxicity of 2-methoxy-4-oxatetradecanoic acid towards *C. neoformans* and *A. niger*. In the case of *C. albicans*, **1** was a better antifungal agent than the corresponding 4-oxa analog. This is not surprising since the 4-oxatetradecanoic acid is known not to be effective against *C. albicans* (Langer et al., 1992).


Thus, in addition to interesting antifungal activities, α -methoxylated fatty acid **1** may be used as a probe to study fatty acid metabolic routes. Further mechanistic and metabolic studies are underway to determine the metabolism of **1**.

Acknowledgements

This work was supported by a grant from the SCORE program of the National Institutes of Health (Grant No. S06GM08102). R. O'Neill thanks the NIH-RISE program for a graduate fellowship. We thank Fred Strobel (Emory University) for the high resolution mass spectral data.

References

- Alterman MA, Chaurasia CS, Lu P, Hanzlik RP. Heteroatom substitution shifts regioselectivity of lauric acid metabolism from omega-hydroxylation to (omega-1)-oxidation. *Biochem. Biophys. Res. Commun* 1995;214:1089–1094. [PubMed: 7575514]
- Carballeira NM, Ortiz D, Parang K, Sardari S. Total synthesis and *in vitro*-antifungal activity of (±)-2-methoxytetradecanoic acid. *Arch. Pharm. Pharm. Med. Chem* 2004;337:152–155.
- Carballeira NM, O'Neill R, Parang K. Racemic and optically active 2-methoxy-4-oxatetradecanoic acids: novel synthetic fatty acids with selective antifungal properties. *Chem. Phys. Lipids* 2005;136:47–54. [PubMed: 15899476]
- Dyroy E, Wergedahl H, Skorve J, Gudbrandsen OA, Songstad J, Berge RK. Thia fatty acids with the sulfur atom in even or odd positions have opposite effects on fatty acid catabolism. *Lipids* 2006;41:169–177. [PubMed: 17707983]
- Galgiani JN. Susceptibility testing of fungi: current status of the standardization process. *Antimicrob. Agents Chemother* 1993;37:2517–2521. [PubMed: 8109912]
- Gudbrandsen OA, Dyroy E, Bohov P, Skorve J, Berge RK. The metabolic effects of thia fatty acids in rat liver depend on the position of the sulfur atom. *Chem. Biol. Interact* 2005;155:71–81. [PubMed: 15949791]
- Hovik R, Osmundsen H, Berge R, Aarsland A, Bergseth S, Bremer J. Effects of thia-substituted fatty acids on mitochondrial and peroxisomal β -oxidation. *Biochem. J* 1990;270:167–173. [PubMed: 2396976]
- Hvattum E, Skrede S, Bremer J, Solbakken M. The metabolism of tetradecylthiopropionic acid, a 4-thia stearic acid, in the rat. *Biochem. J* 1992;286:879–887. [PubMed: 1417748]
- Kishore NS, Wood DC, Mehta PP, Wade AC, Lu T, Gokel GW, Gordon JI. Comparison of the acyl chain specificities of human myristoyl-CoA synthetase and human myristoyl-CoA:protein *N*-myristoyltransferase. *J. Biol. Chem* 1993;268:4889–4902. [PubMed: 8444867]
- Langer CA, Lodge JA, Travis SJ, Caldwell JE, Lu T, Li Q, Bryant ML, Devadas B, Gokel GW, Kobayashi GS, Gordon JI. 4-Oxatetradecanoic acid is fungicidal for *Cryptococcus neoformans* and inhibits replication of human immunodeficiency virus I. *J. Biol. Chem* 1992;267:17159–17169. [PubMed: 1512254]
- Nam NH, Sardari S, Selecky M, Parang K. Carboxylic acids and phosphate ester derivatives of fluconazole: synthesis and antifungal activities. *Bioorg. Med. Chem* 2004;12:6255–6269. [PubMed: 15519168]
- Parang K, Knauss EE, Wiebe LI, Sardari S, Daneshlab M, Csizmadia F. Synthesis and antifungal activities of myristic acid analogs. *Arch. Pharm. Pharm. Med. Chem* 1996;329:475–482.
- Skrede S, Wu P, Osmundsen H. Effects of tetradecylthiopropionic acid and tetradecylthioacrylic acid on rat liver lipid metabolism. *Biochem. J* 1995;305:591–597. [PubMed: 7832778]
- Skrede S, Sørensen HN, Larsen LN, Steineger HH, Høvik K, Spydevold ØS, Horn R, Bremer J. Thia fatty acids, metabolism and metabolic effects. *Biochim. Biophys. Acta* 1997;1344:115–131. [PubMed: 9030189]



i) NaH, 2-(2-bromomethyl)-1,3-dioxolane, THF, 0°C, 5h; *ii*) Amberlyst, acetone-H₂O (1:1), 60°C, 24h; *iii*) trimethylsilyl cyanide/Et₃N, CH₂Cl₂, 0°C, 3h; *iv*) HCl (conc.), 65°C, 24h; *v*) NaH, CH₃I, THF, rt, 4h.

Scheme 1.

Synthesis of (±)-2-hydroxy-4-thiatetradecanoic acid (**5**) and (±)-2-methoxy-4-thiatetradecanoic acid (**1**).

Table 1

Antifungal activity (MIC values, mM)^a against *Candida albicans* (SDB), *Cryptococcus neoformans* (SDB), and *Aspergillus niger* (SDB) at 35-37°C after 24-48 h

Compound	<i>C. albicans</i> ATCC 60193	<i>C. neoformans</i> ATCC 66031	<i>A. niger</i> ATCC 16404
14:0 ^b	3.80 ^c	1.90	3.80
4-thia-14:0	12.66	4.75	6.33
2-OH-4-thia-14:0	1.04	0.17	5.53
2-OMe-4-thia-14:0	1.24	1.24	0.83
2-OMe-14:0 ^b	0.11 ^c	0.11	0.11
2-OMe-4-oxa-14:0 ^b	3.50	0.08	0.22
FLC	>1.0	0.002	1.50
AMB	<0.0003	<0.0003	<0.0009
DMSO	>5	>5	>5

^aThe result is the average of three separate experiments. The upper limit of the standard error of the mean (SEM) was $\pm 10\%$.

^bAntifungal data taken from Carballeira et al., 2005.

^cThese MIC values were measured against *C. albicans* 14053, also taken from Carballeira et al., 2005.