

2015

The Effect of Δ^9 -tetrahydrocannabinol, Cannabidiol, Menthol and Propofol on 5-hydroxytryptamine Type 3 Receptors--A Computational Approach

Andreas C. Schilbach


Tatiana Prytkova

Chapman University, prytkova@chapman.edu

Keun-Hang Susan Yang

Chapman University, kyang@chapman.edu

Follow this and additional works at: http://digitalcommons.chapman.edu/scs_articles

 Part of the [Neurosciences Commons](#), and the [Other Pharmacy and Pharmaceutical Sciences Commons](#)

Recommended Citation

Schilbach et al.: The effect of Δ^9 -tetrahydrocannabinol, cannabidiol, menthol and propofol on 5-hydroxytryptamine type 3 receptors—a computational approach. *BMC Neuroscience* 2015 16(Suppl 1):P290.
doi: 10.1186/1471-2202-16-S1-P290

This Article is brought to you for free and open access by the Science and Technology Faculty Articles and Research at Chapman University Digital Commons. It has been accepted for inclusion in Mathematics, Physics, and Computer Science Faculty Articles and Research by an authorized administrator of Chapman University Digital Commons. For more information, please contact laughtin@chapman.edu.

The Effect of $\Delta 9$ -tetrahydrocannabinol, Cannabidiol, Menthol and Propofol on 5-hydroxytryptamine Type 3 Receptors--A Computational Approach

Comments

This article was originally published in *BMC Neuroscience*, volume 16, supplement 1, in 2015. DOI: [10.1186/1471-2202-16-S1-P290](https://doi.org/10.1186/1471-2202-16-S1-P290)

Creative Commons License

Creative

Commons

This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

License

Copyright

The authors

POSTER PRESENTATION

Open Access

The effect of Δ^9 -tetrahydrocannabinol, cannabidiol, menthol and propofol on 5-hydroxytryptamine type 3 receptors—a computational approach

Andreas C Schilbach*, Tatiana Prytkova, Susan Keun-Hang Yang

From 24th Annual Computational Neuroscience Meeting: CNS*2015
Prague, Czech Republic. 18-23 July 2015

This study investigates the function of 5-HT type 3 (5-HT₃) receptors using a computational approach. Antagonists of the 5-HT₃ receptor are currently one of the most effective therapeutic agents in treatment of chemotherapy-induced nausea, vomiting, and irritable bowel syndrome. Several experimental studies have shown the effect of pharmacological agents such as 9-tetrahydrocannabinol (THC), the psychoactive component of Cannabis, Cannabidiol (CBD), a non-psychoactive ingredient of Cannabis plant, Menthol, Propofol, and etc. on the functional human 5-HT₃ receptors expressed in *Xenopus* oocyte as well as rat nodose ganglion neurons [1,2]. 5-HT evoked currents recorded by a two-electrode clamp technique were inhibited by ligands in a concentration dependent manner. Simulations of allosteric inhibition were modeled using Vina docking techniques with the 5-HT₃ structure (see Tables 1 and 2 for results). The 5-HT₃ structure was found using homology sequence similarity techniques with the neuronal nicotinic acetylcholine receptor (nACh) and inhibitory neurotransmitter receptor for GABA(A). Results of studies with other members of the superfamily of ligand gated ion channels signified key residues involved in ligand binding sites within the transmembrane region of 5-HT₃ [3]. Flexible and rigid docking simulations around key residues resulted in a number of low-energy (high affinity) configurations of ligand binding (Figure 2). The predicted residues TYR and THR may constitute a naturally occurring binding site for 5-HT₃.

Table 1 Docking of Ligands on lower (THR361) binding sites of 5-HT₃A

Compound - (THR361 binding site)	Binding energy (kcal/mol)
Cyclohexane	-4.5
Eucalyptol	-4.5
Menthol	-5.6
Propofol	-6.0

Conclusion

Experimental inhibition of 5-HT₃ shows similar trend in computational binding energies to the lower binding site (THR 361). Docking calculations provide explanation of molecular basis of difference in inhibition by menthol like compounds. Similar binding energies for THC and CBD corresponds to their similar inhibition of membrane currents measured in experiment.

Published: 18 December 2015

References

1. Yang KHS, Isaev D, Morales M, Petroianu G, Galadari S, Oz M: **The effect of Δ^9 -tetrahydrocannabinol on 5-HT₃ receptors depends on the current density.** *Neuroscience* 2010, **171**:40-49.
2. Ashoor A, Nordman JC, Veltri D, Yang KHS, Shuba Y, Kury LA, Bassem Sadek B, Howarth FC, Shehu A, Kabbani N, Oz M: **Menthol inhibits 5-HT₃ receptor-mediated currents.** *J Pharmacol Exp Ther* 2013, **347**(2):398-409.
3. Trattinnig SM, Harpsøe K, Thygesen SB, Rahr LM, Ahring PK, Balle T: **Discovery of a novel allosteric modulator of 5-HT₃ receptor: Inhibition and potentiation of Cys-loop receptor signaling through a conserved transmembrane intersubunit site.** *Journal of Biological Chemistry* 2012, **287**(30):25241-25254.

* Correspondence: schil105@mail.chapman.edu
Schmid College of Science and Technology, Chapman University, Orange
CA, 92866, USA

Table 2 Docking of Ligands on upper (TYR346) binding sites of 5-HT3A

Compound - (TYR346 binding site)	Rigid Docking - Binding energy (kcal/mol)	Flexible Docking - Binding energy (kcal/mol)
Δ^9 -tetrahydrocannabinol (THC)	-7.0	-9.2
Cannabidiol (CBD)	-7.0	-8.8




Figure 2 Interaction diagrams of Propofol with lower binding site (THR361) of 5-HT3A. A. 2-D Ligand interaction diagram. Residues are annotated with their 3-letter amino acid code, and their position classification. Hydrophilic interactions: include the hydroxyl group with residue Asp266 and Met263 (Blue). Hydrophobic interactions: include the hydroxyl group with residue Asp266 and Met263 (Blue). Hydrophobic interactions: include the hydroxyl group with residue Asp266 and Met263 (Blue). Hydrophobic interactions: include the hydroxyl group with residue Asp266 and Met263 (Blue).




Figure 1 Experimental comparison [3] of the effects of compounds structurally related to menthol on the 5-HT3 receptors.

doi:10.1186/1471-2202-16-S1-P290

Cite this article as: Schilbach *et al.*: The effect of Δ^9 -tetrahydrocannabinol, cannabidiol, menthol and propofol on 5-hydroxytryptamine type 3 receptors—a computational approach. *BMC Neuroscience* 2015 16(Suppl 1):P290.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

