

Chapman University

Chapman University Digital Commons

Biology, Chemistry, and Environmental Sciences
Faculty Articles and Research

Science and Technology Faculty Articles and
Research

6-5-2018

Codon Usage Revisited: Lack of Correlation Between Codon Usage and the Number of tRNA Genes in Enterobacteria

Joaquín Rojas
Universidad de Chile

Gabriel Castillo
Universidad de Chile

Lorenzo Eugenio Leiva
Universidad de Chile

Sara Elgamal
The Ohio State University

Omar Orellana
Universidad de Chile

See next page for additional authors

Follow this and additional works at: https://digitalcommons.chapman.edu/sees_articles



Part of the [Amino Acids, Peptides, and Proteins Commons](#), [Biochemistry Commons](#), [Cellular and Molecular Physiology Commons](#), [Molecular Biology Commons](#), [Nucleic Acids, Nucleotides, and Nucleosides Commons](#), and the [Other Biochemistry, Biophysics, and Structural Biology Commons](#)

Recommended Citation

Rojas, J., Castillo, G., Leiva, L.E., Elgamal, S., Orellana, O., Ibba, M. and Katz, A. (2018) Codon usage revisited: Lack of correlation between codon usage and the number of tRNA genes in enterobacteria. *Bioch. Biophys. Res. Comm.* 502, 450-455. <https://doi.org10.1016/j.bbrc.2018.05.168>

This Article is brought to you for free and open access by the Science and Technology Faculty Articles and Research at Chapman University Digital Commons. It has been accepted for inclusion in Biology, Chemistry, and Environmental Sciences Faculty Articles and Research by an authorized administrator of Chapman University Digital Commons. For more information, please contact laughtin@chapman.edu.

Codon Usage Revisited: Lack of Correlation Between Codon Usage and the Number of tRNA Genes in Enterobacteria

Comments

NOTICE: this is the author's version of a work that was accepted for publication in *Biochemical and Biophysical Research Communications*. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in *Biochemical and Biophysical Research Communications*, volume 502, in 2018. <https://doi.org/10.1016/j.bbrc.2018.05.168>

The Creative Commons license below applies only to this version of the article.

Creative Commons License



This work is licensed under a [Creative Commons Attribution-Noncommercial-No Derivative Works 4.0 License](https://creativecommons.org/licenses/by-nc-nd/4.0/).

Copyright

Elsevier

Authors

Joaquín Rojas, Gabriel Castillo, Lorenzo Eugenio Leiva, Sara Elgamal, Omar Orellana, Michael Ibba, and Assaf Katz



HHS Public Access

Author manuscript

Biochem Biophys Res Commun. Author manuscript; available in PMC 2019 August 25.

Published in final edited form as:

Biochem Biophys Res Commun. 2018 August 25; 502(4): 450–455. doi:10.1016/j.bbrc.2018.05.168.

Codon usage revisited: Lack of correlation between codon usage and the number of tRNA genes in enterobacteria

Joaquín Rojas^a, Gabriel Castillo^a, Lorenzo Eugenio Leiva^a, Sara Elgamal^b, Omar Orellana^a, Michael Ibba^b, and Assaf Katz^{a,*}

^aPrograma de Biología Celular y Molecular, ICBM, Facultad de Medicina, Universidad de Chile, Santiago 8380453, Chile

^bDepartment of Microbiology and The Center for RNA Biology, Ohio State University, Columbus, Ohio 43210, USA

Abstract

It is widely believed that if a high number of genes are found for any tRNA in a rapidly replicating bacteria, then the cytoplasmic levels of that tRNA will be high and an open reading frame containing a higher frequency of the complementary codon will be translated faster. This idea is based on correlations between the number of tRNA genes, tRNA concentration and the frequency of codon usage observed in a limited number of strains as well as from the fact that artificially changing the number of tRNA genes alters translation efficiency and consequently the amount of properly folded protein synthesized. tRNA gene number may greatly vary in a genome due to duplications, deletions and lateral transfer which in turn would alter the levels and functionality of many proteins. Such changes are potentially deleterious for fitness and as a result it is expected that changes in tRNA gene numbers should be accompanied by a modification of the frequency of codon usage. In contrast to this model, when comparing the number of tRNA genes and the frequency of codon usage of several *Salmonella enterica* and *Escherichia coli* strains we found that changes in the number of tRNA genes are not correlated to changes in codon usage. Furthermore, these changes are not correlated with a change in the efficiency of codon translation. These results suggest that once a genome gains or loses tRNA genes, it responds by modulating the concentrations of tRNAs rather than modifying its frequency of codon usage.

Keywords

tRNA; codon usage; efficiency of translation; enterobacteria; *Salmonella enterica*; *Escherichia coli*

*To whom correspondence should be addressed. Assaf Katz, Programa de Biología Celular y Molecular, ICBM, Facultad de Medicina, Universidad de Chile, Santiago 8380453, Chile, Tel.: +56-2 29789584, akatz@med.uchile.cl.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Introduction

Proteins, which are essential for the physiology of all organisms, are coded in nucleic acids. In order to translate the genetic message contained in nucleic acids into a polypeptide, organisms use a code where each three contiguous nucleotides (a codon) are translated to a specific amino acid. tRNAs are essential for this process. In one side of the folded tRNAs there is a loop that contains a three nucleotide sequence called the anticodon, that can interact specifically with the complementary codons in mRNAs during translation. The 3'-extreme of the tRNA is able to carry an amino acid that can be transferred to a nascent peptide. This reaction is catalyzed by the ribosome that additionally ensures a correct matching between the codon in the mRNA and the anticodon in tRNA [1–3].

Although there are some exceptions, in most organisms and growth conditions each codon codes for a single amino acid with high precision [4–7]. Nevertheless, the standard genetic code is redundant with 61 codons that code for only 20 canonical amino acids. Thus, while Met and Trp are coded by a single codon, Arg, Leu and Ser are coded by as many as 6 different codons [8]. Thanks to wobble interactions -where the ribosome allows interactions between non Watson-Crick base pairs at the third codon position- a tRNA can recognize several codons. This allows one tRNA to decode several codons (coding for the same amino acid) and also a single codon to be decoded by several tRNAs (carrying the same amino acid). There has been a long debate regarding the role of such redundancy in protein synthesis. One of the most accepted ideas is that different codons coding for the same amino acid will be translated at a different speed. Thus, highly expressed proteins will require their genes to be coded mostly by codons that are efficiently translated which are expected to correlate with high cellular levels of the corresponding tRNAs. This idea is based on the fact that translation elongation speed depends on the concentration of aa-tRNAs [8]. Also, it has been shown that in rapidly replicating organisms, there is a positive correlation between tRNA concentration (or the number of tRNA genes, see below) and the usage of the codons they decode in highly expressed genes [8,9]. Consistent with gene copy number effects, artificially changing the concentration of tRNAs in an organism [10] or the frequency of codons in a gene [11,12] alters gene expression and cell fitness [13]. Additionally, usage of infrequent codons that are expected to be translated at slower rates has been associated with modulation of processes coupled to translation such as protein folding or secretion [3,8,11,14]. In fact, changing codon usage in genes may induce the production of incorrectly folded proteins [3,11,14].

Bacteria can easily obtain genetic material through lateral transfer, a process that is essential for a rapid adaptation of the genome to new environments. Many of the mechanisms involved in the transfer of genetic material involve the transfer of tRNA genes. For instance, viral genomes, genomic islands and plasmids have been shown to carry one or several tRNA genes [15–18]. Internal recombination of different areas of a genome can also duplicate or produce the loss of genes coding for tRNAs [18]. Thus, the number of genes coding for tRNAs in bacterial species can change in a single generation. Concentration of tRNAs has been shown to be correlated to the number of genes coding for them, at least in bacteria such as *Escherichia coli* [19,20]. Thus, changes in the number of tRNA genes are expected to alter the cellular concentration of the corresponding tRNAs. As discussed above, this is

expected to modify the speed of translation of some of the codons altering the levels and functionality of at least part of the proteome.

McDonald *et al.* have shown that changes in the number of tRNA genes derived from lateral transfer events correlate with codon usage of accompanying genes acquired by such events in genomes of *Escherichia coli* and *Shigella*. Potentially, these correlations allow a more efficient translation of the newly acquired genes [18]. We have found previously a similar trend while analyzing codon usage and tRNA gene content of an integrative conjugative element present in the genome of the chemolithoautotrophic bacterium *Acidithiobacillus ferrooxidans* [15,21]. Nevertheless, recent reports have shown that most tRNAs in this mobile element are expressed at very low levels [22], questioning their ability to improve gene expression. If it is common that changes in tRNA gene copy numbers produce only small changes in tRNA levels, then we would expect these have minor effects on gene expression and, consequently, on the frequency of codon usage in most genes of the genome. To test this hypothesis, we studied the relationship between the number of tRNA genes and the frequency of codon usage in genomes of enterobacteria. We have selected as models two well studied enterobacteria, *Escherichia coli* and *Salmonella enterica*. Our results indicate that changes in the number of genes coding for tRNAs have only minor effects on codon usage of both the whole genome and highly expressed genes. Correspondingly, fusions of codons translated by these tRNAs to *gfp*, have little effect on its translation efficiency. These results suggest that expression of tRNA genes acquired by lateral transfer in enterobacteria is rapidly adapted to the requirements of the host genome.

Materials and methods

Selection of analyzed genomes

RefSeq versions of genomes from *E. coli* and *Salmonella* were downloaded from NCBI ftp site. Only genomes annotated as being at “Chromosome” or “Complete Genome” levels were used for further analyses. In order to reduce excessive sequence redundancy, SNP based phylogenetic trees were constructed using kSNP3 [23]. Based on these trees, a single strain was randomly selected from each group of similar strains. For example, only one genome was retained from ~30 genomes that grouped with *E. coli* K12 strains (most of which were annotated as K12 strains or strains derived from it). Using this selection method we retained 206 strains of *E. coli* and 139 strains of *S. enterica* (Supplementary file 1).

Counts of codons and tRNA genes

To prevent differences in tRNA number arising from potentially different criteria used during genome annotation, tRNA genes were re-annotated using tRNAScan-SE 2.0 [24]. Frequency of usage of each codon from each gene was calculated using in home written Perl 5 scripts. Then, average and standard deviation values were calculated from the complete set of codon usage values for each annotated gene in the genome. Additionally, a similar calculation was performed using only genes of 5 highly expressed genes selected based on a report by Karlin *et al.*, 2001 [25]. Selected genes were *rpIL* (50S ribosomal subunit protein L7/L12), *rpsA* (30S ribosomal subunit protein S1), *rpsF* (30S ribosomal subunit protein S6) *groEL* (60 kDa chaperonin) and *eno* (enolase). The *rpsF* gene was absent in one of the

selected *E. coli* strains (RefSeq_assembly_accession: GCF_000013305.1), so this genome was eliminated from codon usage analyzes of highly expressed genes.

Determination of codon translation efficiency

Plasmids carrying fusions of Glu and Ala codons were constructed as indicated in supplementary methods and transformed in *S. typhimurium* ATCC strain 14028, designated 14028s, *S. enteritidis* PT4 NCTC 13349 [26] and *S. typhi* STH2370 [27]. M9 media (47.7mM Na₂HPO₄, 22.0mM KH₂PO₄, 8.6mM NaCl, 18.7mM NH₄Cl, 2mM MgSO₄, 0.1mM CaCl₂, 0.4% Glycerol) supplemented with 50μM L-Cys, 50μM L-Trp, MgCl₂ and 100μg/ml ampicillin were inoculated with each strain from a saturated overnight culture (20h in M9 media supplemented with 0.1% tryptone and 100μg/ml ampicilin) and grown at 37°C in an orbital shaker. When bacteria reached mid-log phase (OD₆₀₀ ~0.4–0.6), 50μl aliquots were used to inoculate fresh 150μl of media supplemented with arabinose (0.4% final concentration) in a 96-well optical-bottom plate. Plates were further shaken for 1.5 hours at 37°C. Then, OD₆₀₀ and fluorescence intensity of GFP (Ex. 480±4.5nm, Em. 515±10nm) and mCherry (Ex. 555±4.5nm, Em. 600±10nm) were measured in a microplate reader (INFINITE M200PRO, TECAN).

Results

In order to compare the number of tRNA genes with the frequency of codon usage, complete sequences of genomes from *Escherichia coli* and *Salmonella enterica*, two very well characterized representatives of the enterobacteriaceae family, were obtained from NCBI and further analyzed as specified above to determine the number of tRNA codons and the frequency of usage of each codon. While the number of genes coding for tRNAs with anticodons such as Ala-UGC and Arg-UCU in *E. coli* or Ala-GGC and Arg-UCU *S. enterica* was variable between strains, others tend to be coded by a single gene copy number (eg Cys-GCA was coded by a single gene in all analyzed strains) (Figure 1 and S1). Interestingly, the group of tRNAs that are coded by a variable number of genes is different between both species. Compared to the variability found in the number of tRNA genes, the frequency of codon usage was much more stable. This is observed for both species as well as when analyzing either the frequency of codon usage of the full genome or for a selected group of highly expressed genes (Figures 1 and S1). The fact that the frequency of codon usage remains constant while the number of genes coding for the corresponding tRNAs varies, suggests that the correlation between the number of tRNA genes and the codon usage frequency is not as strong as usually expected.

Changes in the number of tRNA genes are not correlated to modifications in the frequency of codon usage

We compared the number of tRNA genes in each strain with the frequency of usage of all codons coding for the corresponding amino acid in a subset of highly expressed genes, where we expect a higher correlation between the number of tRNA genes and frequency of codon usage. A graph for each possible tRNA anticodon was produced. As observed in figure 2 for tRNA^{Glu}_{UUC}, tRNA^{Ala}_{GGC} and tRNA^{Ala}_{UGC} in *S. enterica* the frequency of codon usage is completely independent from the number of genes coding for the tRNAs

decoding those codons or those decoding the other codons for the same amino acid. Similar results were obtained for all other tRNAs in either *S. enterica* or *E. coli* (Figures S2, S3). Also, similar results were obtained analyzing the full genome (Data not shown). Thus, our data indicate that when the number of tRNA genes vary in a genome there is no effect on the frequency of codon usage. This observation does not directly imply an absence of correlation between both values. In fact, when we compare the average codon usage with the average number of tRNA genes reading the corresponding codons by direct Watson-Crick complementarity, we observe a positive correlation similar to what has been previously observed (Figure S4). Nevertheless, as mentioned previously, variability in the number of tRNA genes is much wider than observed for the frequency of codon usage.

Changes in the number of tRNA genes are not correlated to changes in the efficiency of codon translation

Changes in the number of tRNA genes are usually expected to modify the concentration of the corresponding tRNAs and as a result also alter the efficiency of codon translation. Due to the broad spectrum of effects that this would produce, it is expected that changes in the number of tRNA genes would constitute a selective pressure to change the frequency of codon usage. As this is not observed, we hypothesize there is no significant change in translation efficiency in strains with altered tRNA gene copy numbers. To test this hypothesis we constructed reporters of codon translation efficiencies by introducing 4 contiguous identical codons near the 5' extreme of the green fluorescent protein gene (*gfp*). In addition to *gfp*, a gene coding for a red fluorescent protein (mCherry) was introduced as a transcriptional fusion to normalize for potential changes in the mRNA levels. Thus, changes in the efficiency of translation of a codon are expected to alter the ratio between fluorescence of GFP and mCherry. Reporters were constructed for codons coding for Glu (GAG and GAA) and Ala (GCG, GCA, GCT, GCC) and tested in three *S. enterica* strains that have different numbers of genes for tRNAs decoding those codons (Table 1). We observed a correlation between the GFP/mCherry ratios of Glu codons and their frequency of usage in highly expressed genes (Figure 3A). Nevertheless, this correlation was not observed for Ala codons, a result in accordance with previous reports showing a similar translation efficiency for all codons coding diverse amino acids [10]. Given that the number of tRNAs that decode each Ala codon are different, this result suggests that efficiency of translation is not correlated with the number of tRNA genes. Furthermore, when the fluorescence ratios are normalized by the ratio of one of the reporters it is evident that relative efficiencies of codon translation are equivalent in the three *S. enterica* strains (Figure 3B and C), although *S. Typhi* has a much lower number of genes coding for tRNA^{Glu}_{UUC} and tRNA^{Ala}_{UGC} than the other strains, supporting the idea that under exponential growth the number of tRNA genes is not affecting codon translation efficiencies.

Discussion

Sudden changes in cellular levels of tRNAs such as those expected by alteration in tRNA gene numbers due to lateral transfer of genes, gene duplication and deletions or mutations of anticodons should produce global changes in the concentration and folding of proteins. Nevertheless, in this work we have observed that changes in the number of tRNA genes is

not correlated to alterations in the frequency of usage or the translation efficiency of the corresponding codons in enterobacteria. Thus, codon usage and translation efficiency seem to be evolutionary robust, while the tRNA gene copy number is apparently less constrained. This is in agreement with a previous report by Satapathy *et al.* who have questioned the role of tRNA gene numbers in selection of codon bias, based on the absence of correlation between the number of tRNA genes and enrichment of C vs U ending codons in the case of 4 amino acids coded by two codons translated by a single tRNA isoacceptor [28]. Probable mechanisms allowing this stability in codon usage in organisms where tRNA gene numbers vary, are silencing of acquired tRNA genes as has been observed in *A. ferrooxidans* [22] or regulation of either transcription or stability of tRNAs which has also been observed under a limited number of experimental conditions [29,30]. Both mechanisms are expected to occur much faster than changes in codons usage, which would require a much larger number of mutations [31]. The fact that genes in mobile elements are enriched for codons decoded by tRNAs contained in these elements [18,21] might be explained by either 1) codon usage of acquired protein coding genes being adapted to genomic tRNA availability slower than silencing of acquired tRNA genes, 2) retaining the original codon usage of acquired genes being advantageous to recipient cells because it reduces the expression of foreign genes that could reduce cellular fitness, or 3) both the tRNA and the protein genes only being expressed in particular conditions, thus reducing the pressure to adapt the codon usage of the rest of the genome.

In any case, based on our observations the estimation of translation efficiency using parameters based on codon usage of highly expressed genes (such as CAI [32]) seems to be more reliable than those based on the count of tRNA genes (such as TAI [33]). Still, caution should be taken when using such parameters as under several culture conditions levels of tRNAs [29,30], their aminoacylation [34], or levels of chemical modification [35] can be altered, changing the efficiency of translation [10,12,36].

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors would like to thank Dr. Carlos Santiviago for the access to *Salmonella* strains and guidance for their usage.

Funding

This work was supported by Fondo Nacional de Desarrollo Científico y Tecnológico [11140222 to A.K.]; Comisión Nacional de Investigación Científica y Tecnológica [79130044 to A.K. and Beca Doctorado Nacional scholarship 21151441 to L.L.]; and the National Institutes of Health [GM65183 to M.I.].

References

1. Ibba M, Becker HD, Stathopoulos C, Tumbula DL, Söll D. The adaptor hypothesis revisited. *Trends Biochem Sci.* 2000; 25:311–316. [PubMed: 10871880]
2. Katz A, Elgamal S, Rajkovic A, Ibba M. Non-canonical roles of tRNAs and tRNA mimics in bacterial cell biology. *Mol Microbiol.* 2016; 101:545–558. [PubMed: 27169680]

3. Rodnina MV. The ribosome in action: Tuning of translational efficiency and protein folding. *Protein Sci Publ Protein Soc.* 2016; 25:1390–1406.
4. Netzer N, Goodenbour JM, David A, Dittmar KA, Jones RB, Schneider JR, Boone D, Eves EM, Rosner MR, Gibbs JS, Embry A, Dolan B, Das S, Hickman HD, Berglund P, Bennink JR, Yewdell JW, Pan T. Innate immune and chemically triggered oxidative stress modifies translational fidelity. *Nature.* 2009; 462:522–526. [PubMed: 19940929]
5. Ling J, Söll D. Severe oxidative stress induces protein mistranslation through impairment of an aminoacyl-tRNA synthetase editing site. *Proc Natl Acad Sci.* 2010; 107:4028–4033. [PubMed: 20160114]
6. Moura GR, Paredes JA, Santos MAS. Development of the genetic code: insights from a fungal codon reassignment. *FEBS Lett.* 2010; 584:334–341. [PubMed: 19941859]
7. Mohler K, Ibba M. Translational fidelity and mistranslation in the cellular response to stress. *Nat Microbiol.* 2017; 2:17117. doi: 10.1038/nmicrobiol.2017.117 [PubMed: 28836574]
8. Komar AA. The Yin and Yang of codon usage. *Hum Mol Genet.* 2016; 25:R77–R85. [PubMed: 27354349]
9. Rocha EPC. Codon usage bias from tRNA's point of view: Redundancy, specialization, and efficient decoding for translation optimization. *Genome Res.* 2004; 14:2279–2286. [PubMed: 15479947]
10. Subramaniam AR, Pan T, Cluzel P. Environmental perturbations lift the degeneracy of the genetic code to regulate protein levels in bacteria. *Proc Natl Acad Sci U S A.* 2013; 110:2419–2424. [PubMed: 23277573]
11. Hu S, Wang M, Cai G, He M. Genetic code-guided protein synthesis and folding in *Escherichia coli*. *J Biol Chem.* 2013; 288:30855–30861. [PubMed: 24003234]
12. Subramaniam AR, Deloughery A, Bradshaw N, Chen Y, O'Shea E, Losick R, Chai Y. A serine sensor for multicellularity in a bacterium. *ELife.* 2013; 2:e01501. [PubMed: 24347549]
13. Brandis G, Hughes D. The Selective Advantage of Synonymous Codon Usage Bias in *Salmonella*. *PLoS Genet.* 2016; 12:e1005926. [PubMed: 26963725]
14. Spencer PS, Siller E, Anderson JF, Barral JM. Silent substitutions predictably alter translation elongation rates and protein folding efficiencies. *J Mol Biol.* 2012; 422:328–335. [PubMed: 22705285]
15. Bustamante P, Covarrubias PC, Levicán G, Katz A, Tapia P, Holmes D, Quatrini R, Orellana O. ICE Afe 1, an actively excising genetic element from the biomining bacterium *Acidithiobacillus ferrooxidans*. *J Mol Microbiol Biotechnol.* 2012; 22:399–407. [PubMed: 23486178]
16. Kane SR, Chakicherla AY, Chain PSG, Schmidt R, Shin MW, Legler TC, Scow KM, Larimer FW, Lucas SM, Richardson PM, Hristova KR. Whole-genome analysis of the methyl tert-butyl ether-degrading beta-proteobacterium *Methylibium petroleiphilum* PM1. *J Bacteriol.* 2007; 189:1931–1945. [PubMed: 17158667]
17. Yoshikawa G, Askora A, Blanc-Mathieu R, Kawasaki T, Li Y, Nakano M, Ogata H, Yamada T. *Xanthomonas citri* jumbo phage XacN1 exhibits a wide host range and high complement of tRNA genes. *Sci Rep.* 2018; 8:4486. doi: 10.1038/s41598-018-22239-3 [PubMed: 29540765]
18. McDonald MJ, Chou C-H, Swamy KB, Huang H-D, Leu J-Y. The evolutionary dynamics of tRNA-gene copy number and codon-use in *E. coli*. *BMC Evol Biol.* 2015; 15doi: 10.1186/s12862-015-0441-y
19. Dong H, Nilsson L, Kurland CG. Co-variation of tRNA abundance and codon usage in *Escherichia coli* at different growth rates. *J Mol Biol.* 1996; 260:649–663. [PubMed: 8709146]
20. Ikemura T. Correlation between the abundance of *Escherichia coli* transfer RNAs and the occurrence of the respective codons in its protein genes: a proposal for a synonymous codon choice that is optimal for the *E. coli* translational system. *J Mol Biol.* 1981; 151:389–409. [PubMed: 6175758]
21. Levicán G, Katz A, Valdés JH, Quatrini R, Holmes DS, Orellana O. A 300 kbp genome segment, including a complete set of tRNA genes, is dispensable for *Acidithiobacillus ferrooxidans*. *Adv Mater Res, Trans Tech Publ.* 2009:187–190.
22. Alamos P, Tello M, Bustamante P, Gutiérrez F, Shmaryahu A, Maldonado J, Levicá G, Orellana O. Functionality of tRNAs encoded in a mobile genetic element from an acidophilic bacterium. *RNA Biol.* 2017; doi: 10.1080/15476286.2017.1349049

23. Gardner SN, Slezak T, Hall BG. kSNP3.0: SNP detection and phylogenetic analysis of genomes without genome alignment or reference genome. *Bioinforma Oxf Engl.* 2015; 31:2877–2878.
24. Lowe TM, Chan PP. tRNAscan-SE On-line: integrating search and context for analysis of transfer RNA genes. *Nucleic Acids Res.* 2016; 44:W54–57. [PubMed: 27174935]
25. Karlin S, Mrázek J, Campbell A, Kaiser D. Characterizations of highly expressed genes of four fast-growing bacteria. *J Bacteriol.* 2001; 183:5025–5040. [PubMed: 11489855]
26. Thomson NR, Clayton DJ, Windhorst D, Vernikos G, Davidson S, Churcher C, Quail MA, Stevens M, Jones MA, Watson M, Barron A, Layton A, Pickard D, Kingsley RA, Bignell A, Clark L, Harris B, Ormond D, Abdellah Z, Brooks K, Cherevach I, Chillingworth T, Woodward J, Norberczak H, Lord A, Arrowsmith C, Jagels K, Moule S, Mungall K, Sanders M, Whitehead S, Chabalgoity JA, Maskell D, Humphrey T, Roberts M, Barrow PA, Dougan G, Parkhill J. Comparative genome analysis of *Salmonella* Enteritidis PT4 and *Salmonella* Gallinarum 287/91 provides insights into evolutionary and host adaptation pathways. *Genome Res.* 2008; 18:1624–1637. [PubMed: 18583645]
27. Bucarey SA, Villagra NA, Martinic MP, Trombert AN, Santiviago CA, Maulén NP, Youderian P, Mora GC. The *Salmonella enterica* serovar Typhi *tsx* gene, encoding a nucleoside-specific porin, is essential for prototrophic growth in the absence of nucleosides. *Infect Immun.* 2005; 73:6210–6219. [PubMed: 16177292]
28. Satapathy SS, Dutta M, Buragohain AK, Ray SK. Transfer RNA gene numbers may not be completely responsible for the codon usage bias in asparagine, isoleucine, phenylalanine, and tyrosine in the high expression genes in bacteria. *J Mol Evol.* 2012; 75:34–42. [PubMed: 23053196]
29. Zhong J, Xiao C, Gu W, Du G, Sun X, He QY, Zhang G. Transfer RNAs Mediate the Rapid Adaptation of *Escherichia coli* to Oxidative Stress. *PLOS Genet.* 2015; 11:e1005302. [PubMed: 26090660]
30. Svenningsen SL, Kongstad M, Stenum TS, Muñoz-Gómez AJ, Sørensen MA. Transfer RNA is highly unstable during early amino acid starvation in *Escherichia coli*. *Nucleic Acids Res.* 2017; 45:793–804. [PubMed: 27903898]
31. Higgs PG, Ran W. Coevolution of codon usage and tRNA genes leads to alternative stable states of biased codon usage. *Mol Biol Evol.* 2008; 25:2279–2291. [PubMed: 18687657]
32. Sharp PM, Li WH. The codon Adaptation Index--a measure of directional synonymous codon usage bias, and its potential applications. *Nucleic Acids Res.* 1987; 15:1281–1295. [PubMed: 3547335]
33. dos Reis M, Wernisch L, Savva R. Unexpected correlations between gene expression and codon usage bias from microarray data for the whole *Escherichia coli* K-12 genome. *Nucleic Acids Res.* 2003; 31:6976–6985. [PubMed: 14627830]
34. Avcilar-Kucukgoze I, Bartholomäus A, Cordero Varela JA, Kaml RF-X, Neubauer P, Budisa N, Ignatova Z. Discharging tRNAs: a tug of war between translation and detoxification in *Escherichia coli*. *Nucleic Acids Res.* 2016; 44:8324–8334. [PubMed: 27507888]
35. Starzyk R. tRNA base modifications and gene regulation. *Trends Biochem Sci.* 1984; 9:333–334.
36. Subramaniam AR, Zid BM, O’Shea EK. An integrated approach reveals regulatory controls on bacterial translation elongation. *Cell.* 2014; 159:1200–1211. [PubMed: 25416955]

Highlights

- Average number of tRNA genes correlate with codon usage in highly expressed genes
- Changes in the number of tRNA genes do not correlate to codon usage
- Altered tRNA gene copy numbers do not correlate to changes of codon translation
- Our data indicates that tRNA gene copies adapt to codon usage, but not the contrary




Figure 1. tRNA gene copy numbers and frequency of codon usage in *Salmonella enterica*
 The box plots show A) the average of tRNA gene copy numbers for each possible anticodon and B) the average of frequency of codon usage in all or C) a subset of highly expressed genes in 139 *Salmonella enterica* strains. In all graphs an “X” symbol indicates the most extreme values, a circle indicates the mean value and the horizontal lines of the box indicate limits where 5, 50 and 95 % of the data is contained. Whiskers indicate standard deviations.




Figure 2. Relations between frequency of codon usage in highly expressed genes and number of tRNA genes in *Salmonella enterica*

Comparison of the number of genes for A) tRNA^{Ala}_{UGC}, B) tRNA^{Ala}_{GGC} and C) tRNA^{Glu}_{UUC} with the frequency of usage of all codons decoding the corresponding amino acid in highly expressed genes of each studied *S. enterica* genome. Cognate codons are symbolized by a “+” symbol while non-cognate codons by an “X” symbol. Genes for tRNAs with other anticodons for these amino acids were not found in *S. enterica* genomes. Thus, corresponding graphs are shown in Figure S2.




Figure 3. Efficiency of codon translation is similar between *Salmonella enterica* strains that have different numbers of tRNA gene copies

Four contiguous identical codons were cloned in an arabinose inducible plasmid as a translational fusion to *gfp* and in transcriptional fusion to *mCherry*. The graphs show the fluorescence of GFP normalized by that of mCherry for each plasmid transformed in three different *S. enterica* serovars. A) GFP/mCherry fluorescence ratios for Glu and Ala codons. Values above the bars indicate the frequency of codon usage (in percent) in highly expressed genes of the corresponding strains. B) and C) show the same data normalized by the average GFP/mCherry fluorescence ratio of B) GAA Glu codon and C) GCT Ala codon. *S. Typhimurium* data is shown in white bars, *S. Enteritidis* in light gray bars and *S. Typhi* in dark gray bars.

Number of tRNA genes for Glu and Ala and the corresponding translated codons in *Salmonella enterica* strains used in translation efficiency experiments

Table 1

	tRNA ^{Glu} _{UUC}	tRNA ^{Glu} _{CUC}	tRNA ^{Ala} _{AGC}	tRNA ^{Ala} _{GCC}	tRNA ^{Ala} _{UGC}	tRNA ^{Ala} _{GCC}
tRNA genes in:						
<i>S. Typhimurium</i> 14028S	4	0	0	2	3	0
<i>S. Typhi</i> STH2370	1	0	0	2	1	0
<i>S. Enteritidis</i> PT4 NCTC13349	4	0	0	2	3	0
Codons translated ^a	GAA, GAG	NA	NA	GCC, GCU	GCA, GCG, GCU	NA

^aNA indicates not applicable, tRNA gene not present in the corresponding genome