

Pharmacy Faculty Articles and Research

School of Pharmacy

2016

Capillary Endothelia from Two ADPKD Patients are Polyploidy

Sarmed H. Kathem *Chapman University*

Wissam A. AbouAlaiwi University of Toledo

Xiaolin Zi University of California

Surya M. Nauli Chapman University, nauli@chapman.edu

Follow this and additional works at: https://digitalcommons.chapman.edu/pharmacy_articles

Part of the Cardiovascular System Commons, Endocrine System Commons, Medical Genetics Commons, Other Pharmacy and Pharmaceutical Sciences Commons, and the Pharmaceutics and Drug Design Commons

Recommended Citation

Kathem SH, AbouAlaiwi WA, Zi X, Nauli SM. Capillary endothelia from two ADPKD patients are polyploidy. *Ann Clin Cytol Pathol.* 2016; 2(2): 1022.

This Article is brought to you for free and open access by the School of Pharmacy at Chapman University Digital Commons. It has been accepted for inclusion in Pharmacy Faculty Articles and Research by an authorized administrator of Chapman University Digital Commons. For more information, please contact laughtin@chapman.edu.

Capillary Endothelia from Two ADPKD Patients are Polyploidy

Comments

This article was originally published in *Annals of Clinical Cytology and Pathology*, volume 2, issue 2, in 2016. DOI:

Creative Commons License



This work is licensed under a Creative Commons Attribution 4.0 License.

Copyright The authors

⊘SciMedCentral

Case Report

Capillary Endothelia from Two ADPKD Patients are Polyploidy

Sarmed H. Kathem^{1,2}, Wissam A. AbouAlaiwi³, Xiaolin Zi² and Surva M. Nauli^{1,2*}

¹Department of Biomedical & Pharmaceutical Sciences, Chapman University, USA ²Department of Urology, University of California, USA

³Department of Pharmacology and Experimental Therapeutics, University of Toledo, USA

Abstract

Bilateral renal cyst formation is the main feature of autosomal dominant polycystic kidney disease (ADPKD). We and other laboratories have previously shown that cystlining epithelia of kidneys from ADPKD patients are characterized by polyploidy. In this report, we show that endothelia from the renal capillary beds of two ADPKD patients are also polyploidy. Spectral karyotyping study further confirms our flow cytometry analyses. We suggest that polyploidy may be used as a potential cellular marker in ADPKD.

INTRODUCTION

Autosomal dominant polycystic kidney disease (ADPKD) is the most common life-threatening hereditary genetic disease. Although kidneys are the major sites of clinical disease, the prevalence of extra renal manifestations in ADPKD is very high. These extra renal manifestations include cyst formation in other ductal organs and various cardiovascular abnormalities [1-3]. Polycystic kidney has also been associated with polyploidy [4-7]. At the molecular level, vascular endothelia from ADPKD have been associated with dysfunctional primary cilia [8] Aberrant cell proliferation has been reported in cells with defective cilia [9,10]. We recently showed that cystic renal epithelia from patients are polyploidy, probably through abnormal regulation of chromosomal passenger complex. Abnormality of this protein complex has also been associated with aneurysm formation in vascular endothelia of mouse polycystic models [11,12]. However, the endothelia of renal capillary beds from ADPKD patients have never been previously studied.

CASE REPORT

Human kidney samples from three subjects were taken; two were from ADPKD patients (54-years-old female and 49-yearsold male), and the other was from a non-ADPKD subject. The nephrectomy was done in both ADPKD patients, because these kidneys had reached the end-stage renal failure and were further complicated by infections. In general, ADPKD kidney has a grossly distorted architecture characterized by fluid-filled cysts (Figure 1). We isolated endothelia from the renal capillary beds. First branches of capillaries from interlobar arteries were pooled. Due to the nature of primary cells from the capillary beds, we were not able to purify these isolated endothelial cells. Successful cell

Annals of Clinical Cytology and Pathology

*Corresponding author

Surya M. Nauli, Chapman University, Harry and Diane Rinker Health Science Campus, 9401 Jeronimo Road, Irvine, CA 92618-1908, USA, Tel: 714-516-5480; Fax: 714-516-5481; Email: nauli@chapman.edu, snauli@uci.edu

Submitted: 10 March 2016 Accepted: 31 April 2016

Published: 25 April 2016

Copyright

© 2016 Nauli et al.

OPEN ACCESS

isolation typically provides homogenous cell morphology based on the side scatter from the flow cytometry. Capillaries were
on the side scatter from the now cytometry, capitalies were
then briefly rinsed with phosphate buffered saline (pH 7.4). To
dissociate the endothelial cells, capillaries were subsequently
incubated with trypsin for 20-30 minutes at 37 $^{\mathrm{o}}\mathrm{C}.$ Because
capillaries contain mainly endothelia, the isolation typically
provides relatively pure endothelial cells. Endothelial cells from
the samples were then processed for flow cytometry and spectral
karyotyping analysis to investigate cellular polyploidy. Propidium
iodide (PI) and 5-bromodeoxyuridine (BrdU) staining were
used to analyze cellular polyploidy from the samples. Generally,
PI was used for DNA content quantification, while BrdU was
used as a marker for cell division. Our flow cytometry analyses
showed abnormal polyploidy peaks in both ADPKD samples but
not in non-ADPKD sample (Figure 2). Cells with >2N represent



Figure 1 Autosomal dominant polycystic kidney a: Polycystic kidney is characterized by the grossly distorted and enlarged kidney. b: A polycystic kidney was cut in half, showing the presence of numerous fluid-filled cysts. Fluid accumulation in the cysts contributes to the heavy weight and large size of an ADPKD kidney.

Cite this article: Kathem SH, AbouAlaiwi WA, Zi X, Nauli SM (2016) Capillary Endothelia from Two ADPKD Patients are Polyploidy. Ann Clin Cytol Pathol 2(2): 1022.

⊘SciMedCentral

polyploidy and have higher DNA content. In non-ADPKD, cells with >2N represent about 0.8%, while in ADPKD patients# 1 and 2 the polyploidy cells constitute of 16.5% and 8.2%, respectively. Interestingly, further analysis of PI and BrdU showed that the abnormal polyploidy cells retain their ability to undergo cell division. This result indicates that endothelia from ADPKD patients are associated with polyploidy and that those polyploidy cells could undergo cell division. Spectral karyotyping (SKY) was also used to confirm our flow cytometry results. Chromosomes from a single endothelium were differentially labeled with probes containing a mixture of fluorescent dyes (Rhodamine, Texas Red, Cy5, FITC and Cy5.5), as describe previously [13]. Unfortunately, we were not successful to karyotype endothelial sample from the second ADPKD patient. From the first ADPKD patient, a single ADPKD endothelium was analyzed, and it showed cellular tetraploidy (Figure 3).

DISCUSSION

We here report that endothelia from renal vascular of ADPKD patients are polyploidy. Those cells could potentially have abnormal cell division that may lead to vascular dysfunction. We previously reported that vascular endothelia from ADPKD patients are less sensitive to fluid-shear stress [8]. This report may thus associate the mechano-dysfunction and polyploidy in ADPKD endothelia. Independent studies have shown centrosomal over-amplification in polycystic kidneys from mice and human [4-7]. The abnormal centrosomal amplification was suggested to occur early in cystic kidney disease. Consequently, abnormal cell division and chromosomal segregation resulted from centrosomal over-duplication lead to polyploidy and genomic instability. Polyploidy has also been observed in vascular endothelial of polycystic kidney mouse models [11,12]. At least in the mice, it



Figure 2 Flow cytometry analysis: Vascular endothelia were isolated from the renal capillary beds of non-APKD (a) and two independent ADPKD (b, c) kidneys. Cells were then labeled with propidium iodide (PI) and 5-bromodeoxyuridine (BrdU). PI and BrdU staining were analyzed individually and interdependently. Brackets indicate the presence of polyploidy cells.



Figure 3 Spectral Karyotype analysis Chromosomes from an endothelium were isolated and processed for labeling with fluorescence probes specific for each individual chromosome. A tetraploidy with a total of 92 chromosomes was observed in the sample obtained from a female ADPKD patient.

⊘SciMedCentral

was postulated that polyploidy in endothelial cells could result in aneurysm formation, probably through abnormal chromosomal passenger complex. Our current study reinforces the previous findings that vascular endothelia and renal epithelia from polycystic mouse models and ADPKD patients are polyploidy.

CONCLUSION

Flow cytometry analyses using PI and BrdU indicate that ADPKD endothelia have higher DNA content and polyploidy. Analysis with spectral karyotyping further confirmed this result showing abnormal chromosomal number and tetraploidy. Our study is preliminary in nature, and a greater number of patients are needed for a future study. If confirmed in a larger scale study, this result might have an impact on the pathogenic event leading to the development of vascular abnormalities in ADPKD. We propose that polyploidy can be a broad cellular feature not only in the renal epithelia but also in vascular endothelia and that polyploidy may be used as a cellular marker to understand disease progression in ADPKD.

REFERENCES

- 1. Ecder T. Cardiovascular complications in autosomal dominant polycystic kidney disease. Curr Hypertens Rev. 2013; 9: 2-11.
- Perrone RD, Malek AM, Watnick T. Vascular complications in autosomal dominant polycystic kidney disease. Nat Rev Nephrol. 2015; 11: 589-598.
- 3. Torres VE, Harris PC. Autosomal dominant polycystic kidney disease: the last 3 years. Kidney Int. 2009; 76: 149-168.
- 4. Jin X, Muntean BS, Aal-Aaboda MS, Duan Q, Zhou J, Nauli SM. L-type calcium channel modulates cystic kidney phenotype. Biochim Biophys Acta. 2014; 1842: 1518-1526.

- Zhou X, Fan LX, Li K, Ramchandran R, Calvet JP, Li X. SIRT2 regulates ciliogenesis and contributes to abnormal centrosome amplification caused by loss of polycystin-1. Hum Mol Genet. 2014; 23: 1644-1655.
- 6. Burtey S, Riera M, Ribe E, Pennenkamp P, Rance R, Luciani J, et al. Centrosome overduplication and mitotic instability in PKD2 transgenic lines. Cell Biol Int. 2008; 32: 1193-1198.
- Battini L, Macip S, Fedorova E, Dikman S, Somlo S, Montagna C, et al. Loss of polycystin-1 causes centrosome amplification and genomic instability. Hum Mol Genet. 2008; 17: 2819-2833.
- 8. AbouAlaiwi WA, Takahashi M, Mell BR, Jones TJ, Ratnam S, Kolb RJ, et al. Ciliary polycystin-2 is a mechanosensitive calcium channel involved in nitric oxide signaling cascades. Circ Res. 2009; 104: 860-869.
- 9. Jonassen JA, San Agustin J, Follit JA, Pazour GJ. Deletion of IFT20 in the mouse kidney causes misorientation of the mitotic spindle and cystic kidney disease. J Cell Biol. 2008; 183: 377-384.
- 10. Qin H, Wang Z, Diener D, Rosenbaum J. Intraflagellar transport protein 27 is a small G protein involved in cell-cycle control. Curr Biol. 2007; 17: 193-202.
- 11. Aboualaiwi WA, Muntean BS, Ratnam S, Joe B, Liu L, Booth RL, et al. Survivin-induced abnormal ploidy contributes to cystic kidney and aneurysm formation. Circulation. 2014; 129: 660-672.
- 12. AbouAlaiwi WA, Ratnam S, Booth RL, Shah JV, Nauli SM. Endothelial cells from humans and mice with polycystic kidney disease are characterized by polyploidy and chromosome segregation defects through survivin down-regulation. Hum Mol Genet. 2011; 20: 354-367.
- 13. AbouAlaiwi WA, Rodriguez I, Nauli SM. Spectral karyotyping to study chromosome abnormalities in humans and mice with polycystic kidney disease. J Vis Exp. 2012; 60.

Cite this article

Kathem SH, AbouAlaiwi WA, Zi X, Nauli SM (2016) Capillary Endothelia from Two ADPKD Patients are Polyploidy. Ann Clin Cytol Pathol 2(2): 1022.