
Spring 5-2021

Sex as a Moderator in the Relationship between Post-operative Pain and Analgesics Administered in Children Undergoing Elective Surgery

Amber M. Osorno
Chapman University, osorno@chapman.edu

Michelle A. Fortier
University of California, Irvine, mfortier@hs.uci.edu

Zeev N. Kain
University of California, Irvine, zkain@hs.usi.edu

Brooke N. Jenkins
Chapman University, bjenkins@chapman.edu

Follow this and additional works at: https://digitalcommons.chapman.edu/cusrd_abstracts



Part of the [Other Psychiatry and Psychology Commons](#), [Pain Management Commons](#), [Pediatrics Commons](#), and the [Surgery Commons](#)

Recommended Citation

Osorno, Amber M.; Fortier, Michelle A.; Kain, Zeev N.; and Jenkins, Brooke N., "Sex as a Moderator in the Relationship between Post-operative Pain and Analgesics Administered in Children Undergoing Elective Surgery" (2021). *Student Scholar Symposium Abstracts and Posters*. 434.
https://digitalcommons.chapman.edu/cusrd_abstracts/434

This Poster is brought to you for free and open access by the Center for Undergraduate Excellence at Chapman University Digital Commons. It has been accepted for inclusion in Student Scholar Symposium Abstracts and Posters by an authorized administrator of Chapman University Digital Commons. For more information, please contact laughtin@chapman.edu.




Sex as a moderator in the relationship between post-operative pain and analgesics administered in children undergoing elective surgery

Amber M. Osorno, Michelle A. Fortier, Zeev N. Kain, & Brooke N. Jenkins
Chapman University
Center on Stress and Health
University of California, Irvine



INTRODUCTION

- Post-operative pain is experienced by 85% of children who undergo surgery. A large percentage of these children do not receive proper pain management after discharge from the post-operative care unit (Fortier, Chou, Maurer, & Kain 2011).
- Among adults, women report more sensations of pain in comparison to men, but women have a higher pain tolerance (Fillingim et al. 2009). The sensation and perceptions of pain may be attributed to physiological characteristics such as hormones or BMI, but the social process of how pain is communicated between the two genders cannot be denied (Hussain et al. 2013).
- Experimental research suggests that adults have biases when it comes to pediatric post-operative pain (Cohen et al. 2014). Results showed that adults rate girls as more sensitive to pain and that they display a greater amount of pain.
- Previous research has revealed that when children are sick, parents sympathize more with their daughters than with their sons (Becker & Hu, 2008).
- A child's post operative pain could be directly related to the amount of analgesics given by the parent. However, looking at previous research, a child's gender could also influence this relationship because of societal biases and sociocultural factors (Cohen et al. 2014).
- The purpose of the current study is to understand the relationship between post-operative pain among children and analgesics given to the child, and how gender moderates this association.



METHOD

Participants: Procedures

112 patients ages 2-13 (Mean = 5.80) at the Children's Hospital of Orange County (CHOC Children's) who underwent elective surgery. 59% male and 41% female. 47% Hispanic, 25% White, and 28% other.

Parents and children were recruited prior to surgery to complete online surveys in a hospital setting where baseline information was gathered. After surgery parents were asked to report pain levels and analgesics given one, three, and seven days post-operatively through an online survey.

MEASURES

- **PPPM - Parent Postoperative Pain Measure** evaluates the child's pain, on days 1, 3, and 7 after surgery (Chambers et al. 1996).
 - Parents answered simply "yes" (coded 1) or "no" (coded 0) to 15 questions such as, "Since discharge from the hospital does your child, hold the sore part or his/her body?" or "Since discharge from the hospital does your child, cry more easily than usual?"
 - PPPM items are summed together. A score of 6 or higher is indicative of "clinically significant pain".
- An **Analgesic Consumption, Follow Up** survey was answered days 1, 3, and 7 post-operatively by parents to gather information on what medication(s) was given, the unit of measurement, and the amount administered. The reported analgesic was then converted into milligrams by the amount of active ingredient, using ratios to convert from tablespoons, ccs, droppers, teaspoons etc. The analgesics were then separated by types, specifically acetaminophen and ibuprofen. To account for weight differences, the amount of acetaminophen or ibuprofen was divided by the weight of the child in kilograms.

RESULTS

Analgesic	Post-Operative Pain and Analgesics	Coefficient of Sex as a Moderator
Post-Operative Day 1		
Ibuprofen	$b = 0.24^*$	$b = 0.27$
Acetaminophen	$b = 0.22^*$	$b = 0.08$
Post-Operative Day 3		
Ibuprofen	$b = 0.72^*$	$b = 0.52$
Acetaminophen	$b = 0.34^*$	$b = 0.39$
Post-Operative Day 7		
Ibuprofen	$b = 0.22$	$b = 0.91$
Acetaminophen	$b = 0.11$	---

Note: * $p < 0.05$

REFERENCES

- Becker, J. B., & Hu, M. (2008). Sex differences in drug abuse. *Frontiers in neuroendocrinology*, 29(1), 36–47.
- Chambers, C.T., Reid, G.J., McGrath, P.J., & Finley, G.A. (1996). Development and preliminary validation of a postoperative pain measure for parents. *Pain*, 68, 307–313.
- Cohen, L. L., Cobb, J., & Martin, S. R. (2014). Gender Biases in Adult Ratings of Pediatric Pain. *Children's Health Care*, 43(2), 87–95. <https://doi-org.libproxy.chapman.edu/10.1080/02739615.2014.849918>
- Fortier, M. A., Chou, J., Maurer, E. L., & Kain, Z. N. (2011). Acute to chronic postoperative pain in children: preliminary findings. *Journal of Pediatric Surgery*, 46(9), 1700–1705.
- Fillingim, R. B., King, C. D., Ribeiro-Dasilva, M. C., Rahim-Williams, B., & Riley, J. L., 3rd (2009). Sex, gender, and pain: a review of recent clinical and experimental findings. *The journal of pain : official journal of the American Pain Society*, 10(5), 447–485.
- Hussain, A. M., Khan, F. A., Ahmed, A., Chawla, T., & Azam, S. I. (2013). Effect of gender on pain perception and analgesic consumption in laparoscopic cholecystectomy: An observational study. *Journal of anaesthesiology, clinical pharmacology*, 29(3), 337–341.

RESULTS

Postoperative Day 1:

- For every one unit increase in pain, there was a 0.24 mg/kg increase in **ibuprofen** administered to the child, $b = 0.24$, $t = 3.55$, $p < 0.05$.
 - Sex did not significantly moderate the relationship between post-operative pain and ibuprofen administered to the child, $b = 0.27$, $t = 1.57$, $p > 0.05$
- For every one unit increase in pain, there was a 0.22mg increase in **acetaminophen** administered to the child, $b = 0.22$, $t = 5.38$, $p < 0.05$.
 - Sex did not significantly moderate the relationship between post-operative pain and acetaminophen administered to the child, $b = 0.08$, $t = 0.89$, $p > 0.05$.

Postoperative Day 3

- For every one unit increase in pain, there was a 0.72 mg/kg increase in **ibuprofen** administered to the child, $b = 0.72$, $t = 4.81$, $p < 0.05$.
 - Sex did not significantly moderate the relationship between post-operative pain and ibuprofen administered to the child, $b = 0.52$, $t = 1.72$, $p > 0.05$
- For every one unit increase in pain, there was a 0.34 mg increase in **acetaminophen** administered to the child, $b = 0.34$, $t = 4.44$, $p < 0.05$.
 - Sex did not significantly moderate the relationship between post-operative pain and acetaminophen administered to the child, $b = 0.39$, $t = 1.81$, $p > 0.05$

Postoperative Day 7

- There was no significant relationship between reported pain post-operatively day seven and given **ibuprofen**, $b = 0.22$, $t = 1.47$, $p > 0.05$
 - Sex did not significantly moderate the relationship between post-operative pain and ibuprofen administered to the child, $b = 0.91$, $t = 1.89$, $p > 0.05$
- There was no significant relationship between reported pain post-operatively day seven and given **acetaminophen**, $b = 0.11$, $t = 1.94$, $p > 0.05$

CONCLUSION

- Gender is not a moderator in the relationship between post-operative pain and analgesics administered to the child postoperatively all 3 days of data collection (days 1,3, and 7 post-operatively). Although it was non-significant, it is important to note the significant relationship between pain levels reported by parents and analgesic consumption. Parents are giving their child dosages of analgesics that are directly correlated to the reported pain values.
- Although gender was not a significant moderator, further research could be done analyzing socio-cultural influences between this association like number of siblings, gender of siblings, and temperament of the child.
- Some limitations of the study include the type of surgery that the participants were undergoing. Although the participants surgeries were elective, differences could lie in the experience of pain. If a participant had back pain due to surgery this could be more impactful on pain scores where other pain could be less.
- Perhaps with current gender equality movements and current education on gender equality makes this relationship not significant. Gender equality has perhaps had a positive influence on parents-perception of their child's pain where both genders are seen equal.
- The data presented could support the idea that both boys and girls are equal in a post-operative care environment, and it is not indicative that gender inequality is initiated at such a young age.
- Future analysis should analyze factors that interact with this association like the gender of the parent that is administering analgesics and the parent's education of analgesic usage.