
11-17-2022

Technical Note: On Uncertainties in Plant Water Isotopic Composition Following Extraction by Cryogenic Vacuum Distillation

Haoyu Diao

Swiss Federal Institute for Forest, Snow and Landscape Research WSL

Philipp Schuler

Swiss Federal Institute for Forest, Snow and Landscape Research WSL

Gregory R. Goldsmith

Chapman University, goldsmit@chapman.edu


Rolf T. W. Siegwolf

Swiss Federal Institute for Forest, Snow and Landscape Research WSL

Matthias Saurer

Swiss Federal Institute for Forest, Snow and Landscape Research WSL

See next page for additional authors
Follow this and additional works at: https://digitalcommons.chapman.edu/sees_articles

 Part of the [Botany Commons](#), [Environmental Chemistry Commons](#), [Hydrology Commons](#), [Organic Chemistry Commons](#), and the [Other Plant Sciences Commons](#)

Recommended Citation

Diao, H., Schuler, P., Goldsmith, G. R., Siegwolf, R. T. W., Saurer, M., and Lehmann, M. M.: Technical note: On uncertainties in plant water isotopic composition following extraction by cryogenic vacuum distillation, *Hydrol. Earth Syst. Sci.*, 26, 5835–5847, <https://doi.org/10.5194/hess-26-5835-2022>, 2022.

This Article is brought to you for free and open access by the Science and Technology Faculty Articles and Research at Chapman University Digital Commons. It has been accepted for inclusion in Biology, Chemistry, and Environmental Sciences Faculty Articles and Research by an authorized administrator of Chapman University Digital Commons. For more information, please contact laughtin@chapman.edu.

Technical Note: On Uncertainties in Plant Water Isotopic Composition Following Extraction by Cryogenic Vacuum Distillation

Comments

This article was originally published in *Hydrology and Earth System Sciences*, volume 26, in 2022.
<https://doi.org/10.5194/hess-26-5835-2022>

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

Copyright

The authors

Authors

Haoyu Diao, Philipp Schuler, Gregory R. Goldsmith, Rolf T. W. Siegwolf, Matthias Saurer, and Marco M. Lehmann



Technical note: On uncertainties in plant water isotopic composition following extraction by cryogenic vacuum distillation

Haoyu Diao^{1,2,3}, Philipp Schuler¹, Gregory R. Goldsmith⁴, Rolf T. W. Siegwolf¹, Matthias Saurer¹, and Marco M. Lehmann¹

¹Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf 8903, Switzerland

²CAS Key Laboratory of Forest Ecology and Management, Institute of Applied Ecology, Chinese Academy of Sciences, Shenyang 110016, China

³University of Chinese Academy of Sciences, Beijing 100049, China

⁴Schmid College of Science and Technology, Chapman University, Orange, CA 92866 USA

Correspondence: Marco M. Lehmann (marco.lehmann@wsl.ch)

Received: 3 May 2022 – Discussion started: 9 May 2022

Revised: 24 October 2022 – Accepted: 28 October 2022 – Published: 17 November 2022

Abstract. Recent studies have challenged the interpretation of plant water isotopes obtained through cryogenic vacuum distillation (CVD) based on observations of a large ²H fractionation. These studies have hypothesized the existence of an H-atom exchange between water and organic tissue during CVD extraction with the magnitude of H exchange related to relative water content of the sample; however, clear evidence is lacking. Here, we systematically tested the uncertainties in the isotopic composition of CVD-extracted water by conducting a series of incubation and rehydration experiments using isotopically depleted water, water at natural isotope abundance, woody materials with exchangeable H, and organic materials without exchangeable H (cellulose triacetate and caffeine). We show that the offsets between hydrogen and oxygen isotope ratios and expected reference values ($\Delta^2\text{H}$ and $\Delta^{18}\text{O}$) have inversely proportional relationships with the absolute amount of water being extracted, i.e. the lower the water amount, the higher the $\Delta^2\text{H}$ and $\Delta^{18}\text{O}$. However, neither $\Delta^2\text{H}$ nor $\Delta^{18}\text{O}$ values, were related to sample relative water content. The $\Delta^2\text{H}$ pattern was more pronounced for materials with exchangeable H atoms than with non-exchangeable H atoms. This is caused by the combined effect of H exchange during the incubation of materials in water and isotopic enrichments during evaporation and sublimation that depend on absolute water amount. The H exchange during CVD extraction itself was negligible. Despite these technical issues, we observed that the water amount-dependent patterns were much less pronounced for samples

at natural isotope abundance and particularly low when sufficiently high amounts of water were extracted ($> 600 \mu\text{L}$). Our study provides new insights into the mechanisms causing isotope fractionation during CVD extraction of water. The methodological uncertainties can be controlled if large samples of natural isotope abundance are used in ecohydrological studies.

1 Introduction

Hydrogen and oxygen isotope ratios ($\delta^2\text{H}$ and $\delta^{18}\text{O}$) in plant water are powerful tools for tracing the water movement of ecosystems and providing information on the source water used by plants (Goldsmith et al., 2017; Flanagan and Ehleringer, 1991; Brinkmann et al., 2019; Nehemy et al., 2019). The most widely used approach for determining the $\delta^2\text{H}$ and $\delta^{18}\text{O}$ of plant water is by first extracting water from plants using cryogenic vacuum distillation (CVD) method (Ehleringer et al., 2000; West et al., 2006; Ingraham and Shadel, 1992; Orłowski et al., 2013). Since no isotopic fractionation generally occurs during plant water uptake (White et al., 1985; Zimmermann et al., 1967; Poca et al., 2019), it is generally believed that the isotope ratios in stem water (root crown water for grasses) derived from the CVD extraction (δ_{CVD}) are equal to that in the plant source water (δ_{source}). However, an increasing number of studies have found that δ_{CVD} , especially $\delta^2\text{H}_{\text{CVD}}$, does not actually re-

flect plant source water, as shown by a strong negative $\delta^2\text{H}$ offset ($\Delta^2\text{H} = \delta^2\text{H}_{\text{CVD}} - \delta^2\text{H}_{\text{source}}$) between CVD-extracted water and source water (Lin and Da S. L. Sternberg, 1993; Zhao et al., 2016; Barbeta et al., 2020; Allen and Kirchner, 2022; Newberry et al., 2017). This issue is important because it may invalidate conclusions on plant water uptake patterns and the partitioning of different pools of water in an ecosystem (Allen et al., 2019; Gessler et al., 2022; Goldsmith et al., 2012; Evaristo and McDonnell, 2017).

A critical hypothesis for the observed $\Delta^2\text{H}$ is the spatial heterogeneity of hydrogen isotope composition of water in stems (Zhao et al., 2016; Barbeta et al., 2020), i.e. isotopic differences between stem conductive and nonconductive tissues, and isotopic differences from inside to outside along the stem radial direction. This hypothesis was rejected by Chen et al. (2020). The authors rehydrated dry stem samples of saplings of nine plant species in excess of a reference water with a known isotopic composition for 24 h and then CVD-extracted the water that had soaked into the material. They concluded that the observed negative $\Delta^2\text{H}$ was not caused by within-stem isotope heterogeneity, but was more likely related to a hydrogen atom (H) exchange between stem water and the exchangeable H of stem organic tissue that occurs during the CVD extraction. In woody tissue, oxygen-bound H atoms can exchange with those in surrounding liquid water and water vapour. Theoretically, about 30 % to 50 % of the H atoms of cellulose and its precursors are exchangeable (Schuler et al., 2022; Filot et al., 2006). If the CVD extraction disturbs the previously established isotopic equilibration between organic tissue and tissue water because of the gradually isotopically enriched tissue water during extraction, the potential H exchange would affect ecohydrological interpretations of $\Delta^2\text{H}$ (Barbeta et al., 2020; De La Casa et al., 2022). However, Chen et al. (2020) were unable to isolate the H exchange that occurred during the extraction. This is because their experimental design included two possible H exchange steps: one that occurs during rehydration (hereinafter referred to as “H exchange during rehydration”) and a second one that occurs during the extraction itself (hereinafter referred to as “H exchange during extraction”). Only the latter is of interest, because it is the H exchange process that theoretically affects the isotopic composition of CVD-extracted water from actual plant samples. However, empirical evidence for the occurrence of H exchange and its effects on $\delta^2\text{H}_{\text{CVD}}$ during rehydration and/or extraction is still lacking.

Moreover, the study of Chen et al. (2020) also found a significant positive correlation between $\Delta^2\text{H}$ and sample relative water content (RWC). They posited that the H exchange effect was probably dampened at high RWC. This correlation has also been found for both $\Delta^2\text{H}$ and $\Delta^{18}\text{O}$ (i.e. $\delta^{18}\text{O}_{\text{CVD}} - \delta^{18}\text{O}_{\text{source}}$) in soil samples (Wen et al., 2021). However, using cellulose in wood as an example, much of the O-bound H is engaged in H bonding that links cellulose fibrils together, i.e. the non-freely exchangeable “bridging hy-

drogen” (Sepall and Mason, 1961; Meier-Augenstein et al., 2014), which requires high temperatures ($> 100^\circ\text{C}$) for accessing (Schuler et al., 2022). As such, the freely exchangeable H in cellulose is theoretically only about 5 % (Sepall and Mason, 1961; Meier-Augenstein et al., 2014). If the magnitude of H exchange effect during extraction is therefore moderate, the observed $\Delta^2\text{H}$ would be more readily attributed to isotopic fractionation and mixing that occur after the water is extracted from the sample. If this were to be true, what really matters would not be the RWC, but rather the absolute amount of water being extracted (hereinafter referred to “absolute water amount (AWA)”). The effects of AWA on extraction uncertainties could influence both $\Delta^2\text{H}$ and $\Delta^{18}\text{O}$ through fractionation during CVD extraction. However, no studies have tested the effect of sample water amount on the CVD extraction biases yet, and the CVD extraction uncertainties during liquid–vapour phase changes are not well known.

During the CVD extraction, successive isotope fractionation processes occur at each phase change of the water (liquid–vapour–solid–liquid). When liquid sample water is distilled to water vapour, isotope fractionation is expected if incomplete extraction occurs (Orlowski et al., 2013; West et al., 2006). When the water vapour is trapped by liquid nitrogen and turns into solid ice, a sublimation isotope fractionation is also expected. Studies testing the sublimation of pure water ice at low temperatures into a vacuum have shown that the sublimated percentage of the bulk water aliquot was $< 10\%$ within 1 h at -100°C , and the sublimation rate decreased drastically with the decrease of temperature in the cold trap (Mortimer et al., 2018; Lécuyer et al., 2017). Given the fact that the extracted water is frozen at such a low temperature (-196°C) during CVD extraction, negligible amounts of water should be released into the vapour phase by sublimation during the extraction. Nevertheless, an observable enrichment of the residual ice is still possible as the water lost to the vapour phase is isotopically depleted (Lécuyer et al., 2017; Mortimer et al., 2018). In addition, isotope fractionation related to evaporation and mixing with water vapour in the laboratory probably occur during the end of the CVD extraction (i.e. mixing of the extracted water with air humidity of the lab air when the water traps are removed from the extraction system). Although this can be partly avoided by flushing the extraction line with N_2 gas, isotope fractionation can occur when the extracted and frozen water thaws in the water collection tube and when it is transferred to storage vials. A better understanding of these individual isotope fractionation processes during CVD extraction would not only aid in identifying the potential sources of methodological biases (Schoppach and Klaus, 2019) and improving the instrumentation, but also contribute to better estimates of plant water sources by mitigating their effects.

To investigate these open questions, we systematically tested for biases induced by CVD extraction of plant water. A series of experiments that involved extracting water from dif-

ferent kinds of materials (with and without exchangeable H) that were incubated or rehydrated using isotopically depleted water and from plant samples at natural isotope abundance were conducted. Specifically, we hypothesized that the $\Delta^2\text{H}$ is influenced by (i) an H-exchange effect, whereas both $\Delta^2\text{H}$ and $\Delta^{18}\text{O}$ are influenced by (ii) tissue AWA, and (iii) evaporation and sublimation enrichments. The lower the AWA, the higher the H-exchange effect and evaporation and sublimation fractionation, and the higher the $\Delta^2\text{H}$ and $\Delta^{18}\text{O}$.

2 Materials and methods

2.1 Experiment 1: testing the overall H-exchange effect

For evaluating the overall effect of H exchange during both rehydration and extraction on the $\Delta^2\text{H}$ and $\Delta^{18}\text{O}$ values of CVD-extracted water, we conducted an incubation experiment by adding different amounts of strongly depleted reference water ($\delta^2\text{H}$: -465.9‰ ; $\delta^{18}\text{O}$: -174.2‰ ; hereafter referred to as δ_{ref}) to the same amount of different dried materials with and without exchangeable H (experiment 1, Fig. 1). The isotopically depleted water was used to make potential isotope effects more evident.

The materials with exchangeable H included stem pieces, stem powder, purified stem cellulose powder, and twig pieces. The stem materials were obtained from the xylem of a trunk disc of a mature *Larix sibirica* grown in Siberia (27 cm diameter, 4 cm thickness). We cut a portion of xylem (a mixture of sap wood and heart wood) from the trunk disc and oven dried it to constant weight before preparing for stem pieces, powder, and cellulose materials. Stem pieces were prepared by cutting the xylem of the disc into 4 mm cubes, while the stem powder was prepared by grinding the same material to a homogeneous powder using a steel-ball mill (MM400, Retsch GmbH, Haan, Germany). The stem cellulose was then extracted from ground stem powder (Schuler et al., 2022). The twig pieces were obtained from five young *Larix decidua* trees growing in a forest in Birmensdorf, Switzerland (47.36° N, 8.45° E). Two twigs per tree were collected in the morning, outer bark and phloem removed. The twig xylem was oven dried at 60 °C for 24 h then cut into 2 mm pieces.

The materials without exchangeable H were cellulose triacetate and caffeine. The cellulose triacetate ($\text{C}_{40}\text{H}_{54}\text{O}_{27}$; Sigma-Aldrich, St. Louis, MO, USA, prod. no. 22199) is a white granule. The caffeine material is $\geq 99\%$ caffeine anhydrous ($\text{C}_8\text{H}_{10}\text{N}_4\text{O}_2$; Fluka Chemie AG, Buchs, Switzerland, prod. no. 27600) in a white crystalline powder form. All H atoms in cellulose triacetate and caffeine are bound to carbon in a form of methyl group ($-\text{CH}_3$) and thus non-exchangeable with vapour or liquid water. Cellulose triacetate and caffeine are anhydrous chemicals and were stored in a cool/dry place.

For each material and each different water amount (50, 100, 200, 400, 600, 800, and 1200 μL), three replicates were prepared by transferring 200 mg of each material into 12 mL gas-tight glass vials (Exetainer; Labco, Lampeter, UK). The amount of material was chosen because of practical considerations and to generate a range of water/biomass ratios. The materials were oven dried in the vials with caps open at 60 °C for 24 h to remove any remaining moisture in the material. In total, the materials with exchangeable H were oven dried at 60 °C for at least 48 h, and the materials without exchangeable H were oven dried at 60 °C for 24 h because they are dried chemicals and were properly stored. After drying, the vials were closed and the samples were cooled down to room temperature, then we opened the cap to inject the reference water into the vials. Depending on the form of the material, the low water amounts (e.g. 50 and 100 μL) could not fully wet the material, whereas the high water amounts (e.g. $> 400 \mu\text{L}$) oversaturated the material. Then, the vials were sealed and the materials were incubated for 24 h at room temperature. According to Chen et al. (2020), this incubation time is thought to lead to a full exchange of H atoms between materials and reference water. As a control, the experiment was repeated without any material by adding only the range of reference water into the vial. All samples were frozen at -20 °C and then extracted with CVD. Δ was calculated as $\delta_{\text{CVD}} - \delta_{\text{ref}}$.


In addition, to evaluate how effective our drying procedure was and how opening the cap for injecting the reference water influenced the dried samples, we conducted a drying test. In the test, 200 mg stem pieces, stem powder, and caffeine with five replicates were transferred into the vials, then caps closed and weighted. We oven dried the samples without caps for 48 h either at 60 or at 105 °C. Every 12 h during the drying, we closed the caps in the oven then transferred the vials to a balance for weighting. After the last weighting, we opened the cap of the vials for 5 s in the lab to simulate the procedure of injecting the reference water. Then the cap closed and samples were weighted again. The percentage of moisture that was removed from the initial sample at different temperatures and hours and the percentage of moisture that was absorbed by the dried sample in the 5 s were calculated (Table S2 in the Supplement).

2.2 Experiment 2: testing the effects of the absolute water amount and relative water content


For separating the H exchange effects on $\Delta^2\text{H}$ and $\Delta^{18}\text{O}$ values occurring during rehydration from those occurring during extraction, we extracted water from a range of rehydrated stem segments with different sizes (experiment 2, Fig. 1).

Large stem segments (10 cm \times 2.5 cm \times 0.3 cm) were cut from the *L. sibirica* disc (see Sect. 2.1) and oven-dried at 60 °C for 24 h before the experiment. We firstly determined the saturated water content of the large stem segments by weighing them before and after they were soaked in ex-


Experiment 1: Testing the overall H-exchange effect



Experiment 2: Testing the effects of the absolute water amount and relative water content



Experiment 3: Testing sublimation and evaporation effects



Experiment 4: Extraction tests using samples at isotope natural abundance




Figure 1. Overview of the experimental designs. Δ , isotopic offset; δ_{CVD} , isotope composition of CVD-extracted water; δ_{ref} , isotope composition of the reference water; $\delta_{\text{ref after rehyd}}$, isotope composition of the reference water after rehydration; $\delta_{\text{ref after evap}}$, isotope composition of the reference water after the evaporation; $\delta_{\text{CVD_ave}}$, the average value of isotope composition of the CVD-extracted water; δ_{tap} , isotope composition of the tap water; RT, room temperature; RH, relative humidity.

cess of deionized water for 24 h. The saturated water content was determined for estimating the AWA of the following smaller stem segment samples. The large stem segments were then oven dried again at 60 °C for 24 h and cut into smaller stem segments with different sizes, aiming to generate a range of samples with a narrow RWC (ca. 40 %), but varying in their AWA from 50–1200 μL . The sizes of the small segments were determined based on the proportional relationship between the size and the saturated water content of the large stem segments. Three replicates were produced for each size. Then, the small stem segments were separately soaked in excess of the isotopic depleted reference water (ca. 25 mL) in sealed glass vials at room temperature for 24 h, following Chen et al. (2020). During the 24 h rehydration, the isotope ratios of the original reference water equilibrated with the exchangeable H in the small stem segments. Thus, by the end of the rehydration, the isotope ratios of water in the small stem segments are assumed equal to the isotope ratio of the reference water after rehydration ($\delta_{\text{ref after rehyd}}$) and not to be equal to the original reference water (δ_{ref}). Subsequently, the small stem segments were taken out of the reference water, the surface water shaken off, and then immediately sealed in vials and frozen at $-20\text{ }^{\circ}\text{C}$ to prevent evaporation. A sample of the reference water after the rehydration was taken from each tube for determining $\delta_{\text{ref after rehyd}}$. The AWA and the RWC of each small stem segment sample were calculated by comparing the sample weights before and after the extraction. Here, the RWC is defined as (wet weight – dry weight)/wet weight, its unit is parts per hundred (%). In experiment 2, Δ was calculated as $\delta_{\text{CVD}} - \delta_{\text{ref after rehyd}}$. This is because it represents the changes in isotope ratios of the water that occurred during the CVD extraction alone.

2.3 Experiment 3: testing sublimation and evaporation effects

For evaluating the potential effects of sublimation and evaporation during CVD extraction on the isotope ratios of the extracted water, different amounts of reference water (50–1200 μL range as above) were added directly into the u-shaped water collection tubes (i.e. in the cold trap) of the CVD extraction system without sample material and then extracted for 2 h following a standard procedure (see Sect. 2.5). The water in the collection tubes was first frozen by the liquid nitrogen cold trap to avoid potential evaporation effects, then the collection tubes were attached to the extraction line for CVD extraction. The experimental design allows us to exclude isotope fractionation related to distillation and condensation occurring when water in the sample vial was relocated to the collection tube (i.e. U-tube), but captures isotope fractionation related to sublimation of the frozen water under the vacuum and evaporation during thawing of water after CVD extraction (experiment 3, Fig. 1). Δ was calculated as $\delta_{\text{CVD}} - \delta_{\text{ref}}$.

For isolating the evaporation effect from the sublimation effect, we also added different amounts of reference water (50–1200 μL range as above) into 2 mL glass vials inside a climate chamber (PGR15, Conviron, Manitoba, Canada). The samples were then left to evaporate with lids open at 25 °C and 50 % relative humidity (RH) for 2 h. The samples in the 2 mL glass vials did not require the CVD extraction. The exposed surface area of the water in the 2 mL glass vials was about 64 mm². Three replicates per water amount were performed for both tests (experiment 3, Fig. 1). Notably, CVD extraction was not used for the evaporation test in the climate chamber, thus the Δ was calculated as the difference between the isotope ratios of the reference water after and before the evaporation (i.e. $\delta_{\text{ref after evap}} - \delta_{\text{ref}}$).

2.4 Experiment 4: extraction tests using samples at isotope natural abundance

In order to test whether the CVD-related isotope fractionation, which was observed with labelled water, can also be seen at natural isotope abundance, we performed extractions with tap water and fresh *L. decidua* twigs with natural isotope abundance to compare with results derived from tests with isotopically depleted reference water. For the tap water (δ_{tap} ; $\delta^2\text{H}$: $-73.91\text{ }‰$, $\delta^{18}\text{O}$: $-14.43\text{ }‰$), the different amounts of water (i.e. 50, 100, 200, 400, 600, 800, and 1200 μL) were added into the vials and kept frozen at $-20\text{ }^{\circ}\text{C}$ until CVD extraction (experiment 4, Fig. 1). The twigs were collected from five young *L. decidua* trees growing in a forest in Birmensdorf, Switzerland (see Sect. 2.1). The twig xylem was cut to different lengths (1, 2, 3, 4, 5, 6, 7, and 8 cm) and sealed in vials. The AWA and RWC of each twig sample were determined by weighing before and after CVD extraction. Three replicates were performed per water amount and twig length (experiment 4, Fig. 1). For the extraction with tap water, the Δ was calculated as $\delta_{\text{CVD}} - \delta_{\text{tap}}$. The isotope ratios of the source water of the *L. decidua* twigs before extraction were unknown, thus the Δ in this case was calculated as the difference between the δ_{CVD} and the average value of δ_{CVD} (i.e. $\delta_{\text{CVD}} - \delta_{\text{CVD_ave}}$).

2.5 Water extraction and stable isotope analysis

We extracted water using a CVD setup similar to the one described by Orłowski et al. (2013). A schematic overview of this setup is shown in Fig. S1 in the Supplement. During extraction, the pressure inside the system was maintained below 0.05 mbar using a vacuum pump. The sample tubes were additionally blocked by PP fiber filters (Nozzle protection filter, Socorex Isba SA, Ecublens, Switzerland) when water was extracted from powder/pieces of the selected materials to avoid particles being drawn into the U-tubes with the extracted water or the vacuum pump (BS2212, Brook Crompton Ltd, Doncaster, UK). The samples in the sample tubes were heated in 80 °C water, and the extracted water was con-

densified and trapped in the collection tubes by liquid nitrogen. The extraction was maintained for 2 h to achieve a complete extraction (West et al., 2006). After the extraction, the vacuum inside the system was released by adding dry nitrogen gas until atmospheric pressure conditions were reestablished. Then the collection tubes with frozen water samples were detached from the system and sealed with rubber plugs. The water in the collection tubes was thawed at room temperature. During this process, the evaporated water vapour usually condensed to form very small water droplets on the inside walls of the collection tube. We consolidated and collected as many of these small water droplets as possible and transferred them into glass vials (350 μL or 2 mL, depending on the extracted water amount; Infochroma AG, Goldau, Switzerland) using a pipette. Syringes and 0.45 μm nylon filters were used if the extracted water appeared to be turbid. The samples were stored at -20°C before and after the extraction.

The $\delta^2\text{H}$ and $\delta^{18}\text{O}$ of water samples were measured with a high temperature conversion elemental analyser coupled to a Delta^{Plus} XP isotope ratio mass spectrometer (TC/EA-IRMS; Finnigan MAT, Bremen, Germany). The $\delta^2\text{H}$ and $\delta^{18}\text{O}$ in the materials without exchangeable H (i.e. cellulose triacetate and caffeine) and $\delta^{18}\text{O}$ in the bulk organic matter of the materials with exchangeable H were measured with a vario PYRO cube (Elementar Analysensysteme GmbH, Langensfeld, Germany) coupled to a Delta^{Plus} XP IRMS. Isotope ratios are reported in per mille (‰) relative to Vienna Standard Mean Ocean Water (VSMOW). Calibration versus the international standards was achieved by analysis of a range of certified water of different isotope ratios, resulting in a precision of analyses of 2 ‰ for $\delta^2\text{H}$ and 0.3 ‰ for $\delta^{18}\text{O}$. The non-exchangeable $\delta^2\text{H}$ ($\delta^2\text{H}_{\text{ne}}$) of the materials with exchangeable H was determined by pre-treating the materials with a high-temperature water vapour equilibration method according to Schuler et al. (2022). The $\delta^2\text{H}$ and $\delta^{18}\text{O}$ values of the materials used in the experiments are shown in Table S1 in the Supplement.

2.6 Statistical analyses

To determine if the sample water $\delta^2\text{H}$ and $\delta^{18}\text{O}$ values were significantly different before and after the CVD extraction, we performed one-sample t-tests for $\Delta^2\text{H}$ and $\Delta^{18}\text{O}$ values to test if they were significantly different from 0. We fit relationships between $\Delta^2\text{H}$ and $\Delta^{18}\text{O}$ as a function of AWA using inversely proportional models, and the relationships between $\Delta^2\text{H}$ and $\Delta^{18}\text{O}$ as a function of RWC using linear models for the best description of the relationships. The relationships between $\Delta^2\text{H}$ and $\Delta^{18}\text{O}$ as a function of RWC were also tested using a linear mixed-effects model with AWA category (AWA < 400 or AWA > 400 μL) as a random effect with the lmerTest package (Kuznetsova et al., 2017). All statistics were performed using R version 4.0.4 (R Core Team, 2021).

3 Results and discussion

3.1 H exchange occurs between sample exchangeable H and water during rehydration

Experiment 1 revealed an inversely proportional relationship between the AWA and the $\Delta^2\text{H}$ and $\Delta^{18}\text{O}$ values of the water derived from material with or without exchangeable H (Fig. 2). In other words, the lower the AWA, the higher the $\Delta^2\text{H}$ and $\Delta^{18}\text{O}$ values. For both $\Delta^2\text{H}$ and $\Delta^{18}\text{O}$, the differences among samples at AWA > 600 μL were much smaller compared to those at AWA < 600 μL . Noticeably, $\Delta^2\text{H}$ of the water derived from materials with exchangeable H showed a more pronounced pattern, which reached about 150 ‰ at 50 μL AWA, compared to those without exchangeable H (Fig. 2a and b). In contrast, the $\Delta^2\text{H}$ pattern of the water derived from materials without exchangeable H was similar to that of the pure reference water, which reached about 30 ‰ at 50 μL AWA (Fig. 2b). For $\Delta^{18}\text{O}$, the patterns were similar among materials with and without exchangeable H, showing an average decrease from 32 ‰ at 50 μL to 1.7 ‰ at 1200 μL (Fig. 2c and d).

The results of experiment 1 suggest that the H exchange during rehydration is mainly responsible for the large $\Delta^2\text{H}$ differences between materials with and without exchangeable H at the low water amounts. This can likely be explained by an isotope mass balance between the exchangeable H in the samples (natural isotope abundance) and H atoms in the reference water (isotopically depleted), with the impact of the exchangeable H of a constant sample size (200 mg) becoming less noticeable as the amount of water increases (from 50 to 1200 μL). Our results thus provide empirical evidence that H exchange occurs during rehydration of material and that this artefact must be considered in rehydration experiments (Chen et al., 2020; Zhao et al., 2022). However, our experiments cannot provide any evidence for the de- and rehydration of plants under natural conditions (i.e. changes in plant relative water content) which can potentially affect isotope composition of plant water (De Deurwaerder et al., 2020; Barbata et al., 2022). The effect of rehydration under field conditions could be quantified by additional experiments using water with an isotopic composition distinct from that normally expected in soils.

Moreover, the $\Delta^2\text{H}$ of the stem and twig pieces were higher than that of the stem and stem cellulose powder when AWA > 600 μL (Fig. 2a), whereas no differences were observed for $\Delta^{18}\text{O}$ (Fig. 2c). At a first glance, this is unexpected because woody pieces should be less prone to H exchange due to lower surface area and because the OH groups are locked in the matrix of the woody structure (Sepall and Mason, 1961). A possible explanation for the observed $\Delta^2\text{H}$ discrepancies might be an isotopic difference in the exchangeable H of pieces and powdered samples before the start of the experiment. However, if this was true, the effect should be more visible at lower water amounts rather than higher water




Figure 2. Absolute water amount effects on $\Delta^2\text{H}$ and $\Delta^{18}\text{O}$ values (**a, b** and **c, d**, respectively) of water derived from materials with and without exchangeable H (**a, c** and **b, d**, respectively), of a constant weight (200 mg). Material with exchangeable H differ in their structural composition (pieces vs. powder). “Reference water” reflects a control performed with water but without any material. $\Delta = \delta_{\text{CVD}} - \delta_{\text{ref}}$ (see Sect. 2.1). The dashed lines represent $\Delta = 0$, with grey shading to represent the SD of the reference water isotope analyses (2‰ for $\delta^2\text{H}_{\text{ref}}$; 1.5‰ for $\delta^{18}\text{O}_{\text{ref}}$). The curves are inversely proportional fits. Mean values ± 1 SD are shown ($n = 3$).

amounts, as observed in this study. We therefore can conclude that the sample matrix may also influence H exchange in rehydration experiments, but that the underlying mechanisms remain speculative. Besides, we additionally tested whether the drying procedure (oven drying at 60 °C) in experiment 1 influenced our results. The test showed that all the tested materials were dried to a constant weight after 36 and 12 h at 60 and 105 °C, respectively, despite more moisture being removed at 105 °C (Table S2). Interestingly, after opening the cap for 5 s to simulate injection of reference water, the amount of absorbed lab water vapour relative to the dry weight was about 1 % higher for the samples which were dried at 105 °C than the samples which were dried at 60 °C (Table S2). These results indicate that a complete drying is very difficult in our case, regardless of drying temperature. However, the test provides evidence that only small amounts of remaining moisture remain in the sample after drying and that the reabsorption should not affect the isotopic results of our study, because strongly depleted reference water was used to amplify the isotopic effect.

Unexpectedly, $\Delta^2\text{H}$ values of the pure reference water and of material without exchangeable H, as well as $\Delta^{18}\text{O}$ values, all show a dependency on AWA (Fig. 2b–d). This observation indicates that Δ values can be biased even when no exchangeable H in plant material is available, and suggests that other factors than H exchange play an important role in biasing isotopic results of CVD-extracted water.

3.2 Absolute water amount not relative water content causes isotope fractionation during CVD extraction

To better understand the unknown isotopic effects during extraction, we performed experiment 2, where we separated the effects between rehydration and extraction on CVD-extracted water along a similar AWA gradient (50–1600 μL), but in a narrow RWC range (36 %–45 %, Fig. 3). The results confirm the inversely proportional pattern of Δ along an AWA gradient as shown by experiment 1 (Fig. 2). More importantly, the results show that Δ values of water extracted from stem material that has been previously rehydrated were similar (Fig. 3a and c) to that extracted from materials with-

out any exchangeable H (Fig. 2b and d), when the isotope ratio of the reference water after rehydration ($\delta_{\text{ref after rehyd}}$) instead of the reference water was considered. This shows that the H-exchange effect during extraction (i.e. all steps after the rehydration procedure) is negligible and cannot be the main reason for observed isotopic offset between expected and measured isotope ratios in water of stem material (Bowers and Williams, 2022).

This finding is in contrast to the conclusions made by Chen et al. (2020), stating that stem water CVD extraction error could originate from a dynamic H exchange between sample and water during extraction. The reason for the lack of H-exchange effect during the 2 h CVD extraction itself is probably because the duration for the H exchange was too short to fully exchange. Previous research shows that a large portion of the water in the sample is expected to be extracted within the first 30 min of extraction (Orlowski et al., 2013; West et al., 2006). Thus, there may not be a strong H exchange occurring between the water and sample tissue, especially when a large portion of exchangeable H atoms are not freely accessible at the extraction temperature of 80 °C (Sepall and Mason, 1961). The duration of rehydration might be another factor that potentially affected our results of the rehydration experiments (experiment 1 and 2). According to Chen et al. (2020), a complete equilibration between sample exchangeable H and reference water during rehydration can be achieved within 24 h, given water of a sufficiently large volume. In experiment 2 of our study, the rehydration was also conducted in an excess of reference water for 24 h. The AWA-dependence of $\Delta^2\text{H}$ values of stem segments in experiment 2 (Fig. 3a) was found to be similar to that of the pure reference water and materials without exchangeable H extractions in experiment 1 (Fig. 2b). This likely confirmed that a full equilibrium was achieved during our 24 h rehydration. Nevertheless, the duration, sample size, and amount of water should be considered in future rehydration studies.

We found a weak, positive linear trend in $\Delta^2\text{H}$ with RWC ($r^2 = 0.28$, $p = 0.02$, Fig. 3b), but not for $\Delta^{18}\text{O}$ ($p = 0.08$, Fig. 3d). The trend in $\Delta^2\text{H}$ can be explained by the fact that samples with a smaller AWA ($< 400 \mu\text{L}$) had a ca. 2 % higher RWC compared to those with a higher AWA ($> 400 \mu\text{L}$). The relationships between both $\Delta^2\text{H}$ and $\Delta^{18}\text{O}$ as a function of RWC were not statistically significant when AWA was considered as a random factor (AWA with 2 levels: AWA < 400 and AWA $> 400 \mu\text{L}$; Fig. 3 and Table S3 in the Supplement). This indicates that the $\Delta^2\text{H}$ and $\Delta^{18}\text{O}$ values are dependent on AWA rather than on RWC of the sample. Our result is not consistent with Chen et al. (2020), who found a significant positive correlation of stem RWC with the $\Delta^2\text{H}$ in a rehydration experiment and further recommended determining sample RWC for correcting CVD artefacts. However, our results are well supported by the recent study of Zhao et al. (2022), who CVD-extracted water from rehydrated samples of 12 woody plant species and analysed Δ values. The RWC of their samples ranged from 30 %–60 % and no significant

relationships were found between both $\Delta^2\text{H}$ and $\Delta^{18}\text{O}$ and RWC. Therefore, the relatively smaller sample RWC range in our study (ca. 10 %) compared to that in Chen et al. (2020), i.e. ca. 20 % for most samples, could not be the reason for the inconsistency of the two studies. Further, given that AWA is typically not reported, we could not test whether the AWA or RWC caused the effect in previous studies. We highlight that the AWA rather than the RWC of a sample should be considered to potentially correct for Δ offsets, and that future studies should report AWA in order to quantify the CVD-induced isotopic biases across laboratories.

3.3 On the processes causing isotope fractionation during CVD extraction

In order to better understand the processes leading to isotope fractionation during CVD, we performed an additional experiment (experiment 3). In this experiment, different amounts of reference water were either injected directly into the collection tubes then subject to CVD extraction, or subject to controlled evaporative conditions in a climate chamber for 2 h. Interestingly, experiment 3 shows that both $\Delta^2\text{H}$ and $\Delta^{18}\text{O}$ values of reference water, that was directly transferred to the cold trap (i.e. U-tubes), also followed the inversely proportional pattern as a function of AWA (Fig. 4a). At the lowest AWA, $\Delta^2\text{H}$ reached about 19 ‰, while $\Delta^{18}\text{O}$ reached about 9 ‰. For AWA $> 600 \mu\text{L}$, the average $\Delta^2\text{H}$ and $\Delta^{18}\text{O}$ were -1.5 ‰ and -2.3 ‰, respectively, and both were significantly different from 0 ($t = -3.13$ and -3.34 , $\text{df} = 8$, $p < 0.01$). We suppose that the negative $\Delta^2\text{H}$ and $\Delta^{18}\text{O}$ values for the AWA $> 600 \mu\text{L}$ were purely caused by the analytical uncertainty, because no incomplete extraction could occur given that the reference water was added directly into the collection tube. The same inversely proportional pattern was observed when water was evaporated under controlled conditions in a climate chamber, whereby $\Delta^2\text{H}$ reached about 46 ‰ and $\Delta^{18}\text{O}$ reached about 23 ‰ at the lowest AWA. For AWA $> 600 \mu\text{L}$, average $\Delta^2\text{H}$ was about 3 ‰ and significantly different from 0 ($t = 12.15$, $\text{df} = 8$, $p < 0.01$). No significant difference from 0 was found for $\Delta^{18}\text{O}$ ($t = 0.23$, $\text{df} = 8$, $p = 0.8$). By comparing these results with the one of 50 μL of reference water that was extracted from the sample tube (Fig. 2b), the sublimation and evaporation isotope fractionation account for 59 % and 38 % of the observed $\Delta^2\text{H}$ and $\Delta^{18}\text{O}$, respectively. The unaccounted isotope fractionation may originate from distillation and condensation, when water in the sample tube was relocated to the water collection tube.

Two potential reasons can explain this water amount-dependent pattern of Δ . On the one hand, a bi-directional exchange between water droplets and water vapour may lead to an isotopic equilibration between the two sources, similar to observations at the leaf level (Lehmann et al., 2020; Goldsmith et al., 2017). This process is expected to occur when the extracted water drop was thawing inside the water




Figure 3. Effects of absolute water amount and relative water content (a, c and b, d, respectively) on $\Delta^2\text{H}$ and $\Delta^{18}\text{O}$ of water derived from the stem segments (a, b and c, d, respectively). The stem segments were rehydrated in excess of the reference water followed by a CVD extraction. $\Delta = \delta_{\text{CVD}} - \delta_{\text{ref after rehyd}}$ (see Sect. 2.2). The relationship between AWA and Δ was fit with an inversely proportional function; the relationship between RWC and Δ was fit with a linear regression. The dashed lines represent $\Delta = 0$. Different colours were used to visually separate data with water amount < 400 (red) and > 400 μL (blue).




Figure 4. Absolute water amount effects on $\Delta^2\text{H}$ (a) and $\Delta^{18}\text{O}$ (b) of the reference water that was directly added into the water collection tube before CVD extraction (blue), or evaporated in a climate chamber (red). For the former, $\Delta = \delta_{\text{CVD}} - \delta_{\text{ref}}$; for the latter, $\Delta = \delta_{\text{ref after evap}} - \delta_{\text{ref}}$ (see Sect. 2.3). The dashed lines represent $\Delta = 0$, with grey shading to represent the SD of the reference water isotope analyses (2‰ for $\delta^2\text{H}_{\text{ref}}$; 1.5‰ for $\delta^{18}\text{O}_{\text{ref}}$). The curves are inversely proportional fits. Mean values ± 1 SD are shown ($n = 3$).

collection tube after extraction. The equilibration effect was presumably large on our extracted water, because of a high isotopic discrepancy between the water vapour of the laboratory (at natural isotope abundance) and the isotopically depleted reference water. The effect is expected to increase as the water pool size decreases, because the smaller the water droplets, the larger the ratio of water vapour volume to water droplets volume in the water collection tube. Therefore, isotope ratios of smaller water amounts were subject to greater influence of the isotopic composition of the laboratory water vapour. On the other hand, the sublimation, evaporation, and/or exchange with surrounding water vapour occurs at the surface of water drops (Stewart, 1975). The smaller the water drop, the larger the ratio of the enriched surface water to the total water drop volume. Thus, the mixing effect of enriched surface water and the rest of the water body would be greater for smaller water drops.

Taken together, our results clearly show that the inversely proportional pattern of Δ along AWA is partly influenced by the isotope fractionation during water sublimation and evaporation during CVD extraction. We therefore conclude that isotope fractionation rather than H-exchange effects with plant material during extraction itself should be considered as factors during CVD extraction of plant water. It should be noted that although our experiments were mainly conducted on samples of *Larix*, the AWA-dependent patterns of Δ were consistent among extractions with different materials and even pure water. This indicates that the CVD biases we observed on *Larix* may also be applicable to other taxa. The magnitude of the biases may be slightly different among species (Zhao et al., 2022; Chen et al., 2020; De La Casa et al., 2022), but the AWA-dependency of the biases should show consistency. We also want to highlight that the observed isotope fractionation likely depends on the setup (e.g. system volume) of the CVD extraction system. Therefore, further laboratory comparisons for plant water extraction using different woody and herbaceous plant species are needed for constraining this extraction bias.

3.4 Relevance for isotope fractionation during CVD extraction for samples at natural isotope abundance

The large isotopic discrepancy between the strongly depleted reference water and the plant tissue organic matter allowed us to investigate the processes and mechanisms driving isotope fractionation during CVD extraction. However, plant water at natural isotope abundance is typically more enriched compared to our reference water ($\delta^2\text{H}$: -465.9‰ ; $\delta^{18}\text{O}$: -174.2‰) and therefore the observed isotopic effects of this study are likely less pronounced at natural isotope abundance. This was true, as shown by the results of experiment 4; the inversely proportional pattern was less pronounced or absent with plant samples (average values in $\delta^2\text{H}$ and $\delta^{18}\text{O}$ of extracted water: -58.16‰ and -7.84‰ , respectively), or

when tap water ($\delta^2\text{H}$: -73.91‰ ; $\delta^{18}\text{O}$: -14.43‰) was used (Fig. 5). For the pure tap water extraction, $\Delta^2\text{H}$ decreased from 3‰ at $50\text{ }\mu\text{L}$ to -3.5‰ at $1200\text{ }\mu\text{L}$, while $\Delta^{18}\text{O}$ decreased from 2‰ at $50\text{ }\mu\text{L}$ to -0.2‰ at $1200\text{ }\mu\text{L}$ (Fig. 5a and c).

The possible explanations for the large difference in the magnitude of the Δ pattern between the reference (Fig. 2b and d) and tap water extractions (Fig. 5a and c) are that the isotope fractionation during CVD extraction is dependent on the initial water isotope ratios and/or the influence from isotope ratios of the surrounding water vapour. Regarding the former, water evaporation tests using water with different isotope ratios ($\delta^2\text{H}$ range: -14.7‰ to -57.5‰ ; $\delta^{18}\text{O}$ range: 3.4‰ to -7.8‰) showed that the $\delta^2\text{H}$ and $\delta^{18}\text{O}$ values of the initial water have little influence on water evaporation isotope fractionation conducted by Hu et al. (2009). In other words, for water with different initial isotope ratios evaporating in the same conditions, the changes in $\delta^2\text{H}$ or $\delta^{18}\text{O}$ would be almost the same. Therefore, the initial water isotope ratios would not be the reason for the significant Δ difference between the reference water and the tap water extraction.

It is therefore possible that the difference in Δ between reference water and tap water extractions is caused by the exchange of extracted water with the water vapour in the laboratory. The isotope ratios of the laboratory water vapour are closer to that of the tap water and twig water, but differ greatly compared to the depleted reference water. Therefore, under these conditions, the initial isotopic signature of the extracted water determines the magnitude of the isotope signature of the extracted water after the exchange with laboratory water vapour and therefore the observed Δ values. This suggests that the observed isotopic variations in our experiments with strongly isotopically depleted reference water might have been less pronounced if water closer to natural isotope abundance would have been used.

For the fresh *L. decidua* twigs water extraction, the AWA-dependent pattern was evident for $\Delta^{18}\text{O}$, with average values decreasing from approximately 1‰ at the lowest water amount to approximately -0.2‰ at the highest water amount (Fig. 5c). In contrast, $\Delta^2\text{H}$ ranged between -6.5‰ and 6.1‰ but without showing a clear inversely proportional pattern (Fig. 5a). While we found no clear explanation for the absence of the AWA dependency for $\Delta^2\text{H}$ in this experiment, we can only speculate that the expected pattern was hidden by processes shaping hydrogen but not oxygen isotopic variations such as H-exchange effects between water vapour and extracted water with similar isotopic compositions. Our results also showed that the Δ values obtained from the extraction of water at natural isotope abundance (Fig. 5a and c) are much smaller compared to those obtained from the extractions of the strongly depleted reference water (Figs. 2 and 3). In addition, we did not find a significant correlation between RWC and $\Delta^2\text{H}$ or $\Delta^{18}\text{O}$ for the *L. decidua* twigs at a RWC range between 45% and 57% (Fig. 5b and d). These results




Figure 5. Absolute water amount and relative water content effects (a, c and b, d, respectively) on $\Delta^2\text{H}$ and $\Delta^{18}\text{O}$ (a, b and c, d, respectively) of water at natural isotope abundance. For water of *Larix decidua* twig xylem, $\Delta = \delta_{\text{CVD}} - \delta_{\text{CVD_ave}}$; while for the tap water, $\Delta = \delta_{\text{CVD}} - \delta_{\text{tap}}$ (see Sect. 2.4). The relationship between water amount and Δ was fit with an inversely proportional function. The dashed lines represent $\Delta = 0$. For (a) and (c), mean values ± 1 SD are shown ($n = 3$).

suggest again that the $\Delta^2\text{H}$ and $\Delta^{18}\text{O}$ of the CVD-extracted water is dependent on AWA rather than on RWC.

4 Conclusions

In conclusion, we provide strong evidence that H exchange with organic H in the sample, despite significantly influencing the $\delta^2\text{H}$ of water during the sample rehydration process, is not the main driver of the broadly observed negative $\Delta^2\text{H}$ value during CVD extraction. Instead, we identified a significant CVD artefact when water is present in small amounts, particularly when $\delta^2\text{H}$ and $\delta^{18}\text{O}$ of the water were below natural isotope abundance. This is linked to the increase of sublimation and evaporation enrichments with the decrease of water amount, as well as the mixing between the extracted water and laboratory water vapour, rather than the effect of sample RWC on the H exchange during extraction. From our results, both $\Delta^2\text{H}$ and $\Delta^{18}\text{O}$ approached steady values and were close to zero when $> 600 \mu\text{L}$ of water was extracted. We therefore recommend extracting more than $600 \mu\text{L}$ of water, especially for studies using labelled water, to avoid large enrichment biases. However, the minimum amount of ex-

tracted water that is necessary to avoid isotopic biases might vary with the CVD setup and therefore laboratory comparison should be conducted. Our results have implications for studies using stable isotopes of water in plant tissue to determine plant water sources (e.g. Evaristo and McDonnell, 2017), trace water through soils with high organic material content (e.g. Koeniger et al., 2016; Sprenger et al., 2016), and reconstruct climate patterns using tree ring tissue (e.g. Loader et al., 2007; Lehmann et al., 2021).

Code availability. The code used for all statistical analyses is available upon request.

Data availability. Data are available from the corresponding author upon reasonable request.

Supplement. The supplement related to this article is available online at: <https://doi.org/10.5194/hess-26-5835-2022-supplement>.

Author contributions. HD, PS, GRG, and MML designed the experiments and HD carried them out. HD prepared the paper with contributions from all co-authors.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Acknowledgements. We acknowledge the technical assistance by Manuela Oettli and Oliver Rehmann at WSL. Haoyu Diao acknowledges a scholarship from the Joint PhD Training Program, University of Chinese Academy of Sciences.

Financial support. This research has been supported by the SNSF Ambizione project "TreeCarbo" (grant no. 179978, granted to Marco M. Lehmann) and the USDA-NIFA (grant no. 2020-67014-30917, granted to Gregory R. Goldsmith).

Review statement. This paper was edited by Markus Weiler and reviewed by two anonymous referees.

References

- Allen, S. T. and Kirchner, J. W.: Potential effects of cryogenic extraction biases on plant water source partitioning inferred from xylem water isotope ratios, *Hydrol. Process.*, 36, e14483, <https://doi.org/10.1002/hyp.14483>, 2022.
- Allen, S. T., Kirchner, J. W., Braun, S., Siegwolf, R. T. W., and Goldsmith, G. R.: Seasonal origins of soil water used by trees, *Hydrol. Earth Syst. Sci.*, 23, 1199–1210, <https://doi.org/10.5194/hess-23-1199-2019>, 2019.
- Barbeta, A., Gimeno, T. E., Clavé, L., Fréjaville, B., Jones, S. P., Delvigne, C., Wingate, L., and Ogée, J.: An explanation for the isotopic offset between soil and stem water in a temperate tree species, *New Phytol.*, 227, 766–779, <https://doi.org/10.1111/nph.16564>, 2020.
- Barbeta, A., Burrell, R., Martín-Gómez, P., Fréjaville, B., Devert, N., Wingate, L., Domec, J.-C., and Ogée, J.: Evidence for distinct isotopic compositions of sap and tissue water in tree stems: consequences for plant water source identification, *New Phytol.*, 233, 1121–1132, <https://doi.org/10.1111/nph.17857>, 2022.
- Bowers, W. H. and Williams, D. G.: Isotopic Heterogeneity of Stem Water in Conifers Is Correlated to Xylem Hydraulic Traits and Supports Multiple Residence Times, *Frontiers in Water*, 4, 861590, <https://doi.org/10.3389/frwa.2022.861590>, 2022.
- Brinkmann, N., Eugster, W., Buchmann, N., and Kahmen, A.: Species-specific differences in water uptake depth of mature temperate trees vary with water availability in the soil, *Plant Biol.*, 21, 71–81, <https://doi.org/10.1111/plb.12907>, 2019.
- Chen, Y., Helliker, B. R., Tang, X., Li, F., Zhou, Y., and Song, X.: Stem water cryogenic extraction biases estimation in deuterium isotope composition of plant source water, *P. Natl. Acad. Sci. USA*, 117, 33345–33350, <https://doi.org/10.1073/pnas.2014422117>, 2020.
- De Deurwaerder, H. P. T., Visser, M. D., Detto, M., Boeckx, P., Meunier, F., Kuehnhammer, K., Magh, R.-K., Marshall, J. D., Wang, L., Zhao, L., and Verbeeck, H.: Causes and consequences of pronounced variation in the isotope composition of plant xylem water, *Biogeosciences*, 17, 4853–4870, <https://doi.org/10.5194/bg-17-4853-2020>, 2020.
- de la Casa, J., Barbeta, A., Rodríguez-Uña, A., Wingate, L., Ogée, J., and Gimeno, T. E.: Isotopic offsets between bulk plant water and its sources are larger in cool and wet environments, *Hydrol. Earth Syst. Sci.*, 26, 4125–4146, <https://doi.org/10.5194/hess-26-4125-2022>, 2022.
- Ehleringer, J. R., Roden, J., and Dawson, T. E.: Assessing ecosystem-level water relations through stable isotope ratio analyses, in: *Methods in ecosystem science*, edited by: Sala, O. E., Jackson, R. B., Mooney, A. A., and Howarth, R. W., Springer, 181–198, https://doi.org/10.1007/978-1-4612-1224-9_13, 2000.
- Evaristo, J. and McDonnell, J. J.: Prevalence and magnitude of groundwater use by vegetation: a global stable isotope meta-analysis, *Sci. Rep.-UK*, 7, 44110, <https://doi.org/10.1038/srep44110>, 2017.
- Filot, M. S., Leuenberger, M., Pazdur, A., and Boettger, T.: Rapid online equilibration method to determine the D/H ratios of non-exchangeable hydrogen in cellulose, *Rapid Commun. Mass Sp.*, 20, 3337–3344, <https://doi.org/10.1002/rcm.2743>, 2006.
- Flanagan, L. B. and Ehleringer, J. R.: Stable Isotope Composition of Stem and Leaf Water: Applications to the Study of Plant Water Use, *Funct. Ecol.*, 5, 270–277, <https://doi.org/10.2307/2389264>, 1991.
- Gessler, A., Bächli, L., Rouholahnejad Freund, E., Treydte, K., Schaub, M., Haeni, M., Weiler, M., Seeger, S., Marshall, J., Hug, C., Zweifel, R., Hagedorn, F., Rigling, A., Saurer, M., and Meusburger, K.: Drought reduces water uptake in beech from the drying topsoil, but no compensatory uptake occurs from deeper soil layers, *New Phytol.*, 233, 194–206, <https://doi.org/10.1111/nph.17767>, 2022.
- Goldsmith, G. R., Muñoz-Villers, L. E., Holwerda, F., McDonnell, J. J., Asbjornsen, H., and Dawson, T. E.: Stable isotopes reveal linkages among ecohydrological processes in a seasonally dry tropical montane cloud forest, *Ecohydrology*, 5, 779–790, <https://doi.org/10.1002/eco.268>, 2012.
- Goldsmith, G. R., Lehmann, M. M., Cernusak, L. A., Arend, M., and Siegwolf, R. T. W.: Inferring foliar water uptake using stable isotopes of water, *Oecologia*, 184, 763–766, <https://doi.org/10.1007/s00442-017-3917-1>, 2017.
- Hu, H.-Y., Bao, W.-M., Wang, T., and Qu, S.-M.: Experimental study on stable isotopic fractionation of evaporating water under varying temperature, *Water Sci. Eng.*, 2, 11–18, 2009.
- Ingraham, N. L. and Shadel, C.: A comparison of the toluene distillation and vacuum/heat methods for extracting soil water for stable isotopic analysis, *J. Hydrol.*, 140, 371–387, [https://doi.org/10.1016/0022-1694\(92\)90249-U](https://doi.org/10.1016/0022-1694(92)90249-U), 1992.
- Koeniger, P., Gaj, M., Beyer, M., and Himmelsbach, T.: Review on soil water isotope-based groundwater

- recharge estimations, *Hydrol. Process.*, 30, 2817–2834, <https://doi.org/10.1002/hyp.10775>, 2016.
- Kuznetsova, A., Brockhoff, P. B., and Christensen, R. H. B.: lmerTest Package: Tests in Linear Mixed Effects Models, *J. Stat. Softw.*, 82, 1–26, <https://doi.org/10.18637/jss.v082.i13>, 2017.
- Lécuyer, C., Royer, A., Fourel, F., Seris, M., Simon, L., and Robert, F.: D/H fractionation during the sublimation of water ice, *Icarus*, 285, 1–7, <https://doi.org/10.1016/j.icarus.2016.12.015>, 2017.
- Lehmann, M. M., Goldsmith, G. R., Mirande-Ney, C., Weigt, R. B., Schönbeck, L., Kahmen, A., Gessler, A., Siegwolf, R. T. W., and Saurer, M.: The 18O -signal transfer from water vapour to leaf water and assimilates varies among plant species and growth forms, *Plant Cell Environ.*, 43, 510–523, <https://doi.org/10.1111/pce.13682>, 2020.
- Lehmann, M. M., Vitali, V., Schuler, P., Leuenberger, M., and Saurer, M.: More than climate: Hydrogen isotope ratios in tree rings as novel plant physiological indicator for stress conditions, *Dendrochronologia*, 65, 125788, <https://doi.org/10.1016/j.dendro.2020.125788>, 2021.
- Lin, G. and da S. L. Sternberg, L.: 31 – Hydrogen Isotopic Fractionation by Plant Roots during Water Uptake in Coastal Wetland Plants, in: *Stable Isotopes and Plant Carbon-water Relations*, edited by: Ehleringer, J. R., Hall, A. E., and Farquhar, G. D., Academic Press, San Diego, 497–510, <https://doi.org/10.1016/B978-0-08-091801-3.50041-6>, 1993.
- Loader, N. J., McCarroll, D., Gagen, M., Robertson, I., and Jalkanen, R.: Extracting climatic information from stable isotopes in tree rings, in: *Terrestrial Ecology*, edited by: Dawson, T. E. and Siegwolf, R. T. W., Elsevier, 25–48, [https://doi.org/10.1016/S1936-7961\(07\)01003-2](https://doi.org/10.1016/S1936-7961(07)01003-2), 2007.
- Meier-Augenstein, W., Kemp, H. F., Schenk, E. R., and Almirall, J. R.: Discrimination of unprocessed cotton on the basis of geographic origin using multi-element stable isotope signatures, *Rapid Commun. Mass Sp.*, 28, 545–552, <https://doi.org/10.1002/rcm.6811>, 2014.
- Mortimer, J., Lécuyer, C., Fourel, F., and Carpenter, J.: D/H fractionation during sublimation of water ice at low temperatures into a vacuum, *Planet. Space Sci.*, 158, 25–33, <https://doi.org/10.1016/j.pss.2018.05.010>, 2018.
- Nehemy, M. F., Benettin, P., Asadollahi, M., Pratt, D., Rinaldo, A., and McDonnell, J. J.: How plant water status drives tree source water partitioning, *Hydrol. Earth Syst. Sci. Discuss.* [preprint], <https://doi.org/10.5194/hess-2019-528>, 2019.
- Newberry, S. L., Nelson, D. B., and Kahmen, A.: Cryogenic vacuum artifacts do not affect plant water-uptake studies using stable isotope analysis, *Ecophysiology*, 10, e1892, <https://doi.org/10.1002/eco.1892>, 2017.
- Orlowski, N., Frede, H.-G., Brüggemann, N., and Breuer, L.: Validation and application of a cryogenic vacuum extraction system for soil and plant water extraction for isotope analysis, *J. Sens. Syst.*, 2, 179–193, <https://doi.org/10.5194/jsss-2-179-2013>, 2013.
- Poca, M., Coomans, O., Urcelay, C., Zeballos, S. R., Bodé, S., and Boeckx, P.: Isotope fractionation during root water uptake by *Acacia caven* is enhanced by arbuscular mycorrhizas, *Plant Soil*, 441, 485–497, <https://doi.org/10.1007/s11104-019-04139-1>, 2019.
- R Core Team: R: A Language and Environment for Statistical Computing, R Foundation for Statistical Computing, <https://www.R-project.org/> (last access: 15 November 2022), 2021.
- Schoppach, R. and Klaus, J.: Is cryogenic vacuum distillation reliable for wood water extraction?, EGU General Assembly 2019, 7–12 April 2019, Vienna, Austria, 4901, 2019.
- Schuler, P., Cormier, M.-A., Werner, R. A., Buchmann, N., Gessler, A., Vitali, V., Saurer, M., and Lehmann, M. M.: A high-temperature water vapor equilibration method to determine non-exchangeable hydrogen isotope ratios of sugar, starch and cellulose, *Plant Cell Environ.*, 45, 12–22, <https://doi.org/10.1111/pce.14193>, 2022.
- Sepall, O. and Mason, S. G.: Hydrogen exchange between cellulose and water: II. Interconversion of accessible and inaccessible regions, *Can. J. Chem.*, 39, 1944–1955, <https://doi.org/10.1139/v61-261>, 1961.
- Sprenger, M., Leistert, H., Gimbel, K., and Weiler, M.: Illuminating hydrological processes at the soil-vegetation-atmosphere interface with water stable isotopes, *Rev. Geophys.*, 54, 674–704, <https://doi.org/10.1002/2015RG000515>, 2016.
- Stewart, M. K.: Stable isotope fractionation due to evaporation and isotopic exchange of falling waterdrops: Applications to atmospheric processes and evaporation of lakes, *J. Geophys. Res.*, 80, 1133–1146, <https://doi.org/10.1029/JC080i009p01133>, 1975.
- Wen, M., Lu, Y., Li, M., He, D., Xiang, W., Zhao, Y., Cui, B., and Si, B.: Correction of cryogenic vacuum extraction biases and potential effects on soil water isotopes application, *J. Hydrol.*, 603, 127011, <https://doi.org/10.1016/j.jhydrol.2021.127011>, 2021.
- West, A. G., Patrickson, S. J., and Ehleringer, J. R.: Water extraction times for plant and soil materials used in stable isotope analysis, *Rapid Commun. Mass Sp.*, 20, 1317–1321, <https://doi.org/10.1002/rcm.2456>, 2006.
- White, J. W. C., Cook, E. R., Lawrence, J. R., and Wallace S. B.: The DH ratios of sap in trees: Implications for water sources and tree ring DH ratios, *Geochim. Cosmochim. Ac.*, 49, 237–246, [https://doi.org/10.1016/0016-7037\(85\)90207-8](https://doi.org/10.1016/0016-7037(85)90207-8), 1985.
- Zhao, L., Wang, L., Cernusak, L. A., Liu, X., Xiao, H., Zhou, M., and Zhang, S.: Significant difference in hydrogen isotope composition between xylem and tissue water in *Populus Euphratica*, *Plant Cell Environ.*, 39, 1848–1857, <https://doi.org/10.1111/pce.12753>, 2016.
- Zhao, P., Sprenger, M., Barzegar, R., Tang, X., and Adamowski, J.: Similar isotopic biases of plant stems bulk water from different water sources by cryogenic vacuum distillation demonstrated through rehydration experiments, *Geophys. Res. Lett.*, 49, e2021GL096474, <https://doi.org/10.1029/2021GL096474>, 2022.
- Zimmermann, U., Ehhalt, D., and Muennich, K. O.: Soil-water movement and evapotranspiration: Changes in the isotopic composition of the water, *Isotopes in Hydrology*, 567–585, <https://inis.iaea.org/search/searchsinglerecord.aspx?recordsFor=SingleRecord&RN=38061083> (last access: 15 November 2022), 1967.