

Mathematics, Physics, and Computer Science Faculty Articles and Research Science and Technology Faculty Articles and Research

5-19-2021

A New Method to Generate Superoscillating Functions and Supershifts

Yakir Aharonov Chapman University, aharonov@chapman.edu

Fabrizio Colombo Politecnico di Milano

Irene Sabadini Politecnico di Milano

Tomer Shushi Ben-Gurion University of the Negev

Daniele C. Struppa Chapman University, struppa@chapman.edu

See next page for additional authors

Follow this and additional works at: https://digitalcommons.chapman.edu/scs_articles

Part of the Other Mathematics Commons, and the Quantum Physics Commons

Recommended Citation

Aharonov Yakir, Colombo Fabrizio, Sabadini Irene, Shushi Tomer, Struppa Daniele C. and Tollaksen Jeff. 2021. A new method to generate superoscillating functions and supershifts. *Proc. R. Soc. A.* **477**: 20210020. http://doi.org/10.1098/rspa.2021.0020

This Article is brought to you for free and open access by the Science and Technology Faculty Articles and Research at Chapman University Digital Commons. It has been accepted for inclusion in Mathematics, Physics, and Computer Science Faculty Articles and Research by an authorized administrator of Chapman University Digital Commons. For more information, please contact laughtin@chapman.edu.

A New Method to Generate Superoscillating Functions and Supershifts

Comments

This is the submitted version of an article that later underwent peer review and was accepted for publication in *Proceedings of the Royal Society A*, volume 477, in 2021. The definitive publisher-authenticated version may differ and is available online at http://doi.org/10.1098/rspa.2021.0020.

Copyright

The authors

Authors

Yakir Aharonov, Fabrizio Colombo, Irene Sabadini, Tomer Shushi, Daniele C. Struppa, and Jeff Tollaksen

A new method to generate superoscillating functions and supershifts

Y. Aharonov^{*}, F. Colombo[†], I. Sabadini[†], T. Shushi[‡], D.C. Struppa[§], J. Tollaksen^{*}

January 19, 2021

Abstract

Superoscillations are band-limited functions that can oscillate faster than their fastest Fourier component. These functions (or sequences) appear in weak values in quantum mechanics and in many fields of science and technology such as optics, signal processing and antenna theory. In this paper we introduce a new method to generate superoscillatory functions that allows us to construct explicitly a very large class of superoscillatory functions.

AMS Classification: 26A09, 41A60.

Key words: Superoscillating functions, New method to generate superoscillations, Supershift.

1 Introduction

Superoscillating functions are band-limited functions that can oscillate faster than their fastest Fourier component. Physical phenomena associated with superoscillatory functions are known since long time and in more recent years there has been a wide interest both from the physical and the mathematical point of view. These functions (or sequences) appeared in weak values in quantum mechanics, see [2, 11, 22], in antenna theory this phenomenon was formulated in [34]. The literature on superoscillations is large, and without claiming completeness we mention the papers [13], [18]-[21], [28]-[30] and [32]. This class of functions has been investigated also from the mathematical point of view, as function theory, but large part of the results are associated with the study of the evolution of superoscillations by quantum fields equations with particular attention to Schrödiger equation. We give a quite complete list of papers [1], [3]-[9], [14]-[16], [23]-[27], [33] where one can find an up-to-date panorama of this field. In order to have an overview of the techniques developed in the recent years to study the evolution of superoscillations and their function theory, we refer the reader to the introductory papers

^{*}Schmid College of Science and Technology, Chapman University, Orange 92866, CA, US, aharonov@chapman.edu, tollakse@chapman.edu

[†]Politecnico di Milano, Dipartimento di Matematica, Via E. Bonardi, 9 20133 Milano, Italy, fabrizio.colombo@polimi.it, irene.sabadini@polimi.it

[‡]Department of Business Administration, Guilford Glazer Faculty of Business and Management, Ben-Gurion University of the Negev, Beer-Sheva, Israel, tomershu@bgu.ac.il

[§]The Donald Bren Distinguished Presidential Chair in Mathematics, Chapman University, Orange, USA, struppa@chapman.edu

[10, 12, 14] and [31]. Finally we mention the *Roadmap on superoscillations*, see [17], where the most recent advances in superoscillations and their applications to technology are well explained by the leading experts in this field.

A fundamental problem is to determine how large is the class of superoscillatory functions. The prototypical superoscillating function that is the outcome of weak values is given by

$$F_n(x,a) = \left(\cos\left(\frac{x}{n}\right) + ia\sin\left(\frac{x}{n}\right)\right)^n = \sum_{j=0}^n C_j(n,a)e^{i(1-2j/n)x}, \quad x \in \mathbb{R},$$
(1)

where a > 1 and the coefficients $C_i(n, a)$ are given by

$$C_j(n,a) = \binom{n}{j} \left(\frac{1+a}{2}\right)^{n-j} \left(\frac{1-a}{2}\right)^j.$$
(2)

If we fix $x \in \mathbb{R}$ and we let n go to infinity, we obtain that

$$\lim_{n \to \infty} F_n(x, a) = e^{iax}.$$

Clearly the name superoscillations comes from the fact that in the Fourier's representation of the function (1) the frequencies 1 - 2j/n are bounded by 1, but the limit function e^{iax} has a frequency *a* that can be arbitrarily larger than 1. A precise definition of superoscillating functions is as follows.

We call generalized Fourier sequence a sequence of the form

$$f_n(x) := \sum_{j=0}^n X_j(n,a) e^{ih_j(n)x}, \quad j = 0, ..., n, \quad n \in \mathbb{N},$$
(3)

where $a \in \mathbb{R}$, $X_j(n, a)$ and $h_j(n)$ are complex and real valued functions of the variables n, a and n, respectively. A generalized Fourier sequence of the form (3) is said to be a superoscillating sequence if $\sup_{j,n} |k_j(n)| \leq 1$ and there exists a compact subset of \mathbb{R} , which will be called a superoscillation set, on which $f_n(x)$ converges uniformly to $e^{ig(a)x}$, where g is a continuous real valued function such that |g(a)| > 1.

The classical Fourier expansion is obviously not a superoscillating sequence since its frequencies are not, in general, bounded. Using infinite order differential operators we can define a class of superoscillatory function applying them to the functions (1) and we obtain superoscillating functions of the form

$$Y_n(x,a) = \sum_{j=0}^n C_j(n,a) e^{ig(1-2j/n)x},$$
(4)

where $C_j(n, a)$ are the coefficients in (2), g are given entire functions, monotone increasing in a, and $x \in \mathbb{R}$. We have shown that

$$\lim_{n \to \infty} Y_n(x, a) = e^{ig(a)x}$$

under suitable conditions on g and the simplest, but important, example is

$$Y_n(x,a) = \sum_{j=0}^n C_j(n,a) e^{i(1-2j/n)^m x}, \text{ for fixed } m \in \mathbb{N}.$$

From the above considerations we deduce that there exists a large class of superoscillating functions taking different functions g.

In this paper we further enlarge the class of superoscillating functions enlarging both the class of the coefficients $C_j(n, a)$ and of the sequence of frequencies 1 - 2j/n that are bounded by 1. A large class of superoscillating functions can be determined solving the following problem.

Problem 1.1. Let $h_j(n)$ be a given set of points in [-1,1], j = 0, 1, ..., n, for $n \in \mathbb{N}$ and let $a \in \mathbb{R}$ be such that |a| > 1. Determine the coefficients $X_j(n)$ of the sequence

$$f_n(x) = \sum_{j=0}^n X_j(n) e^{ih_j(n)x}, \quad x \in \mathbb{R}$$

in such a way that

$$f_n^{(p)}(0) = (ia)^p$$
, for $p = 0, 1, ..., n$.

Remark 1.2. The conditions $f_n^{(p)}(0) = (ia)^p$ mean that the functions $x \mapsto e^{iax}$ and $f_n(x)$ have the same derivatives at the origin, for p = 0, 1, ..., n, so they have the same Taylor polynomial of order n.

Under the condition that the points $h_j(n)$ for j = 0, ..., n, (often denoted by h_j) are distinct we obtain an explicit formula for the coefficients $X_j(n, a)$ given by

$$X_{j}(n,a) = \prod_{k=0, \ k \neq j}^{n} \Big(\frac{h_{k}(n) - a}{h_{k}(n) - h_{j}(n)} \Big),$$

so the superoscillating sequence $f_n(x)$, that solves Problem 1.1, takes the explicit form

$$f_n(x) = \sum_{j=0}^n \prod_{k=0, \ k \neq j}^n \left(\frac{h_k(n) - a}{h_k(n) - h_j(n)} \right) \ e^{ih_j(n)x}, \quad x \in \mathbb{R},$$

as shown in Theorem 2.2. Observe that, by construction, this function is band limited and it converges to e^{iax} with arbitrary |a| > 1, so it is superoscillating.

Observe that different sequences $X_j(n)$ can be explicitly computed when we fix the points $h_j(n)$. See, for example, the case of the sequence

$$h_j(n) = 1 - 2j/n^p$$
 for $j = 0, ..., n, n \in \mathbb{N}$ and for fixed $p \in \mathbb{N}$,

in Section 3.

We consider now the frequencies $h_j(n) = (1 - 2j/n)^m$, for fixed $m \in \mathbb{N}$, to explain some facts.

(I) If we consider the sequence (4), with coefficients $C_j(n,a)$ given by (2), we obtain

$$\lim_{n \to \infty} \sum_{j=0}^{n} C_j(n,a) e^{i(1-2j/n)^m x} = e^{ia^m x}, \quad \text{ for fixed } m \in \mathbb{N}.$$

Note that, in this case, we could have used the frequencies $h_j(n) = (1-2j/n)$ and the coefficients $\tilde{C}_j(n,a) := C_j(n,a^m)$ to get as limit function $e^{ia^m x}$. Thus the same limit function $e^{ia^m x}$ can be obtained by tuning the frequencies and the coefficients.

(II) By solving Problem 1.1 with the frequences $h_j(n) = (1 - 2j/n)^m$, we can determine the coefficients $X_j(n) = X_j(n, a)$ such that we obtain as limit function e^{iax} , namely

$$\lim_{n \to \infty} \sum_{j=0}^{n} X_j(n,a) e^{i(1-2j/n)^m x} = e^{iax}, \quad \text{for fixed} \quad m \in \mathbb{N}.$$
 (5)

Changing the coefficients $\tilde{X}_j(n,a)$ we can get, as limit function, $e^{ia^m x}$.

(III) The coefficients X_j and C_j in the procedures (I) and (II) are different form each other because the two methods to generate superoscillations are different as explained in Section 3.

In section 4 we will also discuss how to generalize this method to obtain analogous results in the case of the supershift property of functions, a mathematical concept that generalizes the notion of superoscillating function.

2 A new class of superoscillating functions

In this section we show the main procedure to determine the coefficients $X_j(n)$ and so to construct explicitly the superoscillating functions solving Problem 1.1.

Theorem 2.1 (Existence and uniqueness of the solution of Problem 1.1). Let $h_j(n)$ be a given set of points in [-1,1], j = 0, 1, ..., n for $n \in \mathbb{N}$ and let $a \in \mathbb{R}$ be such that |a| > 1. If $h_j(n) \neq h_i(n)$, for every $i \neq j$, then there exists a unique solution $X_j(n)$ of the linear system

$$f_n^{(p)}(0) = (ia)^p$$
, for $p = 0, 1, ..., n$,

in Problem 1.1.

Proof. For the sake of simplicity, we denote $h_j(n)$ by h_j . Observe that the derivatives of order p of $f_n(x)$ are

$$f_n^{(p)}(x) = \sum_{j=0}^n X_j(n)(ih_j)^p e^{ih_j x}, \quad x \in \mathbb{R},$$

so if we require that these derivatives are equal to the derivatives of order p for p = 0, 1, ..., n of the function $x \mapsto e^{iax}$ at the origin we obtain the linear system

$$\sum_{j=0}^{n} X_j(n)(ih_j)^p = (ia)^p, \quad p = 0, 1, ..., n$$
(6)

from which we deduce

$$\sum_{j=0}^{n} X_j \ (h_j)^p = a^p, \quad p = 0, 1, ..., n,$$
(7)

where we have written X_j instead of $X_j(n)$. Now we write explicitly the linear system (7) of (n+1) equations and (n+1) unknowns $(X_0, ..., X_n)$

$$X_0 + X_1 + \dots + X_n = 1$$
$$X_0 h_0 + X_1 h_1 + \dots + X_n h_n = a$$
$$\dots$$
$$X_0 h_0^n + X_1 h_1^n + \dots + X_n h_n^n = a^n$$

and, in matrix form, it becomes

$$H(n)X = B(a) \tag{8}$$

where H is the $(n+1) \times (n+1)$ matrix

$$H(n) = \begin{pmatrix} 1 & 1 & \dots & 1 \\ h_0 & h_1 & \dots & h_n \\ \dots & \dots & \dots & \dots \\ h_0^n & h_1^n & \dots & h_n^n \end{pmatrix}$$
(9)

and

$$X = \begin{pmatrix} X_0 \\ X_1 \\ \dots \\ X_n \end{pmatrix} \quad \text{and} \quad B(a) = \begin{pmatrix} 1 \\ a \\ \dots \\ a^n \end{pmatrix}.$$
(10)

Observe that the determinant of H is the Vandermonde determinant, so it is given by

$$\det(H(n)) = \prod_{0 \le i < j \le n} (h_j(n) - h_i(n)).$$

Thus, if $h_j(n) \neq h_i(n)$ for every $i \neq j$, there exists a unique solution of the system.

Theorem 2.2 (Explicit solution of Problem 1.1). Let $h_j(n)$ be a given set of points in [-1, 1], j = 0, 1, ..., n for $n \in \mathbb{N}$ and let $a \in \mathbb{R}$ be such that |a| > 1. If $h_j(n) \neq h_i(n)$, for every $i \neq j$, the unique solution of system (8) is given by

$$X_j(n,a) = \prod_{k=0, \ k \neq j}^n \Big(\frac{h_k(n) - a}{h_k(n) - h_j(n)} \Big).$$
(11)

As a consequence, the superoscillating function takes the form

$$f_n(x) = \sum_{j=0}^n \prod_{k=0, \ k \neq j}^n \left(\frac{h_k(n) - a}{h_k(n) - h_j(n)} \right) \, e^{ih_j x}, \quad x \in \mathbb{R}.$$

Proof. In Theorem 2.1 we proved that, if $h_j \neq h_i$ for every $i \neq j$, there exists a unique solution of the system (8). The solution is given by

$$X_j(n,a) = \frac{\det(H_j(n,a))}{\det(H(n))}$$
(12)

for

$$H_{j}(n,a) = \begin{pmatrix} 1 & 1 & \dots & 1 & \dots & 1 \\ h_{0} & h_{1} & \dots & a & \dots & h_{n} \\ \dots & \dots & \dots & \dots & \dots & \dots \\ h_{0}^{n} & h_{1}^{n} & \dots & a^{n} & \dots & h_{n}^{n} \end{pmatrix}$$
(13)

where the j^{th} -column contains a and its powers. The explicit form of the determinant of the matrix H is given by:

$$det(H(n)) = (h_1 - h_0) \cdot (h_2 - h_0)(h_2 - h_1) \cdot (h_3 - h_0)(h_3 - h_1)(h_3 - h_2)$$
$$\cdot (h_4 - h_0)(h_4 - h_1)(h_4 - h_2)(h_4 - h_3) \cdot \ldots \cdot (h_n - h_0)(h_n - h_1)(h_n - h_2)(h_n - h_3)....(h_n - h_{n-1}).$$

The matrix $H_j(n, a)$ is still of Vandermonde type and its determinant can be computed similarly. So we have that the solution $(X_0(n, a), \ldots, X_n(n, a))$ is such that

$$\begin{aligned} X_0(n,a) &= \frac{(h_1 - a) \cdot (h_2 - a) \cdot (h_3 - a) \cdot (h_4 - a) \cdot \dots \cdot (h_n - a)}{(h_1 - h_0) \cdot (h_2 - h_0) \cdot (h_3 - h_0) \cdot (h_4 - h_0) \cdot \dots \cdot (h_n - h_0)} \\ &= \frac{\prod_{k=0, \ k \neq 0}^n (h_k - a)}{\prod_{k=0, \ k \neq 0}^n (h_k - h_0)}, \\ X_1(n,a) &= \frac{(a - h_0) \cdot (h_2 - a) \cdot (h_3 - a) \cdot (h_4 - a) \cdot \dots \cdot (h_n - a)}{(h_1 - h_0) \cdot (h_2 - h_1) \cdot (h_3 - h_1) \cdot (h_4 - h_1) \cdot \dots \cdot (h_n - h_1)} \\ &= \frac{\prod_{k=0, \ k \neq 1}^n (h_k - a)}{\prod_{k=0, \ k \neq 1}^n (h_k - h_1)}, \end{aligned}$$

and so on, up to

$$X_n(n,a) = \frac{1 \cdot 1 \cdot 1 \cdot 1 \cdot \dots \cdot (a-h_0)(a-h_1)(a-h_2)(a-h_3)\dots(a-h_{n-1})}{1 \cdot 1 \cdot 1 \cdot 1 \cdot \dots \cdot (h_n-h_0)(h_n-h_1)(h_n-h_2)(h_n-h_3)\dots(h_n-h_{n-1})}$$
$$= \frac{\prod_{k=0, \ k \neq n}^n (h_k-a)}{\prod_{k=0, \ k \neq n}^n (h_k-h_n)}.$$

So we get the statement with the recursive formula.

3 The methods to generate superoscillations and examples

Below we compare the superoscillating functions obtained by solving Problem 1.1 and the superoscillating functions obtained via the sequence $F_n(x, a)$ and infinite order differential operators. For different methods see also [30].

(I) Observe that the limit

$$\lim_{n \to \infty} \left(\cos\left(\frac{x}{n}\right) + ia\sin\left(\frac{x}{n}\right) \right)^n = e^{iax}$$

is a direct consequence of

$$\lim_{n \to \infty} \left(1 + ia \frac{x}{n} \right)^n = e^{iax},$$

while the construction method to generate superoscillations in Theorem 2.2 has a different nature because we require that the linear system (6) in the n + 1 unknowns $X_j(n)$ are determined in such a way that

$$\sum_{j=0}^{n} X_j(n)(ih_j)^p = (ia)^p, \quad p = 0, 1, ..., n$$
(14)

so the derivatives

$$f_n^{(p)}(x) = \sum_{j=0}^n X_j(n)(ih_j)^p e^{ih_j(n)x}, \quad x \in \mathbb{R},$$

at x = 0, are equal to the derivatives of the exponential function e^{iax} at the origin. This means that the sequence of functions

$$f_n(x) = \sum_{j=0}^n X_j(n) e^{ih_j(n)x}, \quad x \in \mathbb{R}$$

has n derivatives equal to the derivatives of exponential function e^{iax} at the origin so the limit

$$\lim_{n \to \infty} f_n(x) = e^{iax}$$

follows by construction of the $f_n(x)$.

(II) In the definition of the superoscillating function (1) the derivatives are given by

$$F_n^{(p)}(x,a) = \sum_{j=0}^n C_j(n,a) \left(i(1-2j/n) \right)^p e^{i(1-2j/n)x}, \quad x \in \mathbb{R}$$

and it is just in the limit that we get the derivatives of order p of the exponential function e^{iax} at the origin, namely we have

$$\lim_{n \to \infty} \sum_{j=0}^{n} C_j(n,a) \left(i(1-2j/n) \right)^p = (ia)^p, \quad p \in \mathbb{N}.$$

(III) With the new procedure proposed in this paper we impose the conditions

$$f_n^{(p)}(0) = (ia)^p, \quad p = 0, 1, 2, ..., n$$

(where we have genuine equalities, not in the limit) and we link n with the derivatives of $f_n(x)$ in order to determine the coefficients $X_j(n)$ in (3), so we have that the Taylor polynomials of the two functions $f_n(x)$ and e^{iax} are the same up to order n, i.e.,

$$e^{iax} = 1 + iax + \frac{(iax)^2}{2!} + \dots + \frac{(iax)^n}{n!} + R_n(x),$$

so we get

$$f_n(x) = \sum_{j=0}^n X_j(n)e^{ih_j x}$$

= $f_n(0) + f_n^{(1)}(0)x + f_n^{(2)}(0)\frac{x^2}{2!} + \dots + f_n^{(n)}(0)\frac{x^n}{n!} + \tilde{R}_n(x), \quad x \in \mathbb{R}$
= $1 + iax + (ia)^2\frac{x^2}{2!} + \dots + (ia)^n\frac{x^n}{n!} + \tilde{R}_n(x), \quad x \in \mathbb{R},$

where $R_n(x)$ and $\tilde{R}_n(x)$ are the errors.

It is now easy to generate a very large class of superoscillatory functions. We write a few examples to further clarify the generality of our new construction of superoscillating sequences: given the sequence $h_j(n)$ we determine the coefficients accordingly for the

$$f_n(x) = \sum_{j=0}^n \prod_{k=0, \ k \neq j}^n \left(\frac{h_k(n) - a}{h_k(n) - h_j(n)} \right) e^{ih_j(n)x}, \quad x \in \mathbb{R}.$$

Example 3.1. Let $n \in \mathbb{N}$ and set

$$h_j(n) = 1 - \frac{2}{n}j$$

where j = 0, ..., n. We have

$$h_k(n) - a = 1 - \frac{2}{n}k - a$$

and

$$h_k(n) - h_j(n) = 1 - \frac{2}{n}k - \left(1 - \frac{2}{n}j\right) = \frac{2}{n}\left(j - k\right).$$

Thus, we obtain

$$f_n(x) = \sum_{j=0}^n \prod_{k=0, \ k \neq j}^n \frac{n}{2} \left(\frac{1 - \frac{2}{n}k - a}{j - k} \right) e^{i(1 - \frac{2}{n}j)x}, \quad x \in \mathbb{R}.$$

Example 3.2. Let $n \in \mathbb{N}$, and set

$$h_j(n) = 1 - \frac{2}{n^p}j$$

where j = 0, ..., n, for a fixed $p \in \mathbb{N}$. We have

$$h_k(n) - a = 1 - \frac{2}{n^p}k - a_k$$

and

$$h_k(n) - h_j(n) = 1 - \frac{2}{n^p}k - \left(1 - \frac{2}{n^p}j\right) = \frac{2}{n^p}(j-k).$$

So, we obtain:

$$f_n(x) = \sum_{j=0}^n \prod_{k=0, \ k \neq j}^n \frac{n^p}{2} \left(\frac{1 - \frac{2}{n^p}k - a}{j - k} \right) \ e^{i(1 - \frac{2}{n^p}j)x}, \quad x \in \mathbb{R}.$$

Example 3.3. Let $n \in \mathbb{N}$, and set

$$h_j(n) = 1 - \left(\frac{2j}{n}\right)^p$$

where j = 0, ..., n, for a fixed $p \in \mathbb{N}$. We have

$$h_k(n) - a = 1 - \left(\frac{2k}{n}\right)^p - a,$$

and

$$h_k(n) - h_j(n) = 1 - \left(\frac{2k}{n}\right)^p - \left(1 - \left(\frac{2j}{n}\right)^p\right) = \frac{2^p}{n^p} \left(j^p - k^p\right).$$

So, we obtain

$$f_n(x) = \sum_{j=0}^n \prod_{k=0, \ k \neq j}^n \frac{n^p}{2^p} \left(\frac{1 - \frac{2k}{n} - a}{j^p - k^p}\right) \ e^{i(1 - (2j/n)^p)x}, \quad x \in \mathbb{R}.$$

4 A new class of supershifts

The procedure to define superoscillatory functions can be extended to supershift. We recall that the supershift property of a function extends the notion of superoscillations and it turned out to be the crucial concept behind the study of the evolution of superoscillatory functions as initial conditions of Schrödinger equation or of any other field equation. We recall the definition before to state our result. **Definition 4.1** (Supershift). Let $\mathcal{I} \subseteq \mathbb{R}$ be an interval with $[-1,1] \subset \mathcal{I}$ and let $\varphi : \mathcal{I} \times \mathbb{R} \to \mathbb{R}$, be a continuous function on \mathcal{I} . We set

$$\varphi_{\lambda}(x) := \varphi(\lambda, x), \quad \lambda \in \mathcal{I}, \quad x \in \mathbb{R}$$

and we consider a sequence of points $(\lambda_{j,n})$ such that

$$(\lambda_{j,n}) \in [-1,1]$$
 for $j = 0, ..., n$ and $n = 0, ..., +\infty$.

We define the functions

$$\psi_n(x) = \sum_{j=0}^n c_j(n)\varphi_{\lambda_{j,n}}(x),\tag{15}$$

where $(c_j(n))$ is a sequence of complex numbers for j = 0, ..., n and $n = 0, ..., +\infty$. If $\lim_{n\to\infty} \psi_n(x) = \varphi_a(x)$ for some $a \in \mathcal{I}$ with |a| > 1, we say that the function $\lambda \to \varphi_\lambda(x)$, for x fixed, admits a supershift in λ .

Remark 4.2. We observe that the definition of supershift of a function given above is not the most general one, but it is useful to explain our new procedure for the supershift case. In the following, we will take the interval \mathcal{I} , in the definition of the supershift, to be equal to \mathbb{R} .

Remark 4.3. Let us stress that the term supershift comes from the fact that the interval \mathcal{I} can be arbitrarily large (it can also be \mathbb{R}) and so also the constant a can be arbitrarily far away from the interval [-1,1] where the function $\varphi_{\lambda_{i,n}}(\cdot)$ is computed, see (15).

Remark 4.4. Superoscillations are a particular case of supershift. In fact, for the sequence (F_n) in (1), we have $\lambda_{j,n} = 1 - 2j/n$, $\varphi_{\lambda_{j,n}}(t,x) = e^{i(1-2j/n)x}$ and $c_j(n)$ are the coefficients $C_j(n,a)$ defined in (2).

Problem 1.1, for the supershift case, is formulated as follows.

Problem 4.5. Let $h_j(n)$ be a given set of points in [-1,1], j = 0, 1, ..., n, for $n \in \mathbb{N}$ and let $a \in \mathbb{R}$ be such that |a| > 1. Suppose that for every $x \in \mathbb{R}$ the function $\lambda \mapsto G(\lambda x)$ is holomorphic and entire in λ . Consider the function

$$f_n(x) = \sum_{j=0}^n Y_j(n) G(h_j(n)x), \quad x \in \mathbb{R}$$

where $\lambda \mapsto G(\lambda x)$ depends on the parameter $x \in \mathbb{R}$. Determine the coefficients $Y_j(n)$ in such a way that

$$f_n^{(p)}(0) = (a)^p G^{(p)}(0) \quad for \quad p = 0, 1, ..., n.$$
 (16)

The solution of the above problem is obtained in the following theorem.

Theorem 4.6. Let $h_j(n)$ be a given set of points in [-1,1], j = 0, 1, ..., n for $n \in \mathbb{N}$ and let $a \in \mathbb{R}$ be such that |a| > 1. If $h_j(n) \neq h_i(n)$ for every $i \neq j$ and $G^{(p)}(0) \neq 0$ for all p = 0, 1, ..., n, then there exists a unique solution $Y_j(n, a)$ of the linear system (16) and it is given by

$$Y_{j}(n,a) = \prod_{k=0, \ k \neq j}^{n} \Big(\frac{h_{k}(n) - a}{h_{k}(n) - h_{j}(n)} \Big),$$

so that

$$f_n(x) = \sum_{j=0}^n \prod_{k=0, \ k \neq j}^n \left(\frac{h_k(n) - a}{h_k(n) - h_j(n)} \right) G(h_j(n)x), \quad x \in \mathbb{R}$$

and, by construction, it is

$$\lim_{n \to \infty} f_n(x) = G(ax), \quad x \in \mathbb{R}.$$

Proof. Observe that we have

$$f_n^{(p)}(x) = \sum_{j=0}^n Y_j(n)(h_j(n))^p G^{(p)}(h_j(n)x), \quad x \in \mathbb{R}$$

where $G^{(p)}$ are the derivatives of order p, for p = 0, ..., n with respect to x of the function $G(\lambda x)$, for $\lambda \in \mathbb{R}$ considered as a parameter. So we get the system

$$f_n^{(p)}(0) = \sum_{j=0}^n Y_j(n)(h_j(n))^p G^{(p)}(0) = a^p G^{(p)}(0).$$

Now, since we have assumed that $G^{(p)}(0) \neq 0$ for all p = 0, 1, ..., n, the system becomes

$$\sum_{j=0}^{n} Y_j(n)(h_j(n))^p = a^p$$

and we can solve it as in Theorem 2.2 to get

$$Y_j(n,a) = \prod_{k=0, \ k \neq j}^n \Big(\frac{h_k(n) - a}{h_k(n) - h_j(n)} \Big).$$

Finally, we get

$$f_n(x) = \sum_{j=0}^n \prod_{k=0, \ k \neq j}^n \left(\frac{h_k(n) - a}{h_k(n) - h_j(n)} \right) G(h_j(n)x), \quad x \in \mathbb{R}$$

and by construction it is

$$\lim_{n \to \infty} f_n(x) = G(ax), \quad x \in \mathbb{R}.$$

References

- D. Alpay, F. Colombo, I. Sabadini, D.C. Struppa, Aharonov-Berry superoscillations in the radial harmonic oscillator potential, Quantum Stud. Math. Found., 7 (2020), 269–283.
- [2] Y. Aharonov, D. Albert, L. Vaidman, How the result of a measurement of a component of the spin of a spin-1/2 particle can turn out to be 100, Phys. Rev. Lett., 60 (1988), 1351-1354.
- [3] Y. Aharonov, J. Behrndt, F. Colombo, P. Schlosser, Schrödinger evolution of superoscillations with δ- and δ'-potentials, Quantum Stud. Math. Found., 7 (2020), 293–305.

- [4] Y. Aharonov, J. Behrndt, F. Colombo, P. Schlosser, Green's Function for the Schrödinger Equation with a Generalized Point Interaction and Stability of Superoscillations, J. Differential Equations, 277 (2021), 153–190.
- [5] Y. Aharonov, F. Colombo, I. Sabadini, D.C. Struppa, J. Tollaksen, Evolution of superoscillations in the Klein–Gordon field, Milan J. Math., 88 (2020), no. 1, 171–189.
- [6] Y. Aharonov, F. Colombo, I. Sabadini, D.C. Struppa, J. Tollaksen, How superoscillating tunneling waves can overcome the step potential, Ann. Physics, 414 (2020), 168088, 19 pp.
- [7] Y. Aharonov, F. Colombo, I. Sabadini, D.C. Struppa, J. Tollaksen, On the Cauchy problem for the Schrödinger equation with superoscillatory initial data, J. Math. Pures Appl., 99 (2013), 165–173.
- [8] Y. Aharonov, F. Colombo, I. Sabadini, D.C. Struppa, J. Tollaksen, Superoscillating sequences in several variables, J. Fourier Anal. Appl., 22 (2016), 751–767.
- [9] Y. Aharonov, F. Colombo, I. Sabadini, D.C. Struppa, J. Tollaksen, The mathematics of superoscillations, Mem. Amer. Math. Soc., 247 (2017), no. 1174, v+107 pp.
- [10] Y. Aharonov, F. Colombo, D.C. Struppa, J. Tollaksen, Schrödinger evolution of superoscillations under different potentials, Quantum Stud. Math. Found., 5 (2018), 485–504.
- [11] Y. Aharonov, D. Rohrlich, Quantum Paradoxes: Quantum Theory for the Perplexed, Wiley-VCH Verlag, Weinheim, 2005.
- [12] Y. Aharonov, I. Sabadini, J. Tollaksen, A. Yger, *Classes of superoscillating functions*, Quantum Stud. Math. Found., 5 (2018), 439–454.
- [13] Y. Aharonov, T. Shushi, A new class of superoscillatory functions based on a generalized polar coordinate system, Quantum Stud. Math. Found., 7 (2020), 307–313.
- [14] T. Aoki, F. Colombo, I. Sabadini, D. C. Struppa, Continuity of some operators arising in the theory of superoscillations, Quantum Stud. Math. Found., 5 (2018), 463–476.
- [15] T. Aoki, F. Colombo, I. Sabadini, D.C. Struppa, Continuity theorems for a class of convolution operators and applications to superoscillations, Ann. Mat. Pura Appl., 197 (2018), 1533–1545.
- [16] J. Behrndt, F. Colombo, P. Schlosser, Evolution of Aharonov-Berry superoscillations in Dirac δ-potential, Quantum Stud. Math. Found., 6 (2019), 279–293.
- [17] M. Berry et al, *Roadmap on superoscillations*, 2019, Journal of Optics 21 053002.
- [18] M. V. Berry, Faster than Fourier, in Quantum Coherence and Reality; in celebration of the 60th Birthday of Yakir Aharonov ed. J.S.Anandan and J. L. Safko, World Scientific, Singapore, (1994), pp. 55-65.
- M. Berry, Exact nonparaxial transmission of subwavelength detail using superoscillations, J. Phys. A 46, (2013), 205203.
- [20] M. V. Berry, Representing superoscillations and narrow Gaussians with elementary functions, Milan J. Math., 84 (2016), 217–230.

- [21] M. V. Berry, S. Popescu, Evolution of quantum superoscillations, and optical superresolution without evanescent waves, J. Phys. A, 39 (2006), 6965–6977.
- [22] M.V. Berry, P. Shukla, Pointer supershifts and superoscillations in weak measurements, J. Phys A, 45 (2012), 015301.
- [23] F. Colombo, J. Gantner, D.C. Struppa, Evolution by Schrödinger equation of Aharonov-Berry superoscillations in centrifugal potential, Proc. A., 475 (2019), no. 2225, 20180390, 17 pp.
- [24] F. Colombo, I. Sabadini, D.C. Struppa, A. Yger, Gauss sums, superoscillations and the Talbot carpet, Journal de Mathématiques Pures et Appliquées, https://doi.org/10.1016/j.matpur.2020.07.011.
- [25] F. Colombo, I. Sabadini, D.C. Struppa, A. Yger, Superoscillating sequences and hyperfunctions, Publ. Res. Inst. Math. Sci., 55 (2019), no. 4, 665–688.
- [26] F. Colombo, D.C. Struppa, A. Yger, Superoscillating sequences towards approximation in S or S'-type spaces and extrapolation, J. Fourier Anal. Appl., 25 (2019), no. 1, 242–266.
- [27] F. Colombo, G. Valente, Evolution of Superoscillations in the Dirac Field, Found. Phys., 50 (2020), 1356–1375.
- [28] P. J. S. G. Ferreira, A. Kempf, Unusual properties of superoscillating particles, J. Phys. A, 37 (2004), 12067-76.
- [29] P. J. S. G. Ferreira, A. Kempf, Superoscillations: faster than the Nyquist rate, IEEE Trans. Signal Processing, 54 (2006), 3732–3740.
- [30] P. J. S. G. Ferreira, A. Kempf, M. J. C. S. Reis, Construction of Aharonov-Berry's superoscillations, J. Phys. A 40 (2007), 5141–5147.
- [31] A. Kempf, Four aspects of superoscillations, Quantum Stud. Math. Found., 5 (2018), 477– 484.
- [32] J. Lindberg, Mathematical concepts of optical superresolution, Journal of Optics, 14 (2012), 083001.
- [33] B. Soda, A. Kempf, Efficient method to create superoscillations with generic target behavior, Quantum Stud. Math. Found., 7 (2020), no. 3, 347–353.
- [34] G. Toraldo di Francia, Super-Gain Antennas and Optical Resolving Power, Nuovo Cimento Suppl., 9 (1952), 426–438.