

4-15-2024

Generalized q-Fock Spaces and Structural Identities

Daniel Alpay

Chapman University, alpay@chapman.edu

Paula Cerejeiras

University of Aveiro

Uwe Kaehler

University of Aveiro

Baruch Schneider

University of Ostrava

Follow this and additional works at: https://digitalcommons.chapman.edu/scs_articles



Part of the [Other Mathematics Commons](#)

Recommended Citation

D. Alpay, P. Cerejeiras, U. Kaehler, and B. Schneider. Generalized q-Fock spaces and structural identities. *Proceedings of the American Mathematical Society*, 2024. <https://doi.org/10.1090/proc/16720>

This Article is brought to you for free and open access by the Science and Technology Faculty Articles and Research at Chapman University Digital Commons. It has been accepted for inclusion in Mathematics, Physics, and Computer Science Faculty Articles and Research by an authorized administrator of Chapman University Digital Commons. For more information, please contact laughtin@chapman.edu.

Generalized q-Fock Spaces and Structural Identities

Comments

This is a pre-copy-editing, author-produced PDF of an article that later underwent peer review and was accepted for publication in *Proceedings of the American Mathematical Society* in 2024. This article may not exactly replicate the final published version. The definitive publisher-authenticated version is available online at <https://doi.org/10.1090/proc/16720>.

Copyright

American Mathematical Society

GENERALIZED q -FOCK SPACES AND A NEW TYPE OF STIRLING NUMBERS

DANIEL ALPAY, PAULA CEREJEIRAS, UWE KAEHLER, AND BARUCH SCHNEIDER

ABSTRACT. Using q -calculus we study a family of reproducing kernel Hilbert spaces which interpolate between the Hardy space and the Fock space. We give characterizations of these spaces in terms of classical operators such as integration and backward-shift, and their q -calculus counterparts. We introduce an apparently new family of numbers, close to, but different from, the q -Stirling numbers of the second kind.

1. INTRODUCTION

1.1. Prologue. The Hardy space of the open unit disk \mathbb{D} , here denoted by $\mathbf{H}_2 = \mathbf{H}_2(\mathbb{D})$, is the reproducing kernel Hilbert space with reproducing kernel

$$\frac{1}{1 - z\bar{w}} = \sum_{n=0}^{\infty} z^n \bar{w}^n, \quad z, w \in \mathbb{D},$$

and plays a key role in operator theory, linear system theory and Schur analysis. On the other hand, the Bargmann-Fock-Segal space, here denoted by \mathcal{F} and called Fock space for short, is the reproducing kernel Hilbert space with reproducing kernel

$$e^{z\bar{w}} = \sum_{n=0}^{\infty} \frac{z^n \bar{w}^n}{n!}, \quad z, w \in \mathbb{C},$$

and plays a key role in quantum mechanics (and more recently in signal processing).

The Hardy space \mathbf{H}_2 can be characterized (up to a positive multiplicative factor for the inner product) as the only Hilbert space of power series converging at the origin and such that

$$(1.1) \quad R_0^* = M_z,$$

where M_z is the operator of multiplication by z and

$$(1.2) \quad R_0 f(z) = \frac{f(z) - f(0)}{z}.$$

Note that in \mathbf{H}_2 we have the identities

$$(1.3) \quad R_0 R_0^* = \mathcal{I}, \quad \text{and} \quad R_0 M_z - M_z R_0 = \mathcal{I} - R_0^* R_0 = C^* C,$$

where $Cf = f(0)$ and \mathcal{I} is the identity operator. We remark that

$$(1.4) \quad \mathcal{I} - R_0^* R_0 = C^* C,$$

2010 *Mathematics Subject Classification.* 30H20; 26A33.

Key words and phrases. Fock space; fractional derivative; q -calculus.

D. Alpay thanks the Foster G. and Mary McGaw Professorship in Mathematical Sciences, which supported his research.

which we will call *structural identity*, is the simplest of a family of identities characterizing de Branges spaces.

Similarly, and besides Bargmann celebrated characterization $\partial^* = M_z$ (see [10, 11]), the Fock space is (still up to a positive multiplicative factor for the inner product) the only Hilbert space of power series converging at the origin and such that

$$(1.5) \quad R_0^* = I,$$

where I is the integration operator (see [7])

$$(1.6) \quad (If)(z) = \int_{[0,z]} f(s)ds.$$

1.2. The paper. The q -calculus allows to define a continuum of spaces between \mathbf{H}_2 and \mathcal{F} , namely the family of reproducing kernel Hilbert spaces $\mathbf{H}_{2,q}$ indexed by $q \in [0, 1]$ and with reproducing kernel

$$K_q(z, w) = \sum_{n=0}^{\infty} \frac{z^n \bar{w}^n}{[n]_q!}, \quad q \in [0, 1], \quad z, w \in \mathbb{D}_{1/1-q},$$

where in the above expression

$$\mathbb{D}_{1/1-q} = \begin{cases} \mathbb{D}_{\infty} = \mathbb{C}, & q = 1, \\ \left\{ z \in \mathbb{C} : |z| < \frac{1}{1-q} \right\}, & q \in [0, 1) \end{cases}.$$

Furthermore, $[0]_q! = 1$ and $[n]_q! = 1 \cdot (1+q) \cdot (1+q+q^2) \cdots (1+q+\cdots+q^{n-1})$, $n \in \mathbb{N}$. Thus, in this notation, we have

$$\mathbf{H}_{2,0} = \mathbf{H}_2 \quad \text{and} \quad \mathbf{H}_{2,1} = \mathcal{F},$$

with

$$K_0(z, w) := k_{2,0}(z, w) = \frac{1}{1 - z\bar{w}} \quad \text{and} \quad K_1(z, w) := k_{2,1}(z, w) = e^{z\bar{w}}.$$

The q -calculus allows to gather into a common umbrella problems pertaining to the classical Hardy space \mathbf{H}_2 of the open unit disk and problems pertaining to the Fock space. Consider now

$$(1.7) \quad R_q f(z) = \frac{f(z) - f(qz)}{(1-q)z}, \quad 0 \leq q < 1,$$

while for $q = 1$, we consider $R_1 = \partial$. In this way we have a progression between two fundamental linear operators in analysis, namely the backward-shift and the differentiation operators. Then, one can introduce the q -Fock space $\mathbf{H}_{2,q}$ as the unique (up to a multiplicative positive constant) space of power series such that $R_q^* = M_z$. The case $q = 1$ corresponds to the classical Fock space (see [10]). It is important to note already at this stage that these operators satisfy a q -commutator relation (see also Lemma 2.2)

$$(1.8) \quad R_q M_z - q M_z R_q = \mathcal{I}.$$

2. q -CALCULUS

2.1. Iterative powers of the operator R_q . Recall that R_q was defined by (1.7).

Proposition 2.1. *Let $\Lambda_q f(z) = f(qz)$. We have*

$$(2.9) \quad R_q^n f(z) = \frac{\prod_{k=1}^n (1 - q^k \Lambda_q)}{(1-q)^n} R_0^n f(z), \quad 0 \leq q < 1, \quad n = 1, 2, \dots$$

Proof. Firstly, we observe the intertwining between R_0 and Λ_q ,

$$R_0 \Lambda_q f(z) = R_0 f(qz) = \frac{f(qz) - f(0)}{z} = q \frac{f(qz) - f(0)}{qz} = q \Lambda_q R_0 f(z).$$

Secondly,

$$\begin{aligned} R_q f(z) &= \frac{f(z) - f(qz)}{(1-q)z} = \frac{f(z) - f(0) - f(qz) + f(0)}{(1-q)z} \\ &= \frac{1}{1-q} \frac{f(z) - f(0)}{z} - \frac{q}{1-q} \frac{f(qz) - f(0)}{qz} \\ &= \frac{(1 - q\Lambda_q)R_0}{1-q} f(z). \end{aligned}$$

Hence,

$$\begin{aligned} R_q^2 f(z) &= \left(\frac{(1 - q\Lambda_q)R_0}{1-q} \right)^2 f(z) \\ &= \frac{(1 - q\Lambda_q)R_0(1 - q\Lambda_q)R_0}{(1-q)^2} f(z) \\ &= \frac{(1 - q\Lambda_q)(1 - q^2\Lambda_q)R_0^2}{(1-q)^2} f(z), \end{aligned}$$

and by induction the result holds:

$$\begin{aligned} R_q^n f(z) &= \left(\frac{(1 - q\Lambda_q)R_0}{1-q} \right)^n f(z) \\ &= \frac{(1 - q\Lambda_q)R_0(1 - q\Lambda_q)R_0 \cdots (1 - q\Lambda_q)R_0}{(1-q)^n} f(z) \\ &= \frac{(1 - q\Lambda_q)(1 - q^2\Lambda_q)R_0^2 \cdots (1 - q\Lambda_q)R_0}{(1-q)^n} f(z) \\ &\vdots \\ &= \frac{(1 - q\Lambda_q)(1 - q^2\Lambda_q) \cdots (1 - q^n\Lambda_q)R_0^n}{(1-q)^n} f(z). \end{aligned}$$

□

As we will see later (see Theorems 4.4 and 4.5), R_q^* has completely different properties depending on which of the spaces at hand we compute the adjoint.

2.2. A new type of q -Stirling numbers.

Lemma 2.2 (q -commutator). *For the q -commutator it holds the following identity:*

$$(2.10) \quad [R_q, M_z]_q := R_q M_z - q M_z R_q = \mathcal{I}.$$

Proof. We have

$$R_q M_z z^n = R_q z^{n+1} = (1 + q + \cdots + q^n) z^n, \quad n = 0, 1, 2, \dots$$

while

$$\begin{aligned} qM_z R_q z^n &= qM_z R_q 1 = 0, \quad n = 0, \\ qM_z R_q z^n &= qM_z (1 + q + \cdots + q^{n-1}) z^{n-1} = (q + q + \cdots + q^n) z^n, \quad n = 1, 2, \dots \end{aligned}$$

so that it holds $(R_q M_z - qM_z R_q) z^n = z^n$, for all $n \in \mathbb{N}_0$. \square

We define our q -Stirling numbers as coefficients $S(n, k)$ of the following commutation relation (see [6]):

$$(2.11) \quad (M_z R_q)^n := \sum_{k=1}^n S(n, k) M_z^k R_q^k, \quad n \in \mathbb{N}.$$

Lemma 2.3. *We have for these q -Stirling numbers the following recursion formula*

$$\begin{aligned} S(1, 1) &= 1; \\ S(n, n) &= S(n-1, n-1) q^{n-1}, \quad n = 2, 3, \dots; \\ S(n, k) &= (1 + q + \cdots + q^{k-1}) S(n-1, k) + q^{k-1} S(n-1, k-1), \quad k = 2, \dots, n-1. \end{aligned}$$

Proof. In order to simplify notation, we write the expression for the q -Stirling numbers as

$$(ab)^n := \sum_{k=1}^n S(n, k) a^k b^k.$$

From the q -commutator we get $ba = 1 + qab$ so that

$$\begin{aligned} b^n a &= b^{n-1} (ba) = b^{n-1} (1 + qab) = b^{n-1} + q(b^{n-1} a) b \\ &= b^{n-1} + q [b^{n-2} + q(b^{n-2} a) b] b = (1 + q) b^{n-1} + q^2 (b^{n-2} a) b^2 \\ &\quad \vdots \\ &= (1 + q + \cdots + q^{n-1}) b^{n-1} + q^n a b^n. \end{aligned}$$

Replacing in the above formula for the q -Stirling numbers we obtain

$$\begin{aligned} (ab)^n &= \sum_{k=1}^n S(n, k) a^k b^k \\ &= (ab)^{n-1} (ab) = \left[\sum_{k=1}^{n-1} S(n-1, k) a^k b^k \right] (ab) \\ &= \sum_{k=1}^{n-1} S(n-1, k) a^k (b^k a) b = \sum_{k=1}^{n-1} a^k S(n-1, k) [(1 + q + \cdots + q^{k-1}) b^{k-1} + q^k a b^k] b \\ &= \sum_{k=1}^{n-1} [(1 + q + \cdots + q^{k-1}) S(n-1, k) a^k b^k + q^k S(n-1, k) a^{k+1} b^{k+1}] \\ &= \sum_{k=1}^{n-1} (1 + q + \cdots + q^{k-1}) S(n-1, k) a^k b^k + \sum_{k=2}^n q^{k-1} S(n-1, k-1) a^k b^k \\ &= S(n-1, 1) ab + \sum_{k=2}^{n-1} [(1 + q + \cdots + q^{k-1}) S(n-1, k) + q^{k-1} S(n-1, k-1)] a^k b^k + q^{n-1} S(n-1, n-1) a^n b^n, \end{aligned}$$

so that we have $S(1, 1) = 1$,

$$S(n, 1) = S(n-1, 1), \quad S(n, n) = q^{n-1} S(n-1, n-1),$$

for $n = 2, 3, \dots$ and

$$S(n, k) = (1 + q + \dots + q^{k-1})S(n-1, k) + q^{k-1}S(n-1, k-1),$$

for $k = 2, \dots, n-1$. □

One can easily see the first q -Stirling numbers

$S(n, k)$	1	2	3	4
1	1			
2	1	q		
3	1	$2q + q^2$	q^3	
4	1	$q^3 + 3q^2 + 3q$	$q^5 + 2q^4 + 3q^3$	q^6

Remark 2.4. We need to point out that the above q -Stirling numbers are not the q -Stirling numbers of the first or of the second kind already existent in the literature. These were obtained by studying the corresponding partition problems in q -calculus (for a review on this topic see [12, 13]). Here we obtain them as coefficients of the expansion of $(M_z R_q)^n$ in (2.11) following a similar idea as in [6]. Only in the classic case of $q = 1$ these type of coefficients coincide with the Stirling numbers of the second kind, i.e. with the numbers of partitions of a set of n objects into k non-empty subsets.

3. THE q -FOCK SPACE

Consider the positive definite function $E_q(z\overline{w})$ given by the q -exponential:

$$(3.1) \quad E_q(z) = \sum_{k=0}^{\infty} \frac{z^k}{[k]_q!} = \frac{1}{\prod_{j=0}^{\infty} (1 - z(1-q)q^j)} =: \frac{1}{(z(1-q); q)_{\infty}}, \quad z \in \mathbb{D}_{1/1-q},$$

evaluated at $z\overline{w}$, with $[0]_q = 1$ and $[k]_q = 1 + q + \dots + q^{k-1}$ for $k = 1, 2, \dots$, and $[k]_q! = \prod_{j=0}^k [j]_q$, i.e.

$$[k]_q! = [1]_q [2]_q \dots [k]_q = 1 \cdot (1+q) \cdot (1+q+q^2) \dots (1+q+\dots+q^{k-1}).$$

The term $(a; q)_n = \prod_{j=0}^{n-1} (1 - aq^j)$ denotes the q -Pochhammer symbol.

Definition 3.1. We denote by $\mathbf{H}_{2,q}$ the reproducing kernel Hilbert space of functions analytic in $|z| < \frac{1}{1-q}$ with reproducing kernel $E_q(z\overline{w})$.

As stated before when $q = 0$ we get back the classical Hardy space of the open unit disk, while $q \rightarrow 1$ leads to the classical Fock space; see e.g. [14, 15, 18] for the former, [20] for the latter.

For functions belonging to the q -Fock space we have the following characterization based on its power series expansion.

Lemma 3.2. $f(z) = \sum_{n=0}^{\infty} a_n z^n$ belongs to $\mathbf{H}_{2,q}$ if and only if

$$(3.2) \quad \sum_{n=0}^{\infty} [n]_q! |a_n|^2 < \infty.$$

Based on the q -Jackson integral (see [16], [17])

$$\int_0^a f(x) d_q x := (1-q)a \sum_{k=0}^{\infty} q^k f(q^k a),$$

we can define the following q -integral transform.

Definition 3.3. Given a bounded function $f : [0, -1+1/(1-q)] \rightarrow \mathbb{R}$ we define its q -integral transform as

$$\mathcal{M}_q f(z) = \int_0^{1/(1-q)} t^{z-1} f(qt) d_q t := \sum_{k=0}^{\infty} q^k \left(\frac{q^k}{1-q} \right)^{z-1} f\left(\frac{q^{k+1}}{1-q} \right).$$

With the help of this q -integral transform we get that the coefficients $\frac{1}{[n]_q!}$ satisfy the moment problem

$$\begin{aligned} [n]_q! = \mathcal{M}_q(E_q^{-1})(n+1) &= \int_0^{1/(1-q)} t^n E_q^{-1}(qt) d_q t \\ &= \frac{(q; q)_{\infty}}{(1-q)^n} \sum_{k=0}^{\infty} \frac{q^{(n+1)k}}{(q; q)_k}, \quad 0 < q < 1, \quad n \in \mathbb{N}_0, \end{aligned}$$

where $(a; q)_n = \prod_{j=0}^{n-1} (1 - aq^j)$ denotes the q -Pochhammer symbol and $(q; q)_n = \frac{(q; q)_{\infty}}{(q^{n+1}; q)_{\infty}}$. For the disk $\mathbb{D}_{1/(1-q)}$ we have the measure (see [19])

$$d\mu_q(z) = (q; q)_{\infty} \sum_{k=0}^{\infty} \frac{q^k}{(q; q)_k} d\lambda_{r_k}(z),$$

where $r_k = \frac{q^{k/2}}{\sqrt{1-q}}$ while $d\lambda_{r_k}$ is the normalized Lebesgue measure in the circle of radius r_k . This leads to the following characterization of the space $\mathbf{H}_{2,q}$.

Theorem 3.4. The space $\mathbf{H}_{2,q}$ corresponds to the space of all analytic functions in the disk $\mathbb{D}_{\frac{1}{1-q}} = \{z : |z| < \frac{1}{1-q}\}$ satisfying the condition

$$\iint_{\mathbb{D}_{\frac{1}{1-q}}} |f(z)|^2 d\mu_q(z) < \infty.$$

The inner product of $\mathbf{H}_{2,q}$ is given by

$$\frac{1}{2\pi} \iint_{\mathbb{D}_{\frac{1}{1-q}}} f(z) \overline{g(z)} d\mu_q(z) = \sum_{n=0}^{\infty} f_n \overline{g_n} [n]_q!.$$

Proof. We have

$$\begin{aligned} \frac{1}{2\pi} \iint_{\mathbb{D}_{\frac{1}{1-q}}} z^n \overline{z}^m d\mu_q(z) &= \frac{(q; q)_{\infty}}{2\pi} \sum_{k=0}^{\infty} \frac{q^k}{(q; q)_k} r_k^{n+m} \underbrace{\int_0^{2\pi} e^{i(n-m)\theta} d\theta}_{=2\pi\delta_{n,m}} \\ &= \delta_{n,m} (q; q)_{\infty} \sum_{k=0}^{\infty} \frac{q^k}{(q; q)_k} r_k^{2n} \\ &= \delta_{n,m} \frac{(q; q)_{\infty}}{(1-q)^n} \sum_{k=0}^{\infty} \frac{q^{(n+1)k}}{(q; q)_k}. \end{aligned}$$

□

Combining this result with our moment problem we obtain

$$[n]_q! = \mathcal{M}_q(E_q^{-1})(n+1) = \int_0^{1/(1-q)} t^n E_q^{-1}(qt) d_q t = \frac{\delta_{n,m}}{2\pi} \iint_{\mathbb{D}_{\frac{1}{1-q}}} z^n \overline{z}^m d\mu_q(z).$$

We observe that for $q \rightarrow 1$ we obtain $d\mu_q(z) = \frac{1}{2} e^{-|z|^2} dx dy$.

Also we get a convolution-type formula for our q -integral transform.

Lemma 3.5. *Given bounded functions $f_1, f_2 : [0, -1 + 1/(1-q)] \rightarrow \mathbb{R}$ it holds (pointwisely)*

$$(3.3) \quad \mathcal{M}_q(f_1)(z)\mathcal{M}_q(f_2)(z) = \left(\frac{1}{1-q}\right)^{z-1} \mathcal{M}_q(f_1 \circ f_2)(z),$$

where

$$(3.4) \quad f_1 \circ f_2 \left(q \frac{q^m}{1-q} \right) := \sum_{k=0}^m f_1 \left(\frac{q^{k+1}}{1-q} \right) f_2 \left(\frac{q^{m+1-k}}{1-q} \right).$$

Proof. From Definition 3.3 we have

$$\begin{aligned} \mathcal{M}_q(f_1)(z)\mathcal{M}_q(f_2)(z) &= \left(\sum_{k=0}^{\infty} q^k \left(\frac{q^k}{1-q} \right)^{z-1} f_1 \left(\frac{q^{k+1}}{1-q} \right) \right) \left(\sum_{n=0}^{\infty} q^n \left(\frac{q^n}{1-q} \right)^{z-1} f_2 \left(\frac{q^{n+1}}{1-q} \right) \right) \\ &= \sum_{k,n=0}^{\infty} q^{k+n} \left(\frac{q^{k+n}}{(1-q)^2} \right)^{z-1} f_1 \left(\frac{q^{k+1}}{1-q} \right) f_2 \left(\frac{q^{n+1}}{1-q} \right) \\ &= \left(\frac{1}{1-q} \right)^{z-1} \sum_{m=0}^{\infty} q^m \left(\frac{q^m}{1-q} \right)^{z-1} \underbrace{\left(\sum_{k=0}^m f_1 \left(\frac{q^{k+1}}{1-q} \right) f_2 \left(\frac{q^{m+1-k}}{1-q} \right) \right)}_{:= f_1 \circ f_2 \left(q \frac{q^m}{1-q} \right)}. \end{aligned}$$

□

For the multiplication operator M_z we have the following fact.

Proposition 3.6. M_z is bounded from $\mathbf{H}_{2,q}$ into itself with norm $\|M_z\| \leq \frac{1}{1-q}$.

Proof. This follows from

$$\frac{\frac{1}{1-q} - z\bar{w}}{\prod_{j=0}^{\infty} (1 - z\bar{w}(1-q)q^j)} = \frac{1}{1-q} \frac{1}{\prod_{j=1}^{\infty} (1 - z\bar{w}(1-q)q^j)}.$$

Since the kernel $\frac{1}{1-q} \frac{1}{\prod_{j=1}^{\infty} (1 - z\bar{w}(1-q)q^j)}$ is positive definite in $\mathbb{D}_{1/1-q}$ so is the kernel

$$\frac{\frac{1}{1-q} - z\bar{w}}{\prod_{j=0}^{\infty} (1 - z\bar{w}(1-q)q^j)},$$

and we conclude with the characterization of multipliers in a reproducing kernel Hilbert space. □

Lemma 3.7. (see e.g. [2, Exercise 4.2.25, pp. 165 and 185])

$$(3.5) \quad (R_q f)(z) = \lambda f(z) \iff f(z) = \frac{c}{\prod_{j=0}^{\infty} (1 - \lambda(1-q)zq^j)}.$$

Proposition 3.8. The q -exponential satisfy

$$(3.6) \quad (R_q E_q(\cdot \bar{w}))(z) = \bar{w} E_q(z \bar{w}).$$

Proof. We note that $E_q(qz\bar{w}) = (1 - z\bar{w}(1 - q))E_q(z\bar{w})$ and so

$$\begin{aligned} (R_q E_q(\cdot\bar{w}))(z) &= \frac{E_q(z\bar{w}) - E_q(qz\bar{w})}{(1 - q)z} \\ &= \frac{E_q(z\bar{w}) - (1 - z\bar{w}(1 - q))E_q(z\bar{w})}{(1 - q)z} \\ &= \bar{w}E_q(z\bar{w}). \end{aligned}$$

□

Theorem 3.9. *Let $q \in [0, 1)$. The only Hilbert space of functions which is analytic in a neighborhood of the origin and for which*

$$(3.7) \quad R_q^* = M_z$$

is $\mathbf{H}_{2,q}$ (up to a multiplicative factor for the inner product).

Proof. We have that $E_q(z\bar{w}) = K_q(z, w)$ is the reproducing for $\mathbf{H}_{2,q}$, i.e. $f(z) = \langle f, E_q(\cdot\bar{z}) \rangle_{\mathbf{H}_{2,q}}$. Using (3.6) we can write:

$$\begin{aligned} (R_q^* E_q(\cdot\bar{w}))(z) &= \langle R_q^* E_q(\cdot\bar{w}), E_q(\cdot\bar{z}) \rangle_{\mathbf{H}_{2,q}} \\ &= \langle E_q(\cdot\bar{w}), R_q E_q(\cdot\bar{z}) \rangle_{\mathbf{H}_{2,q}} \\ &= \langle E_q(\cdot\bar{w}), \bar{z} E_q(\cdot\bar{z}) \rangle_{\mathbf{H}_{2,q}} \\ &= z E_q(z\bar{w}). \end{aligned}$$

□

Proposition 3.10. *The space $\mathbf{H}_{2,q}$ is a de Branges-Rovnyak space.*

Proof. This follows from [3, Theorem 2.1 p. 51], since the sequence $[k]_q!$, for $k = 0, 1, \dots$ is an increasing sequence with initial term 1. □

We now compute the adjoint of R_q in $\mathbf{H}_{2,q}$. Since

$$(3.8) \quad \langle z^n, z^m \rangle_{\mathbf{H}_{2,q}} = [n]_q! \delta_{n,m},$$

we have

$$\begin{aligned} \langle z^n, R_q z^m \rangle_{\mathbf{H}_{2,q}} &= \left\langle z^n, \frac{z^m - q^m z^m}{(1 - q)z} \right\rangle_{\mathbf{H}_{2,q}} = (1 + q + \dots + q^{m-1}) \langle z^n, z^{m-1} \rangle_{\mathbf{H}_{2,q}} \\ &= (1 + q + \dots + q^n) [n]_q! = [n + 1]_q! \\ &= \langle z^{n+1}, z^m \rangle_{\mathbf{H}_{2,q}} =: \langle R_q^* z^n, z^m \rangle_{\mathbf{H}_{2,q}}. \end{aligned}$$

Therefore, we obtain $R_q^* = M_z$.

In the case $q = 1$ the Fock space can be characterized (up to a multiplicative positive factor in the inner product) as the only Hilbert space of power series converging in a convex neighborhood of the origin and such that

$$(3.9) \quad (R_0^* f)(z) = \int_{[0,z]} f(s) ds,$$

that is, R_0^* coincides with the integration operator. It is therefore natural to try and define the integral in $\mathbf{H}_{2,q}$ by R_q^* for $q \in (0, 1)$.

Lemma 3.11. *The operator R_0 is bounded in $\mathbf{H}_{2,q}$ and it holds that (with $e_k(z) = z^k$)*

$$(3.10) \quad R_0^* e_k = \frac{e_{k+1}}{1 + q + \dots + q^k}, \quad k = 0, 1, \dots$$

Proof. We have for $k \geq 1$ and $\ell \geq 0$

$$\begin{aligned} \langle R_0 e_k, e_\ell \rangle_{\mathbf{H}_{2,q}} &= \langle e_{k-1}, e_\ell \rangle_{\mathbf{H}_{2,q}} \\ &= \delta_{k-1,\ell} [\ell]_q! \\ &= \delta_{k-1,\ell} \langle e_k, e_k \rangle_{\mathbf{H}_{2,q}} \frac{[\ell]_q!}{[k]_q!} \\ &= \delta_{k-1,\ell} \langle e_k, e_k \rangle_{\mathbf{H}_{2,q}} \frac{1}{1 + q + \dots + q^\ell} \\ &= \langle e_k, R_0^* e_\ell \rangle_{\mathbf{H}_{2,q}}, \end{aligned}$$

with

$$(3.11) \quad R_0^* e_\ell = \frac{e_{\ell+1}}{1 + q + \dots + q^\ell}.$$

□

Consider the q -Jackson integral

$$\int_0^a f(x) d_q x := (1 - q)a \sum_{k=0}^{\infty} q^k f(q^k a),$$

which is said to converge provided that the sum on the right-hand-side converges absolutely.

Lemma 3.12.

$$(3.12) \quad \int_0^z x^\ell d_q x = z^{\ell+1} \frac{1}{1 + q + \dots + q^\ell}.$$

Proof. By definition we have

$$\begin{aligned} \int_0^z x^\ell d_q x &= z(1 - q) \sum_{k=0}^{\infty} q^k (q^k z)^\ell = z^{\ell+1} (1 - q) \left(\sum_{k=0}^{\infty} (q^{1+\ell})^k \right) \\ &= z^{\ell+1} (1 - q) \frac{1}{1 - q^{\ell+1}} = z^{\ell+1} \frac{1}{1 + q + \dots + q^\ell}. \end{aligned}$$

□

It is well know that

$$(3.13) \quad \partial^* = M_z$$

in the Fock space, and that in fact the Fock space is characterized (up to a positive multiplicative constant in the inner product) by this equality; see [10]. In [8] it is proved that in the Hardy space we have

$$(3.14) \quad \partial^* = M_z \partial M_z,$$

and that the above equality does characterize the Hardy space (as usual, up to a positive multiplicative constant in the inner product). We now prove a formula which is valid for $q \in [0, 1]$ and englobes the two above formulas.

Theorem 3.13. *Let $q \in [0, 1]$. Then in $\mathbf{H}_{2,q}$ it holds that*

$$(3.15) \quad \partial^* = M_z \partial R_0^*$$

and this equality characterizes the space $\mathbf{H}_{2,q}$ up to a positive multiplicative constant in the inner product.

When $q = 0$ (Hardy space) we have R_0^* that $R_0^* = M_z$ and so (3.15) reduces to

$$M_z \partial M_z,$$

i.e. (3.14). When $q = 1$ (Fock space), we have $R_0^* = I$ (the integration operator) and $\partial I e_k = e_k$, $k = 0, 1, \dots$. We thus get back (3.13).

Proof of Theorem 3.13. Let $k \in \mathbb{N}_0$. Let us set *a priori* $\partial^* e_k = a_{k,q} e_{k+1}$ for some $a_{k,q} \in \mathbb{C}$. We have on the one hand

$$\begin{aligned} \langle \partial^* e_k, e_{k+1} \rangle_{\mathbf{H}_{2,q}} &= \langle e_k, \partial e_{k+1} \rangle_{\mathbf{H}_{2,q}} \\ &= (k+1) \langle e_k, e_k \rangle_{\mathbf{H}_{2,q}} \\ &= (k+1) [k]_q! \end{aligned}$$

and on the other hand, with $\partial^* e_k = a_{k,q} e_{k+1}$ we have

$$\begin{aligned} \langle \partial^* e_k, e_{k+1} \rangle_{\mathbf{H}_{2,q}} &= a_{k,q} \langle e_{k+1}, e_{k+1} \rangle_{\mathbf{H}_{2,q}} \\ &= a_{k,q} [k+1]_q!. \end{aligned}$$

Thus

$$a_{k,q} [k+1]_q! = (k+1) [k]_q!$$

from which we get

$$(3.16) \quad a_{k,q} = \frac{k+1}{1+q+\dots+q^k}.$$

In view of (3.11), we can write

$$\begin{aligned} \partial^* e_k &= \frac{(k+1) M_z e_k}{1+q+\dots+q^k} \\ &= (k+1) R_0^* e_k \\ &= M_z \partial R_0^* e_k \end{aligned}$$

since

$$M_z \partial e_{k+1} = (k+1) e_{k+1}, \quad k = 0, 1, \dots$$

□

Note that in (3.16), we set $a_{k,0} = k+1$ and $a_{k,1} = 1$, as it should be.

Theorem 3.14. *We have*

$$(3.17) \quad M_z^* = R_q M_z R_0.$$

Proof. For $m = 1, 2, \dots$ we get

$$\langle z^n, R_q M_z R_0 z^m \rangle_{\mathbf{H}_{2,q}} = \langle z^n, R_q M_z z^{m-1} \rangle_{\mathbf{H}_{2,q}} = \langle z^n, R_q z^m \rangle_{\mathbf{H}_{2,q}}.$$

By Proposition 3.10 we obtain

$$\langle z^n, R_q M_z R_0 z^m \rangle_{\mathbf{H}_{2,q}} = \langle z^n, R_q z^m \rangle_{\mathbf{H}_{2,q}} = \langle M_z z^n, z^m \rangle_{\mathbf{H}_{2,q}}.$$

We conclude our proof with the observation that $0 = \langle z^n, R_q M_z R_0 z^0 \rangle_{\mathbf{H}_{2,q}} = \langle M_z z^n, z^0 \rangle_{\mathbf{H}_{2,q}}$. \square

4. THE SPACE $\mathcal{F}_{2,q}$

The space $\mathcal{F}_{2,q}$ appeared in [5] motivated by a study of discrete analytic functions.

Definition 4.1. *Consider the reproducing kernel*

$$K_{2,q}(z, w) = \sum_{n=0}^{\infty} \frac{z^n \bar{w}^n}{([n_q]!)^2}.$$

Then the corresponding reproducing kernel Hilbert space $\mathcal{F}_{2,q}$ is the space of all functions $f(z) = \sum_{n=0}^{\infty} f_n z^n$ such that $\sum_{n=0}^{\infty} |f_n|^2 ([n_q]!)^2 < \infty$.

In this way, we have $K_q =: K_{1,q}$ and $K_{2,q}$ as the reproducing kernels of $\mathbf{H}_{2,q}$ and $\mathcal{F}_{2,q}$, respectively. As both kernels are positive definite and the same holds for its difference $K_{1,q} - K_{2,q}$ we get that $\mathcal{F}_{2,q}$ is contractively included in $\mathbf{H}_{2,q}$ (see [1, 9]).

Remark 4.2. We observe that for $f_1(z) = f_2(z) = E_q^{-1}(z)$ we have

$$\begin{aligned} [\mathcal{M}_q(E_q^{-1})(n+1)]^2 &= \left(\frac{1}{1-q}\right)^n \sum_{m=0}^{\infty} q^m \left(\frac{q^m}{1-q}\right)^{z-1} \left[\sum_{k=0}^m E_q^{-1}\left(\frac{q^{k+1}}{1-q}\right) E_q^{-1}\left(\frac{q^{m+1-k}}{1-q}\right) \right] \\ ([n]_q!)^2 &= \left(\frac{1}{1-q}\right)^n \sum_{m=0}^{\infty} q^m \left(\frac{q^m}{1-q}\right)^n \sum_{k=0}^m (q^{k+1}; q)_{\infty} (q^{m+1-k}; q)_{\infty}. \end{aligned}$$

Hence, we get as density $\omega_{2,q}$ of our q -Fock space $\mathcal{F}_{2,q}$

$$(4.1) \quad \omega_{2,q}(|z|) := \left(\frac{1}{1-q}\right)^{|z|^2-1} (E_q^{-1} \circ E_q^{-1})(|z|^2),$$

and satisfying to

$$(4.2) \quad \mathcal{M}_q(\omega_{2,q})(n+1) = \left(\frac{1}{1-q}\right)^n \mathcal{M}_q(E_q^{-1} \circ E_q^{-1})(n+1) = ([n]_q!)^2.$$

Now, we can define $T_q : \mathbf{H}_2 \mapsto \mathcal{F}_{2,q}$ given as $z^n \rightarrow \frac{z^n}{[n]_q!}$.

Lemma 4.3. *In \mathbf{H}_2 it holds:*

$$(4.3) \quad R_q T_q = T_q R_0.$$

Proof. The case of $n = 0$ is immediate. For $n = 1, 2, \dots$ we have

$$\begin{aligned} R_q T_q z^n &= R_q \left(\frac{z^n}{[n]_q!} \right) = \frac{1}{[n]_q!} (1 + q + \dots + q^{n-1}) z^{n-1} \\ &= \frac{1}{[n-1]_q!} z^{n-1} = T_q z^{n-1} = T_q R_0 z^n. \end{aligned}$$

\square

Theorem 4.4. *The map T_q is an isometry from \mathbf{H}_2 onto $\mathcal{F}_{2,q}$.*

Proof. We have $\langle e_n, e_m \rangle_{\mathcal{F}_{2,q}} = ([n]_q!)^2 \delta_{n,m}$. Hence, we get

$$\begin{aligned} \langle T_q e_n, T_q e_m \rangle_{\mathcal{F}_{2,q}} &= \frac{1}{[n]_q!} \frac{1}{[m]_q!} \langle e_n, e_m \rangle_{\mathcal{F}_{2,q}} \\ &= \delta_{n,m} \frac{([n]_q!)^2}{([n]_q!)^2} \\ &= \delta_{n,m} \\ &= \langle e_n, e_m \rangle_{\mathbf{H}_2}. \end{aligned}$$

□

Theorem 4.5. *In $\mathcal{F}_{2,q}$ it holds that*

$$(4.4) \quad R_q^* e_n = \frac{e_{n+1}}{[n]_q}, \quad n = 0, 1, \dots$$

and

$$(4.5) \quad \mathcal{I} - R_q^* R_q = C^* C,$$

and this structural identity characterizes the space $\mathcal{F}_{2,q}$ up to a multiplicative factor.

Proof. To prove (4.4) we write

$$\begin{aligned} \langle R_q^* e_n, e_m \rangle_{\mathcal{F}_{q,2}} &= \langle e_n, R_q e_m \rangle_{\mathcal{F}_{2,q}} \\ &= [m]_q \langle e_n, e_{m-1} \rangle_{\mathcal{F}_{q,2}} \\ &= \delta_{m-1,n} ([n]_q!)^2 [m]_q. \end{aligned}$$

On the other hand we show that one can assume that $R_q^* e_n = \alpha_n e_{n+1}$; we have

$$\begin{aligned} \langle R_q^* e_n, e_m \rangle &= \alpha_n \langle e_{n+1}, e_m \rangle_{\mathcal{F}_{2,q}} \\ &= \alpha_n \delta_{n+1,m} ([n+1]_q!)^2. \end{aligned}$$

Comparing these equalities we obtain

$$\alpha_n ([n+1]_q!)^2 = ([n]_q!)^2 [n]_q,$$

so that $\alpha_n = \frac{1}{[n]_q}$. It follows that

$$R_q^* R_q e_n = \begin{cases} 0, & n = 0, \\ e_n, & n = 1, 2, \dots \end{cases}$$

and hence the result. □

From the previous computations we also have:

Proposition 4.6. *R_0^* is an isometry in $\mathcal{F}_{2,q}$.*

Proof. This is a direct consequence of the fact that

$$R_0 R_0^* e_n = e_n,$$

for all $n \in \mathbb{N}_0$. □

From Lemma 3.12 we have:

Proposition 4.7. *In $\mathcal{F}_{2,q}$, it holds*

$$R_q^* = I,$$

where I is the integration operator.

We now use well a known method in characteristic function theory (see e.g. [4] in the case of Pontryagin spaces) and rewrite (4.5) as

$$\begin{pmatrix} R_q \\ C \end{pmatrix}^* \begin{pmatrix} R_q \\ C \end{pmatrix} = \mathcal{I}.$$

The operator

$$\begin{pmatrix} \mathcal{I} & 0 \\ 0 & 1 \end{pmatrix} - \begin{pmatrix} R_q \\ C \end{pmatrix} \begin{pmatrix} R_q \\ C \end{pmatrix}^*$$

is therefore positive and for instance using its square root, one can find a Hilbert space $\tilde{\mathcal{H}}$ and operators B and D ,

$$\begin{pmatrix} B \\ D \end{pmatrix} : \tilde{\mathcal{H}} \longrightarrow \mathcal{F}_{2,q} \oplus \mathbb{C},$$

such that

$$\begin{pmatrix} \mathcal{I} & 0 \\ 0 & 1 \end{pmatrix} - \begin{pmatrix} R_q \\ C \end{pmatrix} \begin{pmatrix} R_q \\ C \end{pmatrix}^* = \begin{pmatrix} B \\ D \end{pmatrix} \begin{pmatrix} B \\ D \end{pmatrix}^*.$$

The operator matrix

$$(4.6) \quad \begin{pmatrix} R_q & B \\ C & D \end{pmatrix}$$

is co-isometric. We set

$$(4.7) \quad S_q(z) = D + zC(\mathcal{I} - zR_q)^{-1}B.$$

We now look into the properties of the matrix (4.6). We observe that

$$\begin{pmatrix} \mathcal{I} & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} R_q & B \\ C & D \end{pmatrix} \begin{pmatrix} R_q & B \\ C & D \end{pmatrix}^* = \begin{pmatrix} R_q R_q^* + BB^* & R_q C^* + BD^* \\ CR_q^* + DB^* & CC^* + DD^* \end{pmatrix},$$

so that we get

$$DD^* = 1 - CC^*, \quad BB^* = \mathcal{I} - R_q R_q^*, \quad BD^* = -R_q C^*.$$

Hence, from (4.7) we get

$$\begin{aligned} S_q(z)[S_q(w)]^* &= [D\mathcal{I} + zC(\mathcal{I} - zR_q)^{-1}B][D^*\mathcal{I} + \bar{w}B^*(\mathcal{I} - \bar{w}R_q^*)^{-1}C^*] \\ &= DD^*\mathcal{I} + zC(\mathcal{I} - zR_q)^{-1}BD^* + \bar{w}DB^*(\mathcal{I} - \bar{w}R_q^*)^{-1}C^* + z\bar{w}C(\mathcal{I} - zR_q)^{-1}BB^*(\mathcal{I} - \bar{w}R_q^*)^{-1}C^* \\ &= (1 - CC^*)\mathcal{I} + zC(\mathcal{I} - zR_q)^{-1}BD^* + \bar{w}DB^*(\mathcal{I} - \bar{w}R_q^*)^{-1}C^* + z\bar{w}C(\mathcal{I} - zR_q)^{-1}BB^*(\mathcal{I} - \bar{w}R_q^*)^{-1}C^* \end{aligned}$$

so that

$$\begin{aligned} &\mathcal{I} - S_q(z)[S_q(w)]^* \\ &= CC^*\mathcal{I} - zC(\mathcal{I} - zR_q)^{-1}BD^* - \bar{w}DB^*(\mathcal{I} - \bar{w}R_q^*)^{-1}C^* - z\bar{w}C(\mathcal{I} - zR_q)^{-1}BB^*(\mathcal{I} - \bar{w}R_q^*)^{-1}C^* \\ &= CC^*\mathcal{I} + zC(\mathcal{I} - zR_q)^{-1}R_q C^* + \bar{w}CR_q^*(\mathcal{I} - \bar{w}R_q^*)^{-1}C^* - z\bar{w}C(\mathcal{I} - zR_q)^{-1}(\mathcal{I} - R_q R_q^*)(\mathcal{I} - \bar{w}R_q^*)^{-1}C^* \\ &= C(\mathcal{I} - zR_q)^{-1} \underbrace{\left[(\mathcal{I} - zR_q)(1 - \bar{w}R_q^*) + zR_q(\mathcal{I} - \bar{w}R_q^*) + \bar{w}(\mathcal{I} - zR_q)R_q^* - z\bar{w}(\mathcal{I} - R_q R_q^*)(\mathcal{I} - \bar{w}R_q^*) \right]}_{(A)} (\mathcal{I} - \bar{w}R_q^*)^{-1}C^*. \end{aligned}$$

Easy calculations give now

$$\begin{aligned}
(A) &= (\mathcal{I} - zR_q)(\mathcal{I} - \bar{w}R_q^*) + zR_q(\mathcal{I} - \bar{w}R_q^*) + \bar{w}(\mathcal{I} - zR_q)R_q^* - z\bar{w}(\mathcal{I} - R_qR_q^*) \\
&= \mathcal{I} - zR_q - \bar{w}R_q^* + z\bar{w}R_qR_q^* + zR_q - z\bar{w}R_qR_q^* + \bar{w}R_q^* - z\bar{w}R_qR_q^* - z\bar{w}\mathcal{I} + z\bar{w}R_qR_q^* \\
&= (1 - z\bar{w})\mathcal{I},
\end{aligned}$$

Hence, it holds that

$$(4.8) \quad \frac{\mathcal{I} - S_q(z)S_q(w)^*}{1 - z\bar{w}} = C(\mathcal{I} - zR_q)^{-1}[(\mathcal{I} - wR_q)^*]^{-1}C^*, \quad z, w \in \mathbb{D}.$$

The operator S_q bears various names in operator theory; it is the characteristic operator function, or the transfer function, or the scattering function, associated to the operator matrix (4.6). From (4.8) one sees that S_q is analytic and contractive in the open unit disk, i.e. is a Schur function.

When $q = 0$ we have for $f(z) = \sum_{n=0}^{\infty} c_n z^n$ that

$$C(\mathcal{I} - zR_0)^{-1}f = f(z), \quad z \in \mathbb{D}.$$

Here, for $0 < q \leq 1$ we define

$$f_q(z) = C(\mathcal{I} - zR_q)^{-1}f, \quad z \in \mathbb{D}.$$

As $CR_q^n f = [n]_q! c_n$ we get that the coefficients

$$(4.9) \quad c_n = \frac{CR_q^n f}{[n]_q!}$$

are independent of q and one has $f_q(z) = f(z)$, that is, we obtain $f(z) = C(\mathcal{I} - zR_q)^{-1}f$ for all $z \in \mathbb{D}$.

REFERENCES

- [1] D. Alpay. *The Schur algorithm, reproducing kernel spaces and system theory*. American Mathematical Society, Providence, RI, 2001. Translated from the 1998 French original by Stephen S. Wilson, Panoramas et Synthèses.
- [2] D. Alpay. *A complex analysis problem book*. Birkhäuser/Springer, Cham, 2016. Second edition.
- [3] D. Alpay, F. Colombo, and I. Sabadini. The Fock space as a De Branges-Rovnyak space. *Integral Equations Operator Theory*, 91(6):Paper No. 51, 12, 2019.
- [4] D. Alpay, A. Dijksma, J. Rovnyak, and H. de Snoo. *Schur functions, operator colligations, and reproducing kernel Pontryagin spaces*, volume 96 of *Operator theory: Advances and Applications*. Birkhäuser Verlag, Basel, 1997.
- [5] D. Alpay, P. Jorgensen, R. Seager, and D. Volok. On discrete analytic functions: Products, rational functions and reproducing kernels. *Journal of Applied Mathematics and Computing*, 41:393–426, 2013.
- [6] D. Alpay and M. Porat. Generalized Fock spaces and the Stirling numbers. *J. Math. Phys.*, 59(6):063509, 12, 2018.
- [7] Daniel Alpay, Paula Cerejeiras, Uwe Kaehler, and Trevor Kling. Commutators on Fock spaces. *J. Math. Phys.*, 64:042102–21pages, 2023.
- [8] N. Alpay. A new characterization of the Hardy space and of other spaces of analytic functions. *ArXiv (2020)*; To appear in *İstanbul Univ., Sci. Fac., J. Math. Phys. Astron.*, pages 1–10, 2023.
- [9] N. Aronszajn. La théorie générale des noyaux reproduisants et ses applications. *Math. Proc. Cambridge Phil. Soc.*, 39:133–153, 1944.

- [10] V. Bargmann. On a Hilbert space of analytic functions and an associated integral transform. *Comm. Pure Appl. Math.*, 14:187–214, 1961.
- [11] V. Bargmann. Remarks on a Hilbert space of analytic functions. *Proceedings of the National Academy of Arts*, 48:199–204, 1962.
- [12] Yue Cai, Richard Ehrenborg, and Margaret Readdy. q -Stirling identities revisited. *Electron. J. Comb.*, 25:1–37, 2018.
- [13] Yue Cai and Margaret Readdy. q -Stirling numbers: a new view. *Adv. Appl. Math.*, 86:50–80, 2017.
- [14] P.L. Duren. *Theory of H^p spaces*. Academic press, New York, 1970.
- [15] K. Hoffman. *Banach spaces of analytic functions*. Dover Publications Inc., New York, 1988. Reprint of the 1962 original.
- [16] F. H. Jackson. On q -definite integrals. *Quart. J.*, 41:193–203, 1910.
- [17] Victor G. Kac and Pokman Cheung. *Quantum calculus*. Universitext (UTX). Springer, New York, 2001.
- [18] W. Rudin. *Real and complex analysis*. McGraw-Hill Book Co., New York, third edition, 1987.
- [19] Hans van Leeuwen and Hans Maassen. A q deformation of the Gauss distribution. *J. Math. Phys.*, 36:4743–4756, 1995.
- [20] Kehe Zhu. *Analysis on Fock spaces*, volume 263 of *Graduate Texts in Mathematics (GTM)*. Springer, New York, NY, 2012.

(DA) FACULTY OF MATHEMATICS, PHYSICS, AND COMPUTATION, SCHMID COLLEGE OF SCIENCE AND TECHNOLOGY, CHAPMAN UNIVERSITY, ONE UNIVERSITY DRIVE ORANGE, CALIFORNIA 92866, USA
Email address: `alpay@chapman.edu`

(PC) CENTER FOR RESEARCH AND DEVELOPMENT IN MATHEMATICS AND APPLICATIONS, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF AVEIRO, CAMPUS UNIVERSITÁRIO DE SANTIAGO, 3810-193 AVEIRO, PORTUGAL
Email address: `pceres@ua.pt`

(UK) CENTER FOR RESEARCH AND DEVELOPMENT IN MATHEMATICS AND APPLICATIONS, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF AVEIRO, CAMPUS UNIVERSITÁRIO DE SANTIAGO, 3810-193 AVEIRO, PORTUGAL
Email address: `ukaehler@ua.pt`

(BS) UNIVERSITY OF OSTRAVA, DEPARTMENT OF MATHEMATICS, 30.DUBNA 22, 70200 OSTRAVA, CZECH REPUBLIC
Email address: `baruch.schneider@osu.cz`